

**Ph.D. Qualifying Exam**  
**Algebra: 901-902. June 1, 1999**

Do a total of four problems, one from each section. In sections III and IV, all rings are assumed to be commutative with identity.

SECTION I: GROUPS

1. Classify up to isomorphism all groups of order 28.
2. Prove that every group of order  $5 \cdot 19 \cdot 29$  is cyclic.

SECTION II: FIELDS

3. Let  $E/K$  be an algebraic extension and  $\sigma : E \rightarrow E$  a field homomorphism fixing  $K$ . Prove that  $\sigma$  is an isomorphism.
4. Let  $E$  be the splitting field of  $x^6 + x^3 + 1$  over  $\mathbb{Q}$ .
  - (a) Find the Galois group of  $E/\mathbb{Q}$ .
  - (b) Find field generators for all of the intermediate fields of  $E/\mathbb{Q}$ .
5. Let  $E/K$  be a separable algebraic extension such that every nonconstant polynomial in  $K[x]$  has a root in  $E$ . Prove that  $E$  is an algebraic closure of  $K$ . (Hint: use the primitive element theorem.)

SECTION III: GENERAL RINGS AND MODULES

6. Let  $I$  be an ideal of  $R$  and  $M$  a finitely generated  $R$ -module such that  $IM = M$ . Prove that there exists  $s \in I$  such that  $(1 - s)M = 0$ .
7. Let  $R$  be a ring with a unique maximal ideal. Prove that every finitely generated projective  $R$ -module is free.
8. Let  $(\#)$  denote the exact sequence of  $R$ -modules  $0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} N \rightarrow 0$ .
  - (a) Prove that  $(\#)$  splits if and only if the natural map  $\text{Hom}_R(N, M) \xrightarrow{g^*} \text{Hom}_R(N, N)$  is surjective.
  - (b) Suppose  $N$  is finitely presented. Prove that  $(\#)$  splits if and only if for all  $p \in \text{Spec } R$ ,  $0 \rightarrow L_p \rightarrow M_p \rightarrow N_p \rightarrow 0$  splits.

SECTION IV: NOETHERIAN RINGS AND MODULES

9. Let  $R$  be a Dedekind domain and  $M$  a finitely generated  $R$ -module. Prove that  $M$  is projective if and only if it is torsion-free.
10. Let  $R \subseteq S$  be Noetherian rings such that  $S$  is integral over  $R$ . Suppose  $R$  is local. Prove that  $S$  has only finitely many maximal ideals.
11. Let  $R$  be a Noetherian ring and  $I$  and  $J$  two ideals of  $R$ . Prove that  $I = J$  if and only if  $I_p = J_p$  for all  $p \in \text{Ass}_R(R/I) \cup \text{Ass}_R(R/J)$ .