

Commutative algebras of Toeplitz operators in action

Nikolai Vasilevski

Department of Mathematics
CINVESTAV del I.P.N., Mexico City, Mexico

e-mail: nvasilev@math.cinvestav.mx

- Commutative algebras of Toeplitz operators on the unit disk.
- Fine structure of the algebra of Toeplitz operators with PC -symbols.
- From the unit disk to the unit ball.

- Commutative algebras of Toeplitz operators on the unit disk.
- Fine structure of the algebra of Toeplitz operators with PC -symbols.
- From the unit disk to the unit ball.

Toeplitz operators

The Toeplitz operator was originally defined in terms of the so-called Toeplitz matrix

$$A = \begin{pmatrix} a_0 & a_{-1} & a_{-2} & \dots \\ a_1 & a_0 & a_{-1} & \dots \\ a_2 & a_1 & a_0 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix},$$

where $a_n \in \mathbb{C}$, $n \in \mathbb{Z}$.

Theorem (O. Toeplitz, 1911)

Matrix A defines a bounded operator on $l_2 = l_2(\mathbb{Z}_+)$, where $\mathbb{Z}_+ = \{0\} \cup \mathbb{N}$, if and only if the numbers $\{a_n\}$ are the Fourier coefficients of a function $a \in L_\infty(S^1)$, where S^1 is the unit circle.

Hardy space version

The (discrete) Fourier transform \mathcal{F} is a unitary operator which maps $L_2(S^1)$ onto $l_2(\mathbb{Z})$ and the Hardy space $H_+^2(S^1)$ onto $l_2(\mathbb{Z}_+)$. Then for the operator \mathbb{A} , defined by the matrix A we have

$$\mathcal{F}^{-1} \mathbb{A} \mathcal{F} = T_a : H_+^2(S^1) \longrightarrow H_+^2(S^1).$$

The operator T_a acts on the Hardy space $H_+^2(S^1)$ by the rule

$$T_a : f(t) \in H_+^2(S^1) \longmapsto (P_+ a f)(t) \in H_+^2(S^1),$$

where $P_+ : L_2(S^1) \longrightarrow H_+^2(S^1)$ is the Szegő orthogonal projection, and the Fourier coefficients of the function a are given by the sequence $\{a_n\}$.

Let H be a Hilbert space, H_0 be its subspace.
Let $P_0 : H \mapsto H_0$ be the orthogonal projection,
and let A be a bounded linear operator on H .

The Toeplitz operator with symbol A

$$T_A : x \in H_0 \mapsto P_0(Ax) \in H_0$$

is the **compression** of A (in our case of a multiplication operator)
onto the subspace H_0 , **representing thus an important model case**
in operator theory.

Bergman space version

Consider now $L_2(\mathbb{D})$, where \mathbb{D} is the unit disk in \mathbb{C} .

The Bergman space $\mathcal{A}^2(\mathbb{D})$ is the subspace of $L_2(\mathbb{D})$ consisting of functions analytic in \mathbb{D} .

The Bergman orthogonal projection $B_{\mathbb{D}}$ of $L_2(\mathbb{D})$ onto $\mathcal{A}^2(\mathbb{D})$ has the form

$$(B_{\mathbb{D}}\varphi)(z) = \frac{1}{\pi} \int_{\mathbb{D}} \frac{\varphi(\zeta) d\mu(\zeta)}{(1 - z\bar{\zeta})^2},$$

The Toeplitz operator T_a with symbol $a = a(z)$ acts as follows

$$T_a : \varphi(z) \in \mathcal{A}^2(\mathbb{D}) \longmapsto (B_{\mathbb{D}} a\varphi)(z) \in \mathcal{A}^2(\mathbb{D}).$$

Consider the unit disk \mathbb{D} endowed with the hyperbolic metric

$$g = ds^2 = \frac{1}{\pi} \frac{dx^2 + dy^2}{(1 - (x^2 + y^2))^2}.$$

A **geodesic** in \mathbb{D} is (a part of) an Euclidean circle or a straight line orthogonal to the boundary $S^1 = \partial\mathbb{D}$.




Each pair of geodesics, say L_1 and L_2 , lie in a geometrically defined object, one-parameter family \mathcal{P} of geodesics, which is called the **pencil** determined by L_1 and L_2 .

Each pencil has an associated family \mathcal{C} of lines, called **cycles**, the orthogonal trajectories to geodesics forming the pencil.

Pencils of hyperbolic geodesics

There are **three** types of pencils of hyperbolic geodesics:


- **parabolic,**
- **elliptic,**
- **hyperbolic.**



Möbius transformations

Each Möbius transformation $g \in \text{Möb}(\mathbb{D})$ is a **movement of the hyperbolic plane**, determines a certain pencil of geodesics \mathcal{P} , and its action is as follows:

each geodesic L from the pencil \mathcal{P} , determined by g , moves along the cycles in \mathcal{C} to the geodesic $g(L) \in \mathcal{P}$, while each cycle in \mathcal{C} is invariant under the action of g








Theorem

*Given a pencil \mathcal{P} of geodesics, consider the set of symbols which are **constant** on corresponding cycles. The C^* -algebra generated by Toeplitz operators with such symbols is **commutative**.*

That is, each pencil of geodesics generates a commutative C^* -algebra of Toeplitz operators.

Model cases



Hyperbolic case

Consider the upper half-plane Π , the space $L_2(\Pi)$, and its Bergman subspace $\mathcal{A}^2(\Pi)$. We construct the operator

$$R : L_2(\Pi) \longrightarrow L_2(\mathbb{R}),$$

whose restriction onto the Bergman space

$$R|_{\mathcal{A}^2(\Pi)} : \mathcal{A}^2(\Pi) \longrightarrow L_2(\mathbb{R})$$

is an **isometric isomorphism**.

The adjoint operator

$$R^* : L_2(\mathbb{R}) \longrightarrow \mathcal{A}^2(\Pi) \subset L_2(\Pi)$$

is an **isometric isomorphism** of $L_2(\mathbb{R})$ onto $\mathcal{A}^2(\Pi)$.

Moreover we have

$$\begin{aligned} R R^* &= I : L_2(\mathbb{R}) \longrightarrow L_2(\mathbb{R}), \\ R^* R &= B_{\Pi} : L_2(\Pi) \longrightarrow \mathcal{A}^2(\Pi). \end{aligned}$$

Theorem

Let $a = a(\theta) \in L_\infty(\Pi)$ be a *homogeneous of order zero function*,
(*a functions depending only on the polar angle θ*).

Then the *Toeplitz operator* T_a acting on $\mathcal{A}^2(\Pi)$ is *unitary equivalent* to the *multiplication operator* $\gamma_a I = R T_a R^*$,
acting on $L_2(\mathbb{R})$.

The function $\gamma_a(\lambda)$ is given by

$$\gamma_a(\lambda) = \frac{2\lambda}{1 - e^{-2\pi\lambda}} \int_0^\pi a(\theta) e^{-2\lambda\theta} d\theta, \quad \lambda \in \mathbb{R}.$$

Symplectic manifold

We consider the pair (\mathbb{D}, ω) , where \mathbb{D} is the unit disk and

$$\omega = \frac{1}{\pi} \frac{dx \wedge dy}{(1 - (x^2 + y^2))^2} = \frac{1}{2\pi i} \frac{d\bar{z} \wedge dz}{(1 - |z|^2)^2}.$$

Poisson brackets:

$$\begin{aligned}\{a, b\} &= \pi(1 - (x^2 + y^2))^2 \left(\frac{\partial a}{\partial y} \frac{\partial b}{\partial x} - \frac{\partial a}{\partial x} \frac{\partial b}{\partial y} \right) \\ &= 2\pi i(1 - z\bar{z})^2 \left(\frac{\partial a}{\partial z} \frac{\partial b}{\partial \bar{z}} - \frac{\partial a}{\partial \bar{z}} \frac{\partial b}{\partial z} \right).\end{aligned}$$

Laplace-Beltrami operator:

$$\begin{aligned}\Delta &= \pi(1 - (x^2 + y^2))^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) \\ &= 4\pi(1 - z\bar{z})^2 \frac{\partial^2}{\partial z \partial \bar{z}}.\end{aligned}$$

Weighted Bergman spaces

Introduce weighted Bergman spaces $\mathcal{A}_h^2(\mathbb{D})$ with the scalar product

$$(\varphi, \psi) = \left(\frac{1}{h} - 1\right) \int_{\mathbb{D}} \varphi(z) \overline{\psi(z)} (1 - z\bar{z})^{\frac{1}{h}} \omega(z).$$

The weighted Bergman projection has the form

$$(B_{\mathbb{D}, h}\varphi)(z) = \left(\frac{1}{h} - 1\right) \int_{\mathbb{D}} \varphi(\zeta) \left(\frac{1 - \zeta\bar{\zeta}}{1 - z\bar{\zeta}}\right)^{\frac{1}{h}} \omega(\zeta).$$

Let $E = (0, \frac{1}{2\pi})$, for each $\hbar = \frac{h}{2\pi} \in E$, and consequently $h \in (0, 1)$, introduce the Hilbert space H_{\hbar} as the weighted Bergman space $\mathcal{A}_{\hbar}^2(\mathbb{D})$.

For each function $a = a(z) \in C^\infty(\mathbb{D})$ consider the family of Toeplitz operators $T_a^{(h)}$ with (anti-Wick) symbol a acting on $\mathcal{A}_h^2(\mathbb{D})$, for $h \in (0, 1)$, and denote by \mathcal{T}_h the $*$ -algebra generated by Toeplitz operators $T_a^{(h)}$ with symbols $a \in C^\infty(\mathbb{D})$.

The **Wick symbols** of the Toeplitz operator $T_a^{(h)}$ has the form

$$\tilde{a}_h(z, \bar{z}) = \left(\frac{1}{h} - 1\right) \int_{\mathbb{D}} a(\zeta) \left(\frac{(1 - |z|^2)(1 - |\zeta|^2)}{(1 - z\bar{\zeta})(1 - \zeta\bar{z})} \right)^{\frac{1}{h}} \omega(\zeta).$$

Star product

For each $h \in (0, 1)$ define the function algebra

$$\tilde{\mathcal{A}}_h = \{\tilde{a}_h(z, \bar{z}) : a \in C^\infty(\mathbb{D})\}$$

with point wise linear operations, and with the multiplication law defined by the product of Toeplitz operators:

$$\tilde{a}_h \star \tilde{b}_h = \left(\frac{1}{h} - 1\right) \int_{\mathbb{D}} \tilde{a}_h(z, \bar{\zeta}) \tilde{b}_h(\zeta, \bar{z}) \left(\frac{(1 - |z|^2)(1 - |\zeta|^2)}{(1 - z\bar{\zeta})(1 - \zeta\bar{z})}\right)^{\frac{1}{h}} \omega.$$

Correspondence principle

The correspondence principle is given by

$$\begin{aligned}\tilde{a}_h(z, \bar{z}) &= a(z, \bar{z}) + O(\hbar), \\ (\tilde{a}_h \star \tilde{b}_h - \tilde{b}_h \star \tilde{a}_h)(z, \bar{z}) &= i\hbar \{a, b\} + O(\hbar^2).\end{aligned}$$

Three term asymptotic expansion

$$\begin{aligned} &(\tilde{a}_h \star \tilde{b}_h - \tilde{b}_h \star \tilde{a}_h)(z, \bar{z}) = \\ &i\hbar \{a, b\} + \\ &i\frac{\hbar^2}{4} (\Delta\{a, b\} + \{a, \Delta b\} + \{\Delta a, b\} + 8\pi\{a, b\}) + \\ &i\frac{\hbar^3}{24} [\{\Delta a, \Delta b\} + \{a, \Delta^2 b\} + \{\Delta^2 a, b\} + \Delta^2\{a, b\} + \\ &\quad \Delta\{a, \Delta b\} + \Delta\{\Delta a, b\} + \\ &\quad 28\pi(\Delta\{a, b\} + \{a, \Delta b\} + \{\Delta a, b\}) + 96\pi^2\{a, b\}] + \\ &o(\hbar^3) \end{aligned}$$

Corollary

Let $\mathcal{A}(\mathbb{D})$ be a subspace of $C^\infty(\mathbb{D})$ such that for each $h \in (0, 1)$ the Toeplitz operator algebra $\mathcal{T}_h(\mathcal{A}(\mathbb{D}))$ is commutative.

Then for all $a, b \in \mathcal{A}(\mathbb{D})$ we have

$$\{a, b\} = 0,$$

$$\{a, \Delta b\} + \{\Delta a, b\} = 0,$$

$$\{\Delta a, \Delta b\} + \{a, \Delta^2 b\} + \{\Delta^2 a, b\} = 0.$$

Let $\mathcal{A}(\mathbb{D})$ be a linear space of smooth functions which generates the commutative C^* -algebra $\mathcal{T}_h(\mathcal{A}(\mathbb{D}))$ of Toeplitz operators for each $h \in (0, 1)$.

Let $\mathcal{A}(\mathbb{D})$ be a linear space of smooth functions which generates the commutative C^* -algebra $\mathcal{T}_h(\mathcal{A}(\mathbb{D}))$ of Toeplitz operators for each $h \in (0, 1)$.

First term: $\{a, b\} = 0$:

Lemma

All functions in $\mathcal{A}(\mathbb{D})$ have (globally) the *same set of level lines* and the same set of *gradient lines*.

Let $\mathcal{A}(\mathbb{D})$ be a linear space of smooth functions which generates the commutative C^* -algebra $\mathcal{T}_h(\mathcal{A}(\mathbb{D}))$ of Toeplitz operators for each $h \in (0, 1)$.

First term: $\{a, b\} = 0$:

Lemma

All functions in $\mathcal{A}(\mathbb{D})$ have (globally) the *same set of level lines* and the same set of *gradient lines*.

Second term: $\{a, \Delta b\} + \{\Delta a, b\} = 0$:

Theorem

The space $\mathcal{A}(\mathbb{D})$ consists of functions whose common *gradient lines* are *geodesics* in the hyperbolic geometry of the unit disk \mathbb{D} .

Let $\mathcal{A}(\mathbb{D})$ be a linear space of smooth functions which generates the commutative C^* -algebra $\mathcal{T}_h(\mathcal{A}(\mathbb{D}))$ of Toeplitz operators for each $h \in (0, 1)$.

First term: $\{a, b\} = 0$:

Lemma

All functions in $\mathcal{A}(\mathbb{D})$ have (globally) the *same set of level lines* and the same set of *gradient lines*.

Second term: $\{a, \Delta b\} + \{\Delta a, b\} = 0$:

Theorem

The space $\mathcal{A}(\mathbb{D})$ consists of functions whose common *gradient lines* are *geodesics* in the hyperbolic geometry of the unit disk \mathbb{D} .

Third term: $\{\Delta a, \Delta b\} + \{a, \Delta^2 b\} + \{\Delta^2 a, b\} = 0$:

Theorem

The space $\mathcal{A}(\mathbb{D})$ consists of functions whose common *level lines* are *cycles*.

Theorem

Let $\mathcal{A}(\mathbb{D})$ be a space of smooth functions. Then the following two statements are equivalent:

- there is a pencil \mathcal{P} of geodesics in \mathbb{D} such that all functions in $\mathcal{A}(\mathbb{D})$ are constant on the cycles of \mathcal{P} ;*
- the C^* -algebra generated by Toeplitz operators with $\mathcal{A}(\mathbb{D})$ -symbols is commutative on each weighted Bergman space $\mathcal{A}_h^2(\mathbb{D})$, $h \in (0, 1)$.*

- Commutative algebras of Toeplitz operators on the unit disk.
- Fine structure of the algebra of Toeplitz operators with PC -symbols.
- From the unit disk to the unit ball.

Continuous symbols

Let $\mathcal{T}(C(\overline{\mathbb{D}}))$ be the C^* -algebra generated by T_a , with $a \in C(\overline{\mathbb{D}})$.

Theorem

The algebra $\mathcal{T} = \mathcal{T}(C(\overline{\mathbb{D}}))$ is irreducible and contains the whole ideal \mathcal{K} of compact on $\mathcal{A}^2(\mathbb{D})$ operators. Each operator $T \in \mathcal{T}(C(\overline{\mathbb{D}}))$ is of the form

$$T = T_a + K, \quad \text{where } a \in C(\overline{\mathbb{D}}), \quad K \in \mathcal{K}.$$

The homomorphism


$$\text{sym} : \mathcal{T} \longrightarrow \text{Sym } \mathcal{T} = \mathcal{T}/\mathcal{K} \cong C(\partial\mathbb{D})$$

is generated by

$$\text{sym} : T_a \longmapsto a|_{\partial\mathbb{D}}.$$


Piece-wise continuous symbols

Fix a finite number of distinct points $T = \{t_1, \dots, t_m\}$ on $\gamma = \partial\mathbb{D}$.
Let ℓ_k , $k = 1, \dots, m$, be the part of the radius of \mathbb{D} starting at t_k .
Let $\mathcal{L} = \bigcup_{k=1}^m \ell_k$.



Piece-wise continuous symbols

Fix a finite number of distinct points $T = \{t_1, \dots, t_m\}$ on $\gamma = \partial\mathbb{D}$.
Let ℓ_k , $k = 1, \dots, m$, be the part of the radius of \mathbb{D} starting at t_k .
Let $\mathcal{L} = \bigcup_{k=1}^m \ell_k$.



Denote by $PC(\overline{\mathbb{D}}, T)$ the set (algebra) of all piece-wise continuous functions on \mathbb{D} which are

- **continuous** in $\overline{\mathbb{D}} \setminus \mathcal{L}$,
- have **one-sided limit values** at each point of \mathcal{L} .

Piece-wise continuous symbols

We consider the C^* -algebra $\mathcal{T}_{PC} = \mathcal{T}(PC(\overline{\mathbb{D}}, \ell))$ generated by all Toeplitz operators T_a with symbols $a(z) \in PC(\overline{\mathbb{D}}, \ell)$.

Piece-wise continuous symbols

We consider the C^* -algebra $\mathcal{T}_{PC} = \mathcal{T}(PC(\overline{\mathbb{D}}, \ell))$ generated by all Toeplitz operators T_a with symbols $a(z) \in PC(\overline{\mathbb{D}}, \ell)$.

Bad news: Let $a(z), b(z) \in PC(\overline{\mathbb{D}}, \ell)$, then

$$[T_a, T_b] = T_a T_b - T_{ab}$$

is not compact in general.

That is

$$T_a T_b \neq T_{ab} + K.$$

The algebra \mathcal{T}_{PC} has a **more complicated** structure.

For piece-wise continuous symbols the C^* -algebra \mathcal{T}_{PC} contains:

- initial generators T_a , where $a \in PC$,



$$\sum_{k=1}^p \prod_{j=1}^{q_k} T_{a_{j,k}}, \quad a_{j,k} \in PC,$$

- uniform limits of sequences of such elements.

Compact set Γ

For each $a_1, a_2 \in PC(\overline{\mathbb{D}}, \ell)$ the commutator $[T_{a_1}, T_{a_2}]$ is compact, thus the algebra $\text{Sym } \mathcal{T}_{PC}$ is commutative.

And thus

$$\text{Sym } \mathcal{T}_{PC} \cong C(\text{over certain compact set } \Gamma).$$

Compact set Γ


For each $a_1, a_2 \in PC(\overline{\mathbb{D}}, \ell)$ the commutator $[T_{a_1}, T_{a_2}]$ is **compact**, thus the algebra $\text{Sym } \mathcal{T}_{PC}$ is **commutative**.

And thus

$$\text{Sym } \mathcal{T}_{PC} \cong C(\text{over certain compact set } \Gamma).$$

The set Γ is the union $\hat{\gamma} \cup (\bigcup_{k=1}^m [0, 1]_k)$, where $\hat{\gamma}$ be the boundary γ , cut by points $t_k \in T$, with the following point identification

$$t_k - 0 \equiv 0_k, \quad t_k + 0 \equiv 1_k.$$



Theorem

The symbol algebra $\text{Sym } \mathcal{T}(PC(\overline{\mathbb{D}}, \ell)) = \mathcal{T}(PC(\overline{\mathbb{D}}, \ell))/\mathcal{K}$ is isomorphic and isometric to $C(\Gamma)$.

The homomorphism

$$\text{sym} : \mathcal{T}(PC(\overline{\mathbb{D}}, \ell)) \rightarrow \text{Sym } \mathcal{T}(PC(\overline{\mathbb{D}}, \ell)) = C(\Gamma)$$

is generated by

$$\text{sym} : T_a \longmapsto \begin{cases} a(t), & t \in \widehat{\gamma} \\ a(t_k - 0)(1 - x) + a(t_k + 0)x, & x \in [0, 1] \end{cases},$$



where $t_k \in T$, $k = 1, 2, \dots, m$.

Auxiliary functions: χ_k

For each $k = 1, \dots, m$, let

$$\chi_k = \chi_k(z)$$

be the **characteristic function** of the **half-disk** obtained by cutting \mathbb{D} by the diameter passing through $t_k \in T$, and such that $\chi_k^+(t_k) = 1$, and thus $\chi_k^-(t_k) = 0$.




Auxiliary functions: v_k

For two **small** neighborhoods $V'_k \subset V''_k$ of the point $t_k \in T$, let

$$v_k = v_k(z) : \overline{\mathbb{D}} \rightarrow [0, 1]$$

be a **continuous** function such that

$$v_k|_{\overline{V'_k}} \equiv 1, \quad v_k|_{\overline{\mathbb{D}} \setminus V''_k} \equiv 0.$$



Canonical form of operators: Generators

Let $a \in PC(\overline{\mathbb{D}}, T)$. Then

$$T_a = T_{s_a} + \sum_{k=1}^m T_{v_k} p_{a,k}(T_{\chi_k}) T_{v_k} + K,$$

where K is compact, $s_a \in C(\overline{\mathbb{D}})$,

$$s_a(z)|_{\gamma} \equiv \left[a(z) - \sum_{k=1}^m [a^-(t_k) + (a^+(t_k) - a^-(t_k))\chi_k(z)] v_k^2(z) \right]_{\gamma},$$

$$p_{a,k}(x) = a^-(t_k)(1-x) + a^+(t_k)x.$$

Canonical form of operators: Sum of products

Let

$$A = \sum_{i=1}^p \prod_{j=1}^{q_i} T_{a_{i,j}},$$

then

$$A = T_{s_A} + \sum_{k=1}^m T_{v_k} p_{A,k}(T_{\chi_k}) T_{v_k} + K_A,$$

where $s_A \in C(\overline{\mathbb{D}})$, $p_{A,k} = p_{A,k}(x)$, $k = 1, \dots, m$, are polynomials, and K_A is compact.

Theorem

Every operator $A \in \mathcal{T}(PC(\overline{\mathbb{D}}, T))$ admits the canonical representations

$$A = T_{s_A} + \sum_{k=1}^m T_{v_k} f_{A,k}(T_{\chi_k}) T_{v_k} + K,$$

where $s_A(z) \in C(\overline{\mathbb{D}})$, $f_{A,k}(x) \in C[0, 1]$, $k = 1, \dots, m$, K is compact.

Toeplitz or not Toeplitz (bounded symbols)

Theorem

An operator

$$A = T_{s_A} + \sum_{k=1}^m T_{v_k} f_{A,k}(T_{\chi_k}) T_{v_k} + K$$

*is a compact perturbation of a Toeplitz operator if and only if every operator $f_{A,k}(T_{\chi_k})$ is a **Toeplitz operator**, where $k = 1, \dots, m$.*

Toeplitz or not Toeplitz (bounded symbols)

Theorem

An operator

$$A = T_{s_A} + \sum_{k=1}^m T_{v_k} f_{A,k}(T_{\chi_k}) T_{v_k} + K$$

is a compact perturbation of a Toeplitz operator if and only if *every operator* $f_{A,k}(T_{\chi_k})$ is a *Toeplitz operator*, where $k = 1, \dots, m$.

Let $f_{A,k}(T_{\chi_k}) = T_{a_k}$ for some $a_k \in L_\infty(\mathbb{D})$. Then $A = T_a + K_A$, where

$$a(z) = s_A(z) + \sum_{k=1}^m a_k(z) v_k^2(z).$$

Example

The Toeplitz operator T_{χ_+} is self-adjoint and $\text{sp } T_{\chi_+} = [0, 1]$.
By functional calculus, for each $f \in C([0, 1])$, the operator $f(T_{\chi_+})$ is well defined and **belongs** to the C^* -algebra generated by T_{χ_+} .

Example


The Toeplitz operator T_{χ_+} is self-adjoint and $\text{sp } T_{\chi_+} = [0, 1]$.
By functional calculus, for each $f \in C([0, 1])$, the operator $f(T_{\chi_+})$ is well defined and **belongs** to the C^* -algebra generated by T_{χ_+} .

For any $\alpha \in (0, 1)$, introduce

$$f_\alpha(x) = x^{2(1-\alpha)} \frac{(1-x)^{2\alpha} - x^{2\alpha}}{(1-x) - x}, \quad x \in [0, 1].$$

Then

$$f_\alpha(T_{\chi_+}) = T_{\chi_{[0, \alpha\pi]}}.$$



Example

Let $p(x) = \sum_{k=1}^n a_k x^k$ be a polynomial of degree $n \geq 2$. Then the bounded operator $p(T_{\chi_+})$ is **not** a **Toeplitz** operator.

Example

Let $p(x) = \sum_{k=1}^n a_k x^k$ be a polynomial of degree $n \geq 2$. Then the bounded operator $p(T_{\chi_+})$ is **not** a **Toeplitz** operator.

Corollary

Let

$$A = \sum_{i=1}^p \prod_{j=1}^{q_i} T_{a_{i,j}} \in \mathcal{T}(PC(\overline{\mathbb{D}}, T)).$$

Then A is a compact perturbation of a **Toeplitz operator** if and only if A is a compact perturbation an **initial generator** T_a , for some $a \in PC(\overline{\mathbb{D}}, T)$.

- Each operator $A \in \mathcal{T}(PC(\overline{\mathbb{D}}, T))$ admits a transparent **canonical representation**

$$A = T_{s_A} + \sum_{k=1}^m T_{v_k} f_{A,k}(T_{\chi_k}) T_{v_k} + K.$$

- Each operator $A \in \mathcal{T}(PC(\overline{\mathbb{D}}, T))$ admits a transparent **canonical representation**

$$A = T_{s_A} + \sum_{k=1}^m T_{v_k} f_{A,k}(T_{\chi_k}) T_{v_k} + K.$$

- All initial generators T_a , $a \in PC(\overline{\mathbb{D}}, T)$ are **Toeplitz operators**.

- Each operator $A \in \mathcal{T}(PC(\overline{\mathbb{D}}, T))$ admits a transparent **canonical representation**

$$A = T_{s_A} + \sum_{k=1}^m T_{v_k} f_{A,k}(T_{\chi_k}) T_{v_k} + K.$$

- All initial generators T_a , $a \in PC(\overline{\mathbb{D}}, T)$ are **Toeplitz operators**.
- **None** of the (non trivial) elements

$$\sum_{i=1}^p \prod_{j=1}^{q_i} T_{a_{i,j}},$$

is a compact perturbation of a Toeplitz operator.






- The uniform closure contains a **huge** amount of Toeplitz operators, with **bounded** and even **unbounded** symbols, which are **drastically different** from the initial generators.

- The uniform closure contains a **huge** amount of Toeplitz operators, with **bounded** and even **unbounded** symbols, which are **drastically different** from the initial generators.
- All these **Toeplitz operators** are uniform limits of sequences of **non-Toeplitz** operators.

- The uniform closure contains a **huge** amount of Toeplitz operators, with **bounded** and even **unbounded** symbols, which are **drastically different** from the initial generators.
- All these **Toeplitz operators** are uniform limits of sequences of **non-Toeplitz** operators.
- The uniform closure contains as well many **non-Toeplitz** operators.

- Commutative algebras of Toeplitz operators on the unit disk.
- Fine structure of the algebra of Toeplitz operators with PC -symbols.
- From the unit disk to the unit ball.

Model cases



Model Maximal Commutative Subgroups

- **Elliptic:** \mathbb{T} , with $z \in \mathbb{D} \mapsto tz \in \mathbb{D}$, $t \in \mathbb{T}$,
- **Hyperbolic:** \mathbb{R}_+ , with $z \in \Pi \mapsto rz \in \Pi$, $r \in \mathbb{R}_+$,
- **Parabolic:** \mathbb{R} , with $z \in \Pi \mapsto z + h \in \Pi$, $h \in \mathbb{R}$.

Unit ball

We consider the unit ball \mathbb{B}^n in \mathbb{C}^n ,

$$\mathbb{B}^n = \{z = (z_1, \dots, z_n) \in \mathbb{C}^n : |z|^2 = |z_1|^2 + \dots + |z_n|^2 < 1\}.$$

Unit ball

We consider the unit ball \mathbb{B}^n in \mathbb{C}^n ,

$$\mathbb{B}^n = \{z = (z_1, \dots, z_n) \in \mathbb{C}^n : |z|^2 = |z_1|^2 + \dots + |z_n|^2 < 1\}.$$

For each $\lambda \in (-1, \infty)$, introduce the measure

$$d\mu_\lambda(z) = c_\lambda (1 - |z|^2)^\lambda dv(z),$$

where $dv(z) = dx_1 dy_1 \dots dx_n dy_n$ and

$$c_\lambda = \frac{\Gamma(n + \lambda + 1)}{\pi^n \Gamma(\lambda + 1)}.$$

The (weighted) **Bergman space** $\mathcal{A}_\lambda^2(\mathbb{B}^n)$ is the subspace of $L_2(\mathbb{B}^n, d\mu_\lambda)$ consisting of functions analytic in \mathbb{B}^n .

The orthogonal **Bergman projection** has the form

$$(B_{\mathbb{B}^n} \varphi)(z) = \int_{\mathbb{B}^n} \varphi(\zeta) \frac{(1 - |\zeta|^2)^\lambda}{(1 - z \cdot \bar{\zeta})^{n+\lambda+1}} c_\lambda dv(\zeta).$$

Unbounded realizations

The standard **unbounded** realization of the unit disk \mathbb{D} is the **upper half-plane**

$$\Pi = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}.$$

The standard **unbounded** realization of the unit ball \mathbb{B}^n is the **Siegel domain** in \mathbb{C}^n

$$D_n = \{z = (z', z_n) \in \mathbb{C}^{n-1} \times \mathbb{C} : \operatorname{Im} z_n - |z'|^2 > 0\},$$

where we use the following notation for the points of $\mathbb{C}^n = \mathbb{C}^{n-1} \times \mathbb{C}$:

$$z = (z', z_n), \quad \text{where } z' = (z_1, \dots, z_{n-1}) \in \mathbb{C}^{n-1}, \quad z_n \in \mathbb{C}.$$

Model Maximal Commutative Subgroups

- **Quasi-elliptic:** \mathbb{T}^n , for each $t = (t_1, \dots, t_n) \in \mathbb{T}^n$:
 $z = (z_1, \dots, z_n) \in \mathbb{B}^n \mapsto tz = (t_1 z_1, \dots, t_n z_n) \in \mathbb{B}^n$;
- **Quasi-hyperbolic:** $\mathbb{T}^{n-1} \times \mathbb{R}_+$, for each $(t, r) \in \mathbb{T}^{n-1} \times \mathbb{R}_+$:
 $(z', z_n) \in D_n \mapsto (r^{1/2} t z', r z_n) \in D_n$;
- **Quasi-parabolic:** $\mathbb{T}^{n-1} \times \mathbb{R}$, for each $(t, h) \in \mathbb{T}^{n-1} \times \mathbb{R}$:
 $(z', z_n) \in D_n \mapsto (t z', z_n + h) \in D_n$;
- **Nilpotent:** $\mathbb{R}^{n-1} \times \mathbb{R}$, for each $(b, h) \in \mathbb{R}^{n-1} \times \mathbb{R}$:
 $(z', z_n) \in D_n \mapsto (z' + b, z_n + h + 2iz' \cdot b + i|b|^2) \in D_n$;
- **Quasi-nilpotent:** $\mathbb{T}^k \times \mathbb{R}^{n-k-1} \times \mathbb{R}$, $0 < k < n - 1$,
for each $(t, b, h) \in \mathbb{T}^k \times \mathbb{R}^{n-k-1} \times \mathbb{R}$:
 $(z', z'', z_n) \in D_n \mapsto (t z', z'' + b, z_n + h + 2iz'' \cdot b + i|b|^2) \in D_n$.

Classification Theorem

Theorem

Given any maximal commutative subgroup G of biholomorphisms of the unit ball \mathbb{B}^n , denote by \mathcal{A}_G the set of all $L_\infty(\mathbb{B}^n)$ -functions which are *invariant* under the action of G .

Then the C^* -algebra generated by Toeplitz operators with symbols from \mathcal{A}_G is *commutative* on each weighted Bergman space $\mathcal{A}_\lambda^2(\mathbb{B}^n)$, $\lambda \in (-1, \infty)$.

Classification Theorem

Theorem

Given any maximal commutative subgroup G of biholomorphisms of the unit ball \mathbb{B}^n , denote by \mathcal{A}_G the set of all $L_\infty(\mathbb{B}^n)$ -functions which are *invariant* under the action of G .

Then the C^* -algebra generated by Toeplitz operators with symbols from \mathcal{A}_G is *commutative* on each weighted Bergman space $\mathcal{A}_\lambda^2(\mathbb{B}^n)$, $\lambda \in (-1, \infty)$.

The result can be alternatively formulated in terms of the so-called *Lagrangian frames*, the multidimensional analog of pencils of geodesics and cycles of the unit disk.

It was firmly expected that the situation for the unit ball is pretty much the same as in the case of the unit disk, that is:

The above algebras exhaust all possible algebras of Toeplitz operators on the unit ball which are commutative on each weighted Bergman space.

It was firmly expected that the situation for the unit ball is pretty much the same as in the case of the unit disk, that is:

The above algebras exhaust all possible algebras of Toeplitz operators on the unit ball which are commutative on each weighted Bergman space.

But:

It turns out that there exist many other Banach algebras generated by Toeplitz operators which are commutative on each weighted Bergman space, non of them is a C^* -algebra, and for $n = 1$ all of them collapse to known commutative C^* -algebras of the unit disk.