

# Asymptotic properties of solutions of a $2n^{\text{th}}$ order differential equation on a time scale

Douglas Anderson  
Department of Mathematics and Computer Science  
Concordia College  
Moorhead, MN 56562  
andersod@cord.edu

Allan Peterson  
Department of Mathematics and Statistics  
University of Nebraska-Lincoln  
Lincoln, NE 68588-0323  
apeterso@math.unl.edu

**ABSTRACT:** In this paper we are concerned with a  $2n$ -th order linear self-adjoint differential equation on a time scale. The results generalize known results for the corresponding ordinary differential equations and for difference equations. We define type I and type II solutions, prove the existence of these solutions, and verify asymptotic properties of these solutions. A quadratic functional corresponding to the differential equation on a time scale is defined and is used to prove several of the results in this paper.

**AMS Subject Classification.** 34B10.

## 1 INTRODUCTION

In this paper we will be concerned with the  $2n$ th-order differential equation

$$Lx(t) = \left[ p(t)x^{\Delta^n}(t) \right]^{\Delta^n} + q(t)x(\sigma^n(t)) = 0 \quad (1)$$

for  $t \in \mathbb{T}$ . We assume  $\mathbb{T}$  is a closed subset of the real numbers  $\mathbb{R}$  that is unbounded above. We call  $\mathbb{T}$  a time scale.

**Definition** For  $t \in \mathbb{T}$  define the forward jump operator  $\sigma$  by

$$\sigma(t) := \inf\{\tau \in \mathbb{T} : \tau > t\}.$$

We define the backwards jump operator  $\rho$  for  $t \in \mathbb{T}$ ,  $t > \inf \mathbb{T}$ , by

$$\rho(t) := \sup\{\tau \in \mathbb{T} : \tau < t\}.$$

If  $\sigma(t) = t$  we say  $t$  is *right-dense*, and if  $\sigma(t) > t$  we say  $t$  is *right-scattered*. Similarly, if  $\rho(t) = t$  then  $t$  is *left-dense*, and if  $\rho(t) < t$  then  $t$  is *left-scattered*. We assume  $\sigma^0(t) = t$ , and for any integer  $n > 0$ , we define  $\sigma^n : \mathbb{T} \rightarrow \mathbb{R}$  by

$$\sigma^n(t) := \sigma(\sigma^{n-1}(t)).$$

We assume throughout that the time scale  $\mathbb{T}$  has the topology it inherits from the standard topology on  $\mathbb{R}$ . We also assume  $p, q : \mathbb{T} \rightarrow \mathbb{R}$  are continuous and

$$p(t) > 0$$

on  $\mathbb{T}$ .

**Definition** Assume  $x : \mathbb{T} \rightarrow \mathbb{R}$  and fix  $t \in \mathbb{T}$ ; define  $x^\Delta(t)$  to be the number (provided it exists) with the property that given any  $\epsilon > 0$ , there is a neighborhood  $U$  of  $t$  such that

$$\left| [x(\sigma(t)) - x(s)] - x^\Delta(t)[\sigma(t) - s] \right| < \epsilon |\sigma(t) - s|$$

for all  $s \in U$ . We call  $x^\Delta(t)$  the *delta derivative* of  $x$  at  $t$ . Note if  $\mathbb{T} = \mathbb{Z}$ , then

$$x^\Delta(t) = \Delta x(t) := x(t+1) - x(t).$$

Moreover, if  $\mathbb{T} = \mathbb{R}$ , then

$$x^\Delta(t) = x'(t).$$

Hence, our results contain differential equations and difference equations as special cases. Finally, if  $h > 0$  and  $\mathbb{T} = h\mathbb{Z} = \{hn : n \in \mathbb{Z}\}$ , then

$$x^\Delta(t) = \frac{x(t+h) - x(t)}{h}.$$

**Definition** For  $n \geq 1$ , define

$$x^{\Delta^n}(t) := \left\{ x^\Delta(t) \right\}^{\Delta^{n-1}};$$

assume  $x^{\Delta^0}(t) = x(t)$ .

**Definition** We say  $x : \mathbb{T} \rightarrow \mathbb{R}$  is *right-dense continuous* provided for any  $t \in \mathbb{T}$  such that  $\rho(t) = t$ , the lefthand limit of  $x$  at  $t$  exists (and is finite). Also, if  $t \in \mathbb{T}$  and  $\sigma(t) = t$ , then

$$\lim_{\tau \rightarrow t^+} x(\tau) = x(t).$$

Note if  $x : \mathbb{T} \rightarrow \mathbb{R}$  is continuous, then  $x(\sigma(t))$  is right-dense continuous on  $\mathbb{T}$ .

**Definition** We define

$$D := \left\{ x : \mathbb{T} \rightarrow \mathbb{R} : x^{\Delta^k}(\sigma^i(t)) \text{ is delta differentiable on } \mathbb{T} \text{ for } 0 \leq k \leq n-1 \right. \\ \left. \text{and } 0 \leq i \leq n; \left[ p(\sigma^i(t)) x^{\Delta^n}(\sigma^i(t)) \right]^{\Delta^k} \text{ is delta differentiable on } \mathbb{T} \text{ for } \right. \\ \left. 0 \leq i, k \leq n-1; \text{ and } [p(t) x^{\Delta^n}(t)]^{\Delta^n} \text{ is right-dense continuous on } \mathbb{T} \right\}.$$

We say  $x$  is a solution of  $Lx = 0$  on  $\mathbb{T}$  provided  $x \in D$  and  $Lx(t) = 0$  for all  $t \in \mathbb{T}$ . Our results will hold only for those time scales  $\mathbb{T}$  for which  $D$  is nonempty. Moreover, we need to assume below that the time scale  $\mathbb{T}$  is such that

$$[x^{\Delta^{n-1}}(\sigma^n(t))]^\Delta = x^{\Delta^n}(\sigma^n(t)).$$

This is true for time scales  $\mathbb{T} = \mathbb{R}$  and  $\mathbb{T} = h\mathbb{Z}$  for any  $h > 0$ , but of course is not true in general.

**Definition** For  $x \in D$ , set

$$\begin{aligned} Fx(t) &:= x^{\Delta^{n-1}}(\sigma^{n-1}(t))p(\sigma^{n-1}(t))x^{\Delta^n}(\sigma^{n-1}(t)) \\ &\quad - (-1)^n \sum_{i=0}^{n-2} (-1)^i x^{\Delta^i}(\sigma^n(t)) \left[ p(\sigma^i(t))x^{\Delta^n}(\sigma^i(t)) \right]^{\Delta^{n-1-i}}; \end{aligned} \quad (2)$$

this quadratic functional  $F$  will be important in the study of the asymptotic properties of solutions of  $Lx(t) = 0$ . Note that  $F[kx(t)] = k^2F(x(t))$ .

In the proof of the following lemma, we will make repeated use of the product formula

$$[f(t)g(t)]^\Delta = f(\sigma(t))g^\Delta(t) + f^\Delta(t)g(t),$$

where  $f$  and  $g$  are interchangeable.

**Lemma 1** *If  $x$  is a solution of (1) on  $\mathbb{T}$ , then*

$$[Fx(t)]^\Delta = p(\sigma^{n-1}(t)) \left[ x^{\Delta^n}(\sigma^{n-1}(t)) \right]^2 + (-1)^n q(t)x^2(\sigma^n(t)) \quad (3)$$

for  $t \in \mathbb{T}$ . In particular, if

$$(-1)^n q(t) \geq 0 \quad (4)$$

on  $\mathbb{T}$ , then  $F$  is nondecreasing along solutions  $x$  of (1) for  $t \in \mathbb{T}$ .

*Proof:* Assume  $x$  is a solution of (1). Then

$$[Fx(t)]^\Delta = x^{\Delta^{n-1}}(\sigma^n(t)) \left[ p(\sigma^{n-1}(t))x^{\Delta^n}(\sigma^{n-1}(t)) \right]^\Delta \quad (5)$$

$$+ x^{\Delta^n}(\sigma^{n-1}(t))p(\sigma^{n-1}(t))x^{\Delta^n}(\sigma^{n-1}(t)) \quad (6)$$

$$- (-1)^n \sum_{i=0}^{n-2} (-1)^i x^{\Delta^i}(\sigma^n(t)) \left[ p(\sigma^i(t))x^{\Delta^n}(\sigma^i(t)) \right]^{\Delta^{n-i}} \quad (7)$$

$$- (-1)^n \sum_{i=0}^{n-2} (-1)^i x^{\Delta^{i+1}}(\sigma^n(t)) \left[ p(\sigma^{i+1}(t))x^{\Delta^n}(\sigma^{i+1}(t)) \right]^{\Delta^{n-i-1}} \quad (8)$$

where (5) and (6) result from the product rule of the first advanced times the delta derivative of the second plus the second times the derivative of the first, while (7) and (8) come from the product rule of the first times the delta derivative of the second

plus the derivative of the first times the second advanced. Next, evaluating the sum in (8) at  $n - 2$  and reindexing, we obtain

$$\begin{aligned}
[Fx(t)]^\Delta &= x^{\Delta^{n-1}}(\sigma^n(t)) \left[ p(\sigma^{n-1}(t)) x^{\Delta^n}(\sigma^{n-1}(t)) \right]^\Delta + p(\sigma^{n-1}(t)) \left[ x^{\Delta^n}(\sigma^{n-1}(t)) \right]^\Delta \\
&\quad - (-1)^n (-1)^{n-2} x^{\Delta^{n-1}}(\sigma^n(t)) \left[ p(\sigma^{n-1}(t)) x^{\Delta^n}(\sigma^{n-1}(t)) \right]^\Delta \\
&\quad - (-1)^n \sum_{i=0}^{n-2} (-1)^i x^{\Delta^i}(\sigma^n(t)) \left[ p(\sigma^i(t)) x^{\Delta^n}(\sigma^i(t)) \right]^\Delta \\
&\quad - (-1)^n \sum_{i=1}^{n-2} (-1)^{i-1} x^{\Delta^i}(\sigma^n(t)) \left[ p(\sigma^i(t)) x^{\Delta^n}(\sigma^i(t)) \right]^\Delta \\
&= p(\sigma^{n-1}(t)) \left[ x^{\Delta^n}(\sigma^{n-1}(t)) \right]^2 - (-1)^n x(\sigma^n(t)) \left[ p(t) x^{\Delta^n}(t) \right]^\Delta \\
&= p(\sigma^{n-1}(t)) \left[ x^{\Delta^n}(\sigma^{n-1}(t)) \right]^2 + (-1)^n q(t) x^2(\sigma^n(t)),
\end{aligned}$$

since  $x$  is a solution of (1).

**Definitions** If  $x(t)$  is a solution of (1) such that  $Fx(t) \leq 0$  in a neighborhood of  $\infty$ , we say  $x(t)$  is a *type I solution* of (1). If  $x(t)$  is a solution of (1) such that  $Fx(t) < 0$  for  $t \in \mathbb{T}$ , then we say  $x(t)$  is a *strict type I solution*. If  $x$  solves (1) and  $Fx(t) > 0$  near  $\infty$ , then  $x$  is a *type II solution*. In view of Lemma 1, all solutions of (1) are type I or type II solutions if (4) holds.

Hartman [3] defined generalized zeros of solutions of  $n^{\text{th}}$ -order linear difference equations. We now would like to define what we mean by a *generalized zero of order  $m$* ,  $0 \leq m \leq 2n - 1$ , for a nontrivial solution  $x(t)$  of (1) at  $t_0 \in \mathbb{T}$ . If  $t_0 = \min \mathbb{T}$ , then we say  $x(t)$  has a generalized zero of order  $m$  at  $t_0$  provided

$$x^{\Delta^k}(t_0) = 0,$$

$0 \leq k \leq m - 1$ . If  $t_0 > \min \mathbb{T}$ , then  $x$  has a generalized zero of order  $m$  at  $t_0$  provided if  $t_0$  is left-dense then

$$x^{\Delta^k}(t_0) = 0,$$

$0 \leq k \leq m - 1$ ; if  $t_0$  is left-scattered, then

$$x(\rho(t_0)) \neq 0 \text{ and } x^{\Delta^k}(t_0) = 0$$

for  $0 \leq k \leq m - 1$ , or  $t_0$  is right-scattered with

$$x^{\Delta^k}(t_0) = 0$$

for  $0 \leq k \leq m - 2$  and

$$(-1)^m x(\rho(t_0)) x(\sigma^{m-1}(t_0)) > 0.$$

(This last case happens only if  $\rho(t_0) < t_0 < \sigma(t_0) < \dots < \sigma^{m-1}(t_0)$ .)

## 2 MAIN RESULTS

**Theorem 2** Assume (4) holds on  $\mathbb{T}$ . Any nontrivial solution  $x$  of (1) with

$$x^{\Delta^k}(\sigma^n(t_0)) = 0,$$

$0 \leq k \leq n-1$ , for some  $t_0 \in \mathbb{T}$  is a type II solution. In particular, (1) has at least  $n$  linearly independent type II solutions.

*Proof:* Let  $t_0 \in \mathbb{T}$ , and let  $x_j(t)$ ,  $0 \leq j \leq n-1$ , be solutions of  $Lx = 0$  satisfying

$$x_j^{\Delta^k}(\sigma^n(t_0)) = 0, \quad 0 \leq k \leq n-1 \quad (9)$$

$$\left[ p(t)x_j^{\Delta^k}(t) \right]^{\Delta^k}(t_0) = \delta_{jk}, \quad 0 \leq j, k \leq n-1.$$

It is easy to see that  $\{x_j(t)\}_{j=0}^{n-1}$  is a set of  $n$  linearly independent solutions of (1), and because of (9),

$$Fx_j(\sigma(t_0)) = 0. \quad (10)$$

Integrating both sides of (3) from  $\sigma(t_0)$  to  $t$  and using (10) we get

$$Fx_j(t) = \int_{\sigma(t_0)}^t \left[ p(\sigma^{n-1}(s)) \left[ x_j^{\Delta^n}(\sigma^{n-1}(s)) \right]^2 + (-1)^n q(s) x_j^2(\sigma^n(s)) \right] \Delta s.$$

Since  $x_j(t)$  is a nontrivial solution, by Lemma 1 we can pick  $\tau > \sigma(t_0)$  sufficiently large so that

$$Fx_j(t) > 0$$

for all  $t \geq \tau$ . Hence,  $x_1, \dots, x_n$  are  $n$  linearly independent type II solutions.

**Theorem 3** Assume (4) holds. Then (1) has  $n$  linearly independent type I solutions.

*Proof:* Let  $a \in \mathbb{T}$ . For each fixed  $s \in \mathbb{T}$ ,  $s > \sigma^{2n}(a)$ , we let  $v_k(t, s)$  be a nontrivial solution of (1) for  $1 \leq k \leq n$ , satisfying the  $2n-1$  boundary conditions

$$\begin{aligned} v_k^{\Delta^i}(\sigma^n(a), s) &= 0 & \text{for } 0 \leq i \leq n-1 \text{ but } i \neq k-1 \\ v_k^{\Delta^i}(\sigma^n(s), s) &= 0 & \text{for } 0 \leq i \leq n-1. \end{aligned} \quad (11)$$

Then define

$$u_k(t, s) := \frac{v_k(t, s)}{\sqrt{[v_k(\sigma^n(a), s)]^2 + [v_k^{\Delta}(\sigma^n(a), s)]^2 + \dots + [v_k^{\Delta^{2n-1}}(\sigma^n(a), s)]^2}} \quad (12)$$

for  $1 \leq k \leq n$  and  $s > \sigma^{2n}(a)$ . Then  $u_k(t, s)$  is a solution of (1) satisfying

$$\sum_{i=0}^{2n-1} \left[ u_k^{\Delta^i}(\sigma^n(a), s) \right]^2 = 1.$$

Thus, for each  $k$ , and  $s \in \mathbb{T}$  with  $s > \sigma^{2n}(a)$  (recall  $\mathbb{T}$  is unbounded above), from the set of points

$$\left\{ (u_k(\sigma^n(a), s), u_k^{\Delta}(\sigma^n(a), s), \dots, u_k^{\Delta^{2n-1}}(\sigma^n(a), s)) \right\}_{s \in \mathbb{T}}$$

in  $\mathbb{R}^{2n}$  we can pick a convergent sequence

$$\left\{ (u_k(\sigma^n(a), s_{j_k}), u_k^\Delta(\sigma^n(a), s_{j_k}), \dots, u_k^{\Delta^{2n-1}}(\sigma^n(a), s_{j_k})) \right\}_{j=1}^\infty.$$

Let

$$v_{i+1,k} := \lim_{j \rightarrow \infty} u_k^{\Delta^i}(\sigma^n(a), s_{j_k}) \quad (13)$$

for  $0 \leq i \leq 2n - 1$ . Then

$$\sum_{i=0}^{2n-1} v_{i+1,k}^2 = 1. \quad (14)$$

Further, let  $y_k$ ,  $1 \leq k \leq n$ , be the solutions of (1) satisfying

$$y_k^{\Delta^i}(\sigma^n(a)) = v_{i+1,k} \quad (15)$$

for  $0 \leq i \leq 2n - 1$ ,  $1 \leq k \leq n$ ; note that the  $y_k$  are nontrivial solutions by (14) and (15). Since  $v_k(t, s)$  satisfies (11), formula (12) implies  $u_k(t, s)$ ,  $1 \leq k \leq n$ , also satisfies (11). Hence, we have

$$Fu_k(\sigma(s_{j_k}), s_{j_k}) = 0;$$

as  $Fu_k(t, s_{j_k})$  is nondecreasing by Lemma 1,

$$Fu_k(t, s_{j_k}) \leq 0$$

for  $t \in \mathbb{T}$  with  $\sigma^n(a) \leq t \leq \sigma(s_{j_k})$ , for  $1 \leq k \leq n$ . So, for each  $t_1 \in \mathbb{T}$  with  $t_1 > \sigma^n(a)$ , there exists  $j_{t_1}$  such that  $\sigma(s_{j_k}) > t_1$  for all  $j_k \geq j_{t_1}$ . Then

$$Fu_k(t_1, s_{j_k}) \leq 0$$

for all  $j_k \geq j_{t_1}$ . Taking the limit as  $j \rightarrow \infty$ , we get that

$$Fy_k(t_1) \leq 0$$

for  $1 \leq k \leq n$ . As  $t_1 > \sigma^n(a)$  was arbitrary,

$$Fy_k(t) \leq 0$$

for all  $t > \sigma^n(a)$ , for  $1 \leq k \leq n$ . It follows that the  $y_k$  are type I solutions of (1) for  $1 \leq k \leq n$ . By (11),(12),(13), and (15) we have that

$$y_k^{\Delta^i}(\sigma^n(a)) = 0 \text{ for } 0 \leq i \leq n - 1 \text{ if } i \neq k - 1.$$

If  $y_k^{\Delta^{k-1}}(\sigma^n(a)) = 0$ , then  $y_k$  would have a zero of order  $n$  at  $\sigma^n(a)$ , so that  $y_k$  would be a type II solution by Theorem 2. Consequently,  $y_k^{\Delta^{k-1}}(\sigma^n(a)) \neq 0$  for  $1 \leq k \leq n$ , and the  $y_k$  are linearly independent.

**Theorem 4** *If (4) holds on  $\mathbb{T}$  and  $x$  is a type I solution of (1), then*

$$(-1)^n \int^\infty q(t)x^2(\sigma^n(t))\Delta t < \infty, \quad (16)$$

and

$$\int^\infty p(t) [x^{\Delta^n}(t)]^2 \Delta t < \infty. \quad (17)$$

*If  $q(t) \neq 0$  in a neighborhood of  $\infty$ , then every nontrivial type I solution of (1) is a strict type I solution.*

*Proof:* Let  $x$  be a type I solution of (1) on  $\mathbb{T}$ ; then  $Fx(t) \leq 0$  for  $t \in \mathbb{T}$ . Let

$$M := \lim_{t \rightarrow \infty} Fx(t) \leq 0.$$

Integrating both sides of (3) from  $t_0 \in \mathbb{T}$  to  $\infty$  yields

$$\begin{aligned} M - Fx(t_0) &= \int_{t_0}^{\infty} p(\sigma^{n-1}(t)) \left[ x^{\Delta^n}(\sigma^{n-1}(t)) \right]^2 \Delta t \\ &+ \int_{t_0}^{\infty} (-1)^n q(t) x^2(\sigma^n(t)) \Delta t. \end{aligned}$$

Thus (16) and (17) hold. Now assume  $q(t) \neq 0$  in a neighborhood of  $\infty$ , and that  $v$  is a nontrivial type I solution of (1); then  $Fv(t) \leq 0$  for  $t \in \mathbb{T}$ . Suppose there exists a  $t_0 \in \mathbb{T}$  such that  $Fv(t_0) = 0$ . Then  $Fv(t) \equiv 0$  on  $[t_0, \infty) \cap \mathbb{T}$ , by Lemma 1. But then  $[Fv(t)]^\Delta = 0$  on  $[t_0, \infty) \cap \mathbb{T}$ , so that from (3) we have

$$p(\sigma^{n-1}(t)) \left[ v^{\Delta^n}(\sigma^{n-1}(t)) \right]^2 + (-1)^n q(t) v^2(\sigma^n(t)) \equiv 0$$

for  $t \geq t_0 \in \mathbb{T}$ . Since (4) holds all terms are nonnegative; moreover,  $q(t) \neq 0$  near  $\infty$  gives that  $v$  is the trivial solution, contrary to assumption. Therefore,  $Fv(t) < 0$  for all  $t \in \mathbb{T}$ , and  $v$  is a strict type I solution of (1).

We conjecture that if (4) holds and  $\liminf_{t \rightarrow \infty} (-1)^n q(t) > 0$ , then (1) has  $n$  linearly independent type I solutions  $v_k$  satisfying  $\lim_{t \rightarrow \infty} v_k(t) = 0$  for  $1 \leq k \leq n$ .

**Theorem 5** *If (4) holds on  $\mathbb{T}$ , then no nontrivial solution of (1) can have a zero of order  $n$  at  $\sigma^n(t_1)$  followed by a generalized zero of order  $n$  at  $\sigma^n(t_2)$ , where  $t_1, t_2 \in \mathbb{T}$  with  $t_2 > \sigma^n(t_1)$  and  $\sigma^n(t_2)$  is left-scattered.*

*Proof:* Assume (4) holds. Assume  $x$  is a nontrivial solution of (1) and  $t_1 \in \mathbb{T}$  such that

$$x^{\Delta^i}(\sigma^n(t_1)) = 0$$

for  $0 \leq i \leq n-1$ . Then  $Fx(\sigma(t_1)) = 0$ ; by Lemma 1,

$$Fx(t) \geq 0 \tag{18}$$

for all  $t \geq \sigma(t_1) \in \mathbb{T}$ . Suppose there is a  $t_2 > \sigma^n(t_1) \in \mathbb{T}$  such that  $x$  has a generalized zero of order  $n$  at  $\sigma^n(t_2)$ . Then  $x(\sigma^{n-1}(t_2)) \neq 0$ , and by (2),

$$\begin{aligned} Fx(t_2) &= x^{\Delta^{n-1}}(\sigma^{n-1}(t_2)) p(\sigma^{n-1}(t_2)) x^{\Delta^n}(\sigma^{n-1}(t_2)) \\ &= \frac{(-1)^{n-1} x(\sigma^{n-1}(t_2)) p(\sigma^{n-1}(t_2)) x^{\Delta^n}(\sigma^{n-1}(t_2))}{\alpha^{n-1}(t_2)} \\ &= \frac{(-1)^{n-1} x(\sigma^{n-1}(t_2)) p(\sigma^{n-1}(t_2))}{\alpha^{n-1}(t_2)} \left[ \frac{x(\sigma^{2n-1}(t_2))}{\beta(t_2)} + \frac{(-1)^n x(\sigma^{n-1}(t_2))}{\alpha^n(t_2)} \right] \\ &= \frac{(-1)^{n-1} x(\sigma^{n-1}(t_2)) x(\sigma^{2n-1}(t_2)) p(\sigma^{n-1}(t_2))}{\alpha^{n-1}(t_2) \beta(t_2)} - \frac{x^2(\sigma^{n-1}(t_2)) p(\sigma^{n-1}(t_2))}{\alpha^{2n-1}(t_2)}, \end{aligned}$$

where

$$\alpha(t) := \sigma^n(t) - \sigma^{n-1}(t)$$

and

$$\beta(t) := [\sigma^{2n-1}(t) - \sigma^{2n-2}(t)][\sigma^{2n-2}(t) - \sigma^{2n-3}(t)] \cdots [\sigma^n(t) - \sigma^{n-1}(t)];$$

we assume  $\alpha(t_2)$  and  $\beta(t_2)$  are nonzero when  $x(\sigma^{2n-1}(t_2)) \neq 0$ . Since  $x$  has a generalized zero of order  $n$  at  $\sigma^n(t_2)$ ,

$$(-1)^n x(\sigma^{n-1}(t_2))x(\sigma^{2n-1}(t_2)) \geq 0.$$

Therefore  $Fx(t_2) < 0$ , a contradiction of (18) since  $t_2 > \sigma(t_1)$ .

**Theorem 6** *Every unbounded solution near  $\infty$  of (1), where*

$$\liminf_{t \rightarrow \infty} q(t) > 0 \tag{19}$$

and

$$0 < \liminf_{t \rightarrow \infty} p(t) < \infty \tag{20}$$

for  $t \in \mathbb{T}$ , is oscillatory.

*Proof:* Let  $x$  be an unbounded solution of (1). Assume that  $x$  is nonoscillatory; then without loss of generality there exists  $t_0$  in  $\mathbb{T}$  such that  $x(t) > 0$  for all  $t \geq t_0$  in  $\mathbb{T}$ . Since  $x$  is a solution of (1), by repeated application of Rolle's Theorem we can assume  $x$  is increasing. From (19) we can also assume  $t_0$  is sufficiently large so that  $q(t) > 0$  for  $t \geq t_0$  in  $\mathbb{T}$ . Therefore

$$\left[ p(t)x^{\Delta^n}(t) \right]^{\Delta^n} = -q(t)x(\sigma^n(t)) < 0 \tag{21}$$

for  $t \geq t_0$  in  $\mathbb{T}$ , and

$$\lim_{t \rightarrow \infty} \left[ p(t)x^{\Delta^n}(t) \right]^{\Delta^n} = \lim_{t \rightarrow \infty} -q(t)x(\sigma^n(t)) = -\infty. \tag{22}$$

And yet

$$\left[ p(t)x^{\Delta^n}(t) \right]^{\Delta^{n-1}} - \left[ p(t_0)x^{\Delta^n}(t_0) \right]^{\Delta^{n-1}} = \int_{t_0}^t \left[ p(s)x^{\Delta^n}(s) \right]^{\Delta^n} \Delta s \rightarrow -\infty. \tag{23}$$

by (22). As a result, there is a  $t_1 > t_0$  in  $\mathbb{T}$  such that

$$\left[ p(t)x^{\Delta^n}(t) \right]^{\Delta^{n-1}} < 0$$

for all  $t \geq t_1$  in  $\mathbb{T}$ ,

$$\liminf_{t \rightarrow \infty} \left[ p(t)x^{\Delta^n}(t) \right]^{\Delta^{n-1}} = -\infty \tag{24}$$

for  $t \in \mathbb{T}$  using (21), (22), (23), and so on. By continuing this process of integrating each successive expression we obtain

$$\liminf_{t \rightarrow \infty} \left[ p(t)x^{\Delta^n}(t) \right]^{\Delta^i} = -\infty$$

for  $t \in \mathbb{T}$ ,  $i = n - 2, n - 3, \dots, 1, 0$ . Using (20) and similar reasoning, we see for  $i = n, n - 1, \dots, 1, 0$  that

$$\liminf_{t \rightarrow \infty} x^{\Delta^i}(t) = -\infty$$

as well. Consequently

$$\liminf_{t \rightarrow \infty} x(t) = -\infty,$$

a contradiction of our initial assumption that  $x(t) > 0$  for  $t \geq t_0$  in  $\mathbb{T}$ . Thus every unbounded solution of (1) is oscillatory under conditions (19) and (20).

**Example 7** *The differential equation*

$$x^{\Delta\Delta\Delta\Delta}(t) + x(\sigma^2(t)) = 0,$$

for  $t \in \mathbb{T}$ , satisfies the hypotheses of most of the theorems in this paper. Note that if  $\mathbb{T} = \mathbb{R}^+$ , where  $\mathbb{R}^+$  is the set of nonnegative real numbers, then

$$\begin{aligned} x_1(t) &= e^{\frac{\sqrt{2}}{2}t} \cos\left(\frac{\sqrt{2}}{2}t\right), \\ x_2(t) &= e^{\frac{\sqrt{2}}{2}t} \sin\left(\frac{\sqrt{2}}{2}t\right), \\ x_3(t) &= e^{-\frac{\sqrt{2}}{2}t} \cos\left(\frac{\sqrt{2}}{2}t\right), \\ x_4(t) &= e^{-\frac{\sqrt{2}}{2}t} \sin\left(\frac{\sqrt{2}}{2}t\right), \end{aligned}$$

are solutions on  $\mathbb{R}^+$ . Note that  $x_1(t)$  and  $x_2(t)$  are unbounded near  $\infty$  and as guaranteed by Theorem 6 are oscillatory near  $\infty$ . It can be shown that

$$Fx_1(t) = Fx_2(t) = \frac{1}{\sqrt{2}}e^{\frac{t}{\sqrt{2}}},$$

and it follows that  $x_1(t)$  and  $x_2(t)$  are type II solutions. Also it can be shown that

$$Fx_3(t) = Fx_4(t) = -\frac{1}{\sqrt{2}}e^{-\frac{t}{\sqrt{2}}},$$

so  $x_3(t)$  and  $x_4(t)$  are type I solutions. Hence, the conjecture noted after Theorem 4 holds.

Note that Peil and Peterson [6] studied the asymptotic behavior of solutions of (1) with  $\mathbb{T} \equiv \mathbb{Z}$ . For fourth order difference equations see Smith and Taylor [8]. For fourth order ordinary differential equations see Jones [4] and Svec [9]. For results of the type in this paper for third order difference equations see Peterson [7]. For results for third order differential equations on a time scale see Morelli and Peterson [5].

## References

- [1] D. Anderson, A  $2n$ th-order Linear Difference Equation, Communications in Applied Analysis, 2, Number 4, (1998), 521-529.
- [2] C. D. Ahlbrandt and A. C. Peterson, The  $(n, n)$ -disconjugacy of a  $2n$ th-order linear difference equation, Computers Math. Applic., **28**, No 1-3(1994), 1-9.
- [3] P. Hartman, Difference Equations: Disconjugacy, Principal Solutions, Green's Functions, Complete Monotonicity, Transactions of the American Mathematical Society, **246**(December, 1978), 1-30.
- [4] G. Jones, Oscillatory solutions of a fourth order linear differential equation, Lecture Notes Pure Appl. Math. (S. Elaydi, ed.), Marcel Dekker 127, 261-266.
- [5] M. Morelli and A. Peterson A third order differential equation on a time scale (preprint).
- [6] T. Peil and A. Peterson, Asymptotic behavior of solutions of a two-term difference equation, Rocky Mountain Journal of Mathematics, **24**, No 1(Winter 1994), 233-251.
- [7] A. Peterson, A quadratic functional for a third order linear difference equation, Journal of Difference Equations, 3 (1998), 463-472.
- [8] B. Smith and W. Taylor, Oscillation and asymptotic behavior of certain fourth order difference equations, Rocky Mountain Journal Mathematics 16 (1986), 403-406.
- [9] M. Svec, Sur le comportement asymptotique des integrals de lequation differentielle  $y^{(4)} + Q(x)y = 0$ , Czech. Math. J. 8 (1958), 230-244.