

Name: _____ Score: _____

Instructions: You must show supporting work to receive full and partial credits.

1(20pts)

- (a) How many partitioning subintervals are needed over interval $[1, 2]$ in order to have an approximating Riemann sum within 0.01 of the exact definite integral $\int_1^2 e^{-x^2} dx$?

- (b) Approximate the integral $\int_1^2 e^{-x^2} dx$ by the left Riemann sum with the same partition number you found in (a).

- (c) Is the left sum approximation an under-estimate or an over-estimate of the true value? Explain your answer.

2(20pts)


(a) Use the definition of derivative to show that $(x^2)' = 2x$.

(b) The following table defines a function f at a few points. Approximate $f'(1.1)$, $f'(1.2)$, $f''(1.2)$ and fill in the blank boxes with your answers.

x	1.1	1.2	1.3
$f(x)$	1.21	1.44	1.69
$f'(x)$			
$f''(x)$			

(c) Label points A, B, C, D on the graph f with the following description

- (i) Point A is a point on the curve where the derivative is zero.
- (ii) Point B is a point on the curve where the function f is not differentiable.
- (iii) Point C is a point on the curve where the derivative is constant in an interval containing C .
- (iv) Point D is a point on the curve where the second derivative f'' is zero.



3(20pts) Find the derivative of

(a) $\ln(1 - \cos x)$

(b) $3^x + 3x^3$

(c) $e^{-2x} \sin x$

(d) $\frac{1 - x^2}{1 + x^2}$

4(20pts) In (a) and (b), find the exact value of the definite integrals by the Fundamental Theorem of Calculus

(a) $\int_0^1 4x^3 dx.$

(b) $\int_0^2 e^{2x} dx.$

(c) Given that $\int_1^2 f(x)dx = 2$, $\int_2^4 f(x)dx = -1$, $\int_1^2 g(x)dx = 0$, $\int_2^4 g(x)dx = 5$. Fill in the following blanks.

(i) $\int_1^4 f(x)dx = \underline{\hspace{2cm}}.$

(ii) $\int_1^2 [3f(x) - 2g(x)]dx = \underline{\hspace{2cm}}.$

(iii) $\int_1^3 f(x)dx + \int_3^4 f(x)dx = \underline{\hspace{2cm}}.$

(iv) $\int_4^2 f(x)dx = \underline{\hspace{2cm}}.$

5(20pts)

(a)

$$\text{Given } \begin{cases} f(1) = -1, g(1) = 4 \\ f'(1) = 3, g'(1) = 2 \\ f(2) = 3, g(2) = 1 \\ f'(2) = 0, g'(2) = -2 \end{cases}$$

$$\text{Find } \begin{cases} \text{(i) } h'(2), & \text{if } h(x) = 2f(x)g(x). \\ \text{(ii) } h'(2), & \text{if } h(x) = f(g(x)). \\ \text{(iii) } h'(1), & \text{if } h(x) = \frac{f(x)}{g(x)}. \\ \text{(iv) } h'(1), & \text{if } h(x) = f(g(x)/2). \end{cases}$$

(i)

(ii)

(iii)

(iv)

(b) An economist is interested in how the price of a certain item affects its sales. Suppose that at a price of $\$p$, a quantity, q , of the item is sold. If $q = f(p)$, explain the meaning of each of the following statements


(i) $f(150) = 2000$

(ii) $f'(150) = -25$

(i)

(ii)

(c) The temperature $T(t)$ in $^{\circ}\text{C}$ as a function of time t in minute is given by the graph. Estimate the average temperature over the time interval $[0, 60]$ and label it on the T -axis.



END

Sample Test 2 Solu Key

1 (a) $f(x) = e^{-x^2}$ monotone decreasing. Use regular partition of n subinterval. then $\Delta x = \frac{b-a}{n} = \frac{2-1}{n} = \frac{1}{n}$. And
 $\left| \int_1^2 e^{-x^2} dx - \text{Riemann sum of regular partition} \right| \leq |f(b) - f(a)| \Delta x = \frac{|e^{-4} - e^{-1}|}{n} \leq 0.01$
 $\Rightarrow n \geq \frac{e^{-1} - e^{-4}}{0.01} = 100(e^{-1} - e^{-4}) = 34.956. \Rightarrow \boxed{n=35}$

(b) $\int_1^2 e^{-x^2} dx \approx 0.14$

(c) Over-estimate because f is decreasing.




2 (a) $(x^2)' = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} = \lim_{h \rightarrow 0} \frac{(x+h-x)(x+h+x)}{h} = \lim_{h \rightarrow 0} \frac{h(2x+h)}{h} = \lim_{h \rightarrow 0} (2x+h) = 2x$

(b)

x	1.1	1.2	1.3
f	1.21	1.44	1.69
f'	2.3	2.5	
f''		2	

$f'(1.1) = \frac{f(1.2) - f(1.1)}{1.2 - 1.1} = \frac{1.44 - 1.21}{0.1} = \frac{0.23}{0.1} = 2.3$
 $f'(1.2) = \frac{f(1.3) - f(1.2)}{1.3 - 1.2} = \frac{1.69 - 1.44}{0.1} = \frac{0.25}{0.1} = 2.5$
 $f''(1.2) = \frac{f'(1.1) - f'(1.2)}{1.1 - 1.2} = \frac{2.3 - 2.5}{-0.1} = \frac{-0.2}{-0.1} = 2$

(c)



3 (a) $(\ln(1 - \cos x))' = \frac{1}{1 - \cos x} \cdot (1 - \cos x)' = \frac{1}{1 - \cos x} (0 - (-\sin x)) = \frac{\sin x}{1 - \cos x}$

(b) $(x^3 + 3x^2)' = (3x^2 \ln 3 + 6x)$

(c) $(e^{-2x} \sin x)' = (e^{-2x})' \sin x + e^{-2x} (\sin x)' = -2e^{-2x} \sin x + e^{-2x} \cos x$

(d) $\left(\frac{1-x^2}{1+x^2}\right)' = \frac{(1-x^2)'(1+x^2) - (1-x^2)(1+x^2)'}{(1+x^2)^2} = \frac{-2x(1+x^2) - 2x(1-x^2)}{(1+x^2)^2} = \frac{-4x}{(1+x^2)^2}$

4 (a) $\int_0^1 4x^3 dx = x^4 \Big|_0^1 = 1^4 - 0^4 = \textcircled{1}$ (b) $\int_0^2 e^{2x} dx = \frac{1}{2} e^{2x} \Big|_0^2 = \frac{1}{2} (e^4 - 1)$

(c) (i) $\int_1^4 f(x) dx = \int_1^2 f dx + \int_2^4 f dx = 2 + (-1) = \textcircled{1}$ (ii) $\int_1^2 (3f - 2g) dx$

$= 3 \int_1^2 f dx - 2 \int_1^2 g dx = 3(2) - 2(0) = \textcircled{6}$ (iii) $\int_1^3 f dx + \int_3^4 f dx = \int_1^4 f dx = \textcircled{1}$

(iv) $\int_4^2 f(x) dx = - \int_2^4 f(x) dx = -(-1) = \textcircled{1}$

5 (a) (i) $h'(2) = 2(f'(2)g'(2) + f(2)g''(2)) = 2(0(1) + 3(-2)) = \textcircled{-12}$

(ii) $h'(2) = f'(2)g'(2) = 3(-2) = \textcircled{-6}$

(iii) $h'(1) = \frac{f(1)g'(1) - f'(1)g(1)}{g^2(1)} = \frac{3(4) - (-1)(2)}{4^2} = \frac{14}{16} = \textcircled{\frac{7}{8}}$

(iv) $h'(1) = f'(1)g'(1)/2 = 3(2)/2 = \textcircled{3}$

(b) (i) 2000 items sold at unit price \$150 each (ii) For ~~each~~ \$1 increase in sale price, 25 items less are sold.

(c) $\bar{T} = \frac{1}{60} \int_0^{60} T(t) dt = \frac{1}{60} \left[\frac{10+30}{2} \cdot 20 + 30(20) + \frac{30}{2}(20) \right] = \textcircled{21.7}$