

Betti numbers for fat point ideals in the plane: a geometric approach.*

Alessandro Gimigliano
Dipartimento di Matematica e CIRM
Università di Bologna
40126 Bologna, Italy
email: gimiglia@dm.unibo.it

Brian Harbourne
Department of Mathematics
University of Nebraska
Lincoln, NE 68588-0130 USA
email: bharbour@math.unl.edu

Monica Idà
Dipartimento di Matematica
Università di Bologna
40126 Bologna, Italy
email: ida@dm.unibo.it

December 28, 2006

Acknowledgments: We thank GNSAGA, and the University of Bologna, which supported visits to Bologna by the second author, who also thanks the NSA and NSF for supporting his research.

Abstract

We consider the open problem of determining the graded Betti numbers for fat point subschemes Z supported at general points of \mathbf{P}^2 . We relate this problem to the open geometric problem of determining the splitting type of the pullback of $\Omega_{\mathbf{P}^2}$ to the normalization of certain rational plane curves. We give a conjecture for the graded Betti numbers which would determine them in all degrees but one for every fat point subscheme supported at general points of \mathbf{P}^2 . We also prove our Betti number conjecture in a broad range of cases. An appendix discusses many more cases in which our conjecture has been verified computationally and provides a new and more efficient computational approach for computing graded Betti numbers in certain degrees. It also demonstrates how to derive explicit conjectural values for the Betti numbers and how to compute splitting types.

1 Introduction

Let $Z = m_1P_1 + \cdots + m_nP_n$ be a fat point subscheme of \mathbf{P}^2 supported at general points P_i . Thus Z is the 0-dimensional subscheme of \mathbf{P}^2 defined by the homogeneous ideal $I(Z) = \cap_i I(P_i)^{m_i}$ in the homogeneous coordinate ring $R = K[\mathbf{P}^2]$ of \mathbf{P}^2 (where we take K to be an algebraically closed field of arbitrary characteristic), where $I(P_i)$ is the ideal generated by all homogeneous forms $f \in R$ vanishing at P_i . The homogeneous component $I(Z)_t$ of $I(Z)$ in degree t is just the K -vector space span of the homogeneous elements of $I(Z)$ of degree t . Thus $I(Z)_t$ consists of all homogeneous polynomials of degree t which vanish at each point P_i to order at least m_i . The Hilbert function of $I(Z)$ is the function h_Z which gives the K -vector space dimension $h_Z(t) = \dim I(Z)_t$ of $I(Z)_t$ as a function of t .

*2000 *Mathematics Subject Classification*. Primary 14C20, 13P10; Secondary 14J26, 14J60.

Key words and phrases. Graded Betti numbers, fat points, splitting types.

1.1 The SHGH Conjecture

The problem of determining h_Z has attracted a lot of attention over the years, and it is still an open problem in general. In fact, even the most fundamental problem is open: it is not known in general what the least t is (which we call $\alpha(Z)$) for which $I(Z)_t \neq 0$. The fact that even the latter problem is open is less surprising given that if one knows $\alpha(Z)$ for every Z , then one can determine $h_Z(t)$ for all t and Z , and conversely.

Given that it is not known how big $I(Z)_t$ is, or even if it is non-zero, it is not surprising that the least value of t such that $I(Z)$ is generated in degrees t or less is not known. More precisely, the graded Betti numbers for the minimal free resolution of $I(Z)$ are not known. There is not even a conjecture in general for the Betti numbers. On the other hand, there is a general conjecture for the values of the Hilbert function h_Z . This is the SHGH Conjecture (due, in various equivalent forms, to Segre [S], Harbourne [Ha4], Gimigliano [G] and Hirschowitz [Hi1]; see Conjecture 2.2.1 or Conjecture A1.2.1). Considerations of geometry lead to a lower bound $e(h_Z, t)$ for $h_Z(t)$ (the definition of $e(h_Z, t)$, which is somewhat complicated, is given in the appendix). The SHGH Conjecture asserts that $e(h_Z, t) = h_Z(t)$. Thus $e(h_Z, t)$ is regarded as the “expected” value of $h_Z(t)$.

In degrees $t > \alpha(Z)$, the SHGH Conjecture implies the following simple statement:

Conjecture 1.1.1 *Let $Z = m_1P_1 + \cdots + m_nP_n$, for nonnegative integers m_i and general points $P_i \in \mathbf{P}^2$. If $t > \alpha(Z)$ and $f_1^{b_1} \cdots f_r^{b_r}$ is a factorization of the greatest common divisor of $I(Z)_t$ as a product of non-associate irreducible factors, then*

$$h_Z(t) = \binom{t+2}{2} - \sum_i \binom{m_i+1}{2} + \sum_l \binom{b_l}{2}.$$

In particular, under the assumptions given, $e(h_Z, t) = \binom{t+2}{2} - \sum_i \binom{m_i+1}{2} + \sum_l \binom{b_l}{2}$. Although it is perhaps not clear from this how to actually compute $e(h_Z, t)$, the full SHGH Conjecture allows one to compute $e(h_Z, t)$ in terms only of the m_i , and also to conjecturally determine the degrees and multiplicities at each point P_i of all curves in the base locus of $I(Z)_t$ (see the appendix). The advantage of the statement above is that it gives insight into the problem of determining h_Z , without the burden of the technicalities needed to state the full SHGH Conjecture.

For example, $I(Z)_t$ consists of all homogeneous polynomials of degree t which vanish at each point P_i to order at least m_i . But the vector space of all forms of degree t has dimension $\binom{t+2}{2}$, and requiring vanishing at P_i to order m_i imposes $\binom{m_i+1}{2}$ independent linear conditions. We do not know that the conditions imposed at one point are independent of those imposed at all of the other points, and in fact they are not always independent. Thus we obtain the bound $h_Z(t) \geq \binom{t+2}{2} - \sum_i \binom{m_i+1}{2}$. What the SHGH Conjecture does in essence is to give a precise measure of the failure of the imposed conditions to be independent. The key insight is that the failure of independence in degree $t > \alpha(Z)$ is due to the gcd not being square-free. What the conjecture above hides is that the curves defined by the forms f_i are thought always to be very special, and this is significant for what we do in this paper.

To make this clearer we strengthen the conjecture above to include the case that $t = \alpha(Z)$. To do this we need a minor technicality. We say a plane curve C defined by an irreducible form of degree d is *contributory* with respect to points P_1, \dots, P_n if C is a rational curve smooth except possibly at the points P_i , such that $\text{mult}_{P_1}(C) + \cdots + \text{mult}_{P_n}(C) = 3d - 1$. We will say a plane curve C defined by an irreducible form of degree d is *negative* with respect to points P_1, \dots, P_n if $(\text{mult}_{P_1}(C))^2 + \cdots + (\text{mult}_{P_n}(C))^2 > d^2$. It follows by the genus formula that curves contributory for P_1, \dots, P_n are also negative.

After extending Conjecture 1.1.1, we have:

Conjecture 1.1.2 *Let $Z = m_1P_1 + \cdots + m_nP_n$, for nonnegative integers m_i and general points $P_i \in \mathbf{P}^2$. Given $t \geq \alpha(Z)$, let $f_1^{c_1} \cdots f_r^{c_r}$ be a factorization of the greatest common divisor of $I(Z)_t$ as a product of non-associate irreducible factors. Then every factor f_j which defines a curve negative for the points P_i defines a contributory curve, and we have*

$$h_Z(t) = \binom{t+2}{2} - \sum_i \binom{m_i+1}{2} + \sum_k \binom{c_{j_k}}{2},$$

where the index k runs over the factors f_{j_k} defining curves contributory for the points P_i .

The point is that the SHGH Conjecture implies that it is only exponents c_j of factors f_j defining contributory curves which contribute to the lack of independence. The SHGH Conjecture also implies in degrees $t > \alpha(Z)$ that every factor f_j defines a contributory curve, and that every curve negative for general points is in fact contributory. (Factors defining non-contributory curves do occur in degree α , however; for example, if $Z = P_1 + \cdots + P_9$ consists of 9 general points, then $\alpha(Z) = 3$ and the gcd in degree 3 is an irreducible cubic defining an elliptic curve C , which is therefore not contributory, but it is also not negative.)

It is known, however, that contributory curves do contribute to the failure of independence. The SHGH Conjecture is that nothing else contributes. In particular, if the greatest common divisor of $I(Z)_t$ is 1 (i.e., if the zero locus of $I(Z)_t$ is at most 0-dimensional), or more generally if the gcd of $I(Z)_t$ is just square-free, then the conjecture is that there is no failure of independence, and hence that $h_Z(t) = \binom{t+2}{2} - \sum_i \binom{m_i+1}{2}$.

1.2 The Betti Number Conjecture

The goal of this paper is to give a theorem and a conjecture for graded Betti numbers for ideals $I(Z)$, mimicking the fact that there is a bound $e(h_Z, t) \leq h_Z(t)$ which by the SHGH Conjecture is an equality, given explicitly by Conjecture 1.1.2.

The SHGH Conjecture specifies how big the ideal $I(Z)$ is in each degree. The next question, for which no general conjecture has yet been posed, is where must one look for generators of $I(Z)$. More precisely, in any minimal set of homogeneous generators, how many generators are there in each degree? The goal of this paper is to develop (and prove cases of) a conjecture for the numbers of generators in each degree bigger than $\alpha(Z) + 1$. The number of generators in degrees less than $\alpha(Z) + 1$ is trivial: there are obviously no generators in degrees $t < \alpha(Z)$, and, since any minimal set of homogeneous generators of $I(Z)$ must include a K -vector space basis of $I(Z)_\alpha$, precisely $h_Z(t)$ generators in degree $t = \alpha(Z)$. There remains the question of how many generators there are in degree $t = \alpha(Z) + 1$. Our approach is to relate the number of generators in a given degree to the splitting of a certain rank 2 bundle on certain curves. In degree $\alpha(Z) + 1$ precisely what curves must be taken into account is more subtle than it is in larger degrees. Thus here we focus on degrees larger than $\alpha(Z) + 1$. We study the subtleties needed for a unified approach that subsumes degree $\alpha(Z) + 1$ in separate papers, beginning with [GHI1].

For any t , the number of homogeneous generators in degree $t + 1$ in any minimal set of homogeneous generators is the dimension of the cokernel of the map $\mu_t : I(Z)_t \otimes R_1 \rightarrow I(Z)_{t+1}$ given on simple tensors by multiplication, $f \otimes g \mapsto fg$. What we will do is to give a conjecture for $\dim \text{cok } \mu_t$ for all Z in all degrees $t > \alpha(Z)$. Clearly $\dim \text{cok } \mu_t = 0$ if $t < \alpha(Z) - 1$ and $\dim \text{cok } \mu_t = h_Z(\alpha(Z))$ if $t = \alpha(Z) - 1$, so our conjecture handles all cases except $t = \alpha(Z)$. The conjecture is in terms of data determined by contributory curves.

Let C' be a curve contributory for points P_1, \dots, P_n . The composition of the normalization map $C \rightarrow C'$ with the inclusion $C' \subset \mathbf{P}^2$ gives a morphism $f : C \rightarrow \mathbf{P}^2$. Pulling back the twisted cotangent bundle $\Omega_{\mathbf{P}^2}(1)$ gives a rank two bundle $f^*(\Omega_{\mathbf{P}^2}(1))$ on C , but C is smooth and rational, so $f^*(\Omega_{\mathbf{P}^2}(1))$ splits as $f^*(\Omega_{\mathbf{P}^2}(1)) \cong \mathcal{O}_C(-a_C) \oplus \mathcal{O}_C(-b_C)$ for some integers $a_C \leq b_C$. We call (a_C, b_C) the *splitting type* of C (and for convenience, we set $a_{C'} = a_C$ and $b_{C'} = b_C$ and refer to $(a_{C'}, b_{C'})$ also as the *splitting type* of C').

Given $t > \alpha(Z)$, clearly $I(Z)_{t-1} \neq 0$, so $I(Z)_{t-1}$ has a well-defined gcd (up to scalar multiple). Let γ_{t-1} be the product of all irreducible factors of the gcd defining curves negative for the points P_1, \dots, P_n . (If $t > \alpha(Z) + 1$, then, as in Conjecture 1.1.1, the SHGH Conjecture implies that γ_{t-1} is itself the gcd.) Let $\gamma_{t-1} = f_1^{c_1} \cdots f_r^{c_r}$ be its factorization into non-associate irreducible factors and let $d_j = \deg(f_j)$. Do likewise for $I(Z)_t$; it is easy to see that we get $\gamma_t = f_1^{c'_1} \cdots f_r^{c'_r}$ where $c'_j \leq c_j$ for all j . Doing the same for $I(Z)_{t+1}$ gives $\gamma_{t+1} = f_1^{c''_1} \cdots f_r^{c''_r}$ with $c''_j \leq c'_j$. Let C_j be the normalization of the curve defined by f_j . The following theorem refines ideas of Fitchett [F1], [F2].

Theorem 1.2.1 *Let $Z = m_1 P_1 + \cdots + m_n P_n$, for nonnegative integers m_i and general points $P_i \in \mathbf{P}^2$. Let $t > \alpha(Z)$ with c_j, c'_j and c''_j defined as above. If the SHGH Conjecture holds, then*

$$\dim \text{cok } \mu_t \leq \sum_j d_j (c'_j - c''_j) - \sum_j \binom{c'_j - c''_j}{2} + \sum_j \left(\binom{c_j - c'_j - a_{C_j}}{2} + \binom{c_j - c'_j - b_{C_j}}{2} \right).$$

In fact, the full SHGH Conjecture is not required for this theorem. One just needs that $I(Z)$ behaves as expected for the specific Z being considered. We also propose the following conjecture:

Conjecture 1.2.2 *Equality holds in Theorem 1.2.1.*

As we show by examples in the appendix, the SHGH Conjecture allows one to determine conjectural values for the exponents c_j, c'_j and c''_j . As we show below, in many cases the splitting type of the curves C_j are also known, and even in those cases where the type is not known, it is much easier to compute the splitting type of each C_j symbolically than it is to compute $\dim \text{cok } \mu_t$ symbolically in the usual way of finding a Gröbner basis of the ideal $I(Z)$. Thus the SHGH Conjecture and Conjecture 1.2.2, if true, allow one to determine the minimal number of homogeneous generators of $I(Z)$ in every degree except possibly degree $\alpha(Z) + 1$. Examples in the appendix show how this is done.

We now put this into the context of minimal free graded resolutions. The minimal free graded resolution of $I(Z)$ is an exact sequence of the form $0 \rightarrow M_1 \rightarrow M_0 \rightarrow I(Z) \rightarrow 0$, where M_0 (the module of generators) and M_1 (the module of syzygies) are free graded R -modules, hence of the form $M_0 = \bigoplus_{i \geq 0} R[-i]^{g_i(Z)}$, and $M_1 = \bigoplus_{j \geq 0} R[-j]^{s_j(Z)}$ for nonnegative integers $g_i(Z)$ and $s_j(Z)$. (By $R[-i]$ we just mean the free R -module of rank 1 with the grading such that $R[-i]_t = R_{t-i}$.) The graded Betti numbers of $I(Z)$ are the sequences of integers $g_i(Z)$ and $s_j(Z)$ (which we write as g_i and s_j if Z is understood). The Betti number g_i is just $\dim \text{cok } \mu_{i-1}$, hence g_i is the number of generators of degree i in any minimal set of homogeneous generators of $I(Z)$. Moreover, it is not hard to show that $g_i - s_i = \Delta^3 h_Z(i)$ for all i , where Δ is the difference operator; i.e., $\Delta h_Z(i) = h_Z(i) - h_Z(i-1)$ (see p. 685 of [FHH]).

1.3 The Structure of The Paper

We obtain our results by reformulating them in terms of complete linear systems on the surface X obtained by blowing up \mathbf{P}^2 at the points P_1, \dots, P_n .

In Section 2 we recall the background necessary for this reformulation, and we state known results needed for our approach. In Section 3 we state our main results and show how they lead to the statement in terms of fat points given above.

We also include an appendix for the purpose of showing how to obtain explicit predictions for the values of Hilbert functions of fat points and how our results lead to explicit numerical predictions for Betti numbers. In addition, Section A2 of the appendix discusses how to compute splitting types and discusses evidence in support of Conjecture 1.2.2, partly based on an approach for computing g_i for $i > \alpha + 1$ which is substantially more efficient than the usual methods, which involve finding a Gröbner basis of $I(Z)$. (A Macaulay 2 script which implements our method is included in Section A2.3.) Even for relatively small values of the multiplicities m_i and even for randomly chosen points P_i over a finite field rather than for generic P_i , finding the graded Betti numbers g_i for $I(Z)$ is beyond what can be done computationally by the usual methods. For example, we were unable to determine the graded Betti numbers of $I(Z)$ for $Z = 77P_1 + \cdots + 77P_7 + 44P_8 + 11P_9 + 11P_{10} + 11P_{11}$ by the usual methods, but our new computational method, based on the results of Section 3, working on an 800MHz computer over the finite field $|K| = 31991$ using randomly chosen points P_i , determined the result in slightly over 5 minutes. The result, of course, is in agreement with our conjectural expected values. See also Example A1.2.3.

1.4 What was Previously Known

A lot of work has been done on the SHGH Conjecture. That the SHGH Conjecture holds for $n \leq 9$ points was known to Castelnuovo [Cas]; a more modern proof is given by Nagata [N2]. The uniform case (i.e., $Z = m(P_1 + \cdots + P_n)$) was proved for $m = 2$ by [AC], $m = 3$ by [Hi2], for $m \leq 12$ by [CM2] and for $m \leq 20$ by [CCMO]. The case that n is a square has likewise seen progressive improvements, with the main difficulty being to show $I(Z)_t = 0$ when it is expected to be. For example, by specializing $n = 16$ points to a smooth curve of degree 4, it is not hard to show that $h_Z(t)$ has its expected value of $\binom{t+2}{2} - 4\binom{m+1}{2}$ for all $t \geq 4m + 1$, while $h_Z(t)$ has its expected value of 0 for $t < 4m + 1$ by [N1] for all m . A generalization of this in [HHF] shows that the SHGH Conjecture holds in all degrees t such that $e(h_z, t) > 0$, if n is any square as long as m is not too small. By more technical arguments one can show that the SHGH Conjecture holds also for small m and t . For example, if n is a power of 4, [E1] showed SHGH holds in the uniform case; [BZ] extended this to n being a product of powers of 4 and 9; and [HR] showed that the SHGH Conjecture holds for infinitely many m for each square n . By [E2] it is now known to hold in the uniform case for any m when n is a square; alternate proofs have been given by [CM3] and [R].

Results for the general case (i.e., such that the multiplicities m_i of $Z = \sum_i m_i P_i$ need not all be equal) are not as comprehensive. That the SHGH Conjecture holds in the case that $m_i \leq 3$ for all i is due to [CM1], improved to $m_i \leq 4$ by [Mi] and then to $m_i \leq 7$ by [Y].

Previous results on graded Betti numbers seem to start with [Cat], which obtained a complete answer for $n \leq 5$ general points. This was extended to $n = 6$ by [F3], then 7 by [Ha1] and 8 by [FHH]. For $n > 8$, almost all results (and even conjectures) are for cases which are either uniform or close to uniform. For example, conjectures in the uniform case were put forward by [Ha2], and in cases close to uniform by [HHF]. (Those conjectures are consistent with Conjecture 1.2.2 due to Corollary 2.3.2.) The Betti numbers in the case of n general points of multiplicity $m = 1$ were determined by [GGR]. The uniform Betti numbers conjecture of [Ha2] was verified for $m = 2$ by [I] and for $m = 3$ by [GI]. More generally, if $Z = m_1 P_1 + \cdots + m_n P_n$ where the points $P_i \in \mathbf{P}^2$ are general and $m_i \leq 3$ for all i , [BI] determines the graded Betti numbers in all degrees. By Theorem 3.2 of [HHF], by applying [E2] it follows that the uniform Betti numbers conjecture of [Ha2] holds for all $m \geq (\sqrt{n} - 2)/4$ when the number n of points is an even square. Additional cases are shown

in [HR] when n is not a square.

2 Background

In this section we set notation and cite well known facts which we will refer to later in the paper. Let P_1, \dots, P_n be distinct (not necessarily general) points of the projective plane \mathbf{P}^2 . Let $p : X \rightarrow \mathbf{P}^2$ be the birational morphism given by blowing up the points.

2.1 Preliminaries

The divisor class group $\text{Cl}(X)$ of divisors on X modulo linear equivalence is the free abelian group with basis L, E_1, \dots, E_n , where E_i is the class of the divisor $p^{-1}(P_i)$ and L is the pullback of the class of a line. Given any divisor F on X , the dimension $h^0(X, \mathcal{O}_X(F))$ of the global sections of $\mathcal{O}_X(F)$ depends only on the class $[F]$ of F . For convenience, we will denote $h^0(X, \mathcal{O}_X(F))$ by either $h^0(X, F)$ or $h^0(X, [F])$, or even $h^0(F)$ or $h^0([F])$ if X is understood.

Given any $F = tL - m_1E_1 - \dots - m_nE_n$, by Riemann-Roch we have

$$h^0(X, F) - h^1(X, F) + h^2(F, X) = \frac{F^2 - K_X \cdot F}{2} + 1,$$

where $K_X = -3L + E_1 + \dots + E_n$ is the canonical class. Since E_i is reduced and irreducible and $m_i = E_i \cdot F$, we have a canonical isomorphism $H^0(X, tL - \sum_{m_i > 0} m_i E_i) \rightarrow H^0(X, F)$. (The idea is that if $|F|$ is nonempty, then $-\sum_{m_i < 0} m_i E_i$ is contained in the base locus of $|F|$, essentially by Bezout's Theorem, and so $|F| = (-\sum_{m_i < 0} m_i E_i) + |tL - \sum_{m_i > 0} m_i E_i|$.) On the other hand, L is nef (meaning that $L \cdot C \geq 0$ for any effective divisor C on X), so $h^0(X, F) = 0$ if $t < 0$. By duality we have $h^2(X, F) = h^0(X, K_X - F)$, so it follows that $h^2(X, F) = 0$ whenever $t \geq 0$ (in fact, whenever $t \geq -2$).

If $t \geq 0$ and $m_i \geq 0$ for all i , then $\frac{F^2 - K_X \cdot F}{2} + 1 = \binom{t+2}{2} - \sum_i \binom{m_i+1}{2}$, so Riemann-Roch gives

$$h^0(X, F) \geq \max\left(0, \binom{t+2}{2} - \sum_i \binom{m_i+1}{2}\right).$$

This is just a manifestation of the canonical identification $H^0(X, tL - m_1E_1 - \dots - m_nE_n) = H^0(\mathbf{P}^2, \mathcal{I}_Z(t)) = I(Z)_t$, where $Z = m_1P_1 + \dots + m_nP_n$ and \mathcal{I}_Z is the sheaf of ideals defining Z . Because of this, given any integer t and a fat point scheme $Z = m_1P_1 + \dots + m_nP_n$, we define $F_t(Z) = tL - m_1E_1 - \dots - m_nE_n$, and hence we have $h_Z(t) = h^0(X, F_t(Z))$ for all t .

Given a divisor F on X , let $\mu_F : H^0(X, F) \otimes H^0(X, L) \rightarrow H^0(X, F + L)$ denote the obvious natural map. By identifying $H^0(X, F_t(Z))$ with $I(Z)_t$ and $H^0(X, L)$ with $H^0(\mathbf{P}^2, \mathcal{O}_{\mathbf{P}^2}(1)) = R_1$, it follows that $\dim \text{cok } \mu_t(Z) = \dim \text{cok } \mu_{F_t(Z)}$. Given a sum $F = H + N$ of effective divisors such that $|F| = N + |H|$ (i.e., such that N is contained in the scheme theoretic base locus of F), Lemma 2.1.1 gives a simple but useful fact which relates $\dim \text{cok } \mu_F$ to $\dim \text{cok } \mu_H$. (For the proof, observe that N is in the base locus of the image $\text{Im } \mu_F$ of μ_F ; i.e., that $\text{Im } \mu_F = N + \text{Im } \mu_H$. See Lemma 2.10(b) of [Ha3].) Obviously μ_F is injective when F is not effective, and, when F is effective, F decomposes as in Lemma 2.1.1. Thus Lemma 2.1.1 reduces the general problem of computing $\dim \text{cok } \mu_F$ to the case that $|F|$ is effective and fixed component free, and thus in particular to the case that F is nef.

Lemma 2.1.1 *Let $F = H + N$ be a sum of effective divisors H and N on the surface X such that $|F| = N + |H|$. Then*

$$\dim \text{cok } \mu_F = (\dim \text{cok } \mu_H) + (h^0(X, F + L) - h^0(X, H + L)).$$

It is easy to give examples such that $F = N$ (and hence μ_F is injective) but that $N + |L|$ is a proper subset of $|F + L|$ (and hence the map μ_F cannot be surjective). For example, if C is an *exceptional* curve (i.e., C is smooth and rational with $C^2 = -1$) with $d = C \cdot L$, then the kernel of $\mu_{L+(d+1)C}$ is non-zero since already μ_L has non-zero kernel, and $\mu_{L+(d+1)C}$ is not onto since C is in the base locus of $|L + (d+1)C|$ but $|2L + (d+1)C|$ is base curve free. In particular, the occurrence of fixed components is one reason that μ_F can fail to have maximal rank (i.e., fail to be either injective or surjective). What motivates this paper is that this is not the only reason. In fact, μ_F can fail to have maximal rank even when F is very ample. (For example, let $F = (3L - E_1 - \cdots - E_7) + m(8L - 3E_1 - \cdots - 3E_7 - E_8)$, where the points P_i are general and $m \geq 1$. Then F is very ample [Ha5], but μ_F fails to have maximal rank [FHH].) The point of this paper is that, when $F - L$ is effective, the failure of μ_F to have maximal rank depends on the fixed components of $|F - L|$.

Note that if C is a plane curve contributory for points P_i , then C' is an exceptional curve, where C' is the proper transform of C on the surface X obtained by blowing up the points P_i . As discussed in the introduction, our geometric approach for determining the dimension of the cokernel of μ_F for certain divisors F thus depends on knowing the splitting $\mathcal{O}_E(-a_E) \oplus \mathcal{O}_E(-b_E)$ of the restriction $p^*\Omega(1)|_E$ of $p^*\Omega(1)$ to exceptional curves E . The following result (see [As], or [F1], [F2]) covers most of what is known:

Lemma 2.1.2 *Let $E \subset X$ be a smooth rational curve, where $p : X \rightarrow \mathbf{P}^2$ is the morphism blowing up distinct points P_i of \mathbf{P}^2 . Let $d = E \cdot L$ and let m be the maximum of $E \cdot E_i$, $1 \leq i \leq n$. Then there are integers $0 \leq a_E \leq b_E \leq d$ with $\min(m, d - m) \leq a_E \leq d - m$ and $d = a_E + b_E$ such that $(p^*\Omega_{\mathbf{P}^2}(1))|_E$ is isomorphic to $\mathcal{O}_E(-a_E) \oplus \mathcal{O}_E(-b_E)$.*

Note that, if $d \leq 2m + 1$, then $a_E = \min(m, d - m)$ and $b_E = \max(m, d - m)$. For cases not covered by Lemma 2.1.2, a_E and b_E can be computed fairly efficiently. We will describe an algorithm for doing so in section A2 of the appendix.

2.2 The SHGH Conjecture

Here we state the version of the SHGH Conjecture given in [Ha4]. This version is simple to state and useful conceptually. We include in the appendix an equivalent version that is more useful for obtaining explicit conjectural values of Hilbert functions.

Conjecture 2.2.1 *Let X be a surface obtained by blowing up n generic points of \mathbf{P}^2 . Then every reduced irreducible curve $C \subset X$ with $C^2 < 0$ is an exceptional curve and either $h^0(X, F) = 0$ or $h^1(X, F) = 0$ for every nef divisor F on X .*

Theorem 2.2.2 *Conjecture 2.2.1 implies Conjectures 1.1.1 and 1.1.2.*

Proof. First, note that if H is nef and effective, then $h^2(X, H) = 0$ by duality (since L is nef but $(K_X - H) \cdot L < 0$). Now let C be nef and effective. Then $H + C$ is also nef and effective. By Conjecture 2.2.1, $h^1(X, H) = 0 = h^1(X, H + C)$ and $h^1(X, C) = 0 = h^2(X, C)$ (hence $(C^2 - C \cdot K_X)/2 \geq 0$ by Riemann-Roch). Assume $H \cdot C > 0$. Applying Riemann-Roch for surfaces now gives $h^0(X, H + C) = h^0(X, H) + (C^2 - C \cdot K_X)/2 + H \cdot C > h^0(X, H)$. Similarly, if C is instead an exceptional curve with $C \cdot H > 0$, then $H + C$ is nef and effective, and $h^0(X, H + C) > h^0(X, H)$. In particular, if H is nef and effective with $|H|$ base curve free, and if C is a prime divisor which is a base curve of $|H + C|$ such that C is either exceptional or $C^2 > 0$ (hence nef), then $C \cdot H = 0$.

Now consider $Z = m_1P_1 + \cdots + m_nP_n$ and let $\alpha = \alpha(Z)$. Then $h_Z(t) = h^0(X, F)$, where $F = tL - m_1E_1 - \cdots - m_nE_n$. Let H be the moving part of $|F|$, and decompose the fixed part as $D + N$, where D is the sum of the curves in the base locus of $|F|$ of nonnegative self-intersection and N is the sum of the curves in the base locus of negative self-intersection. By Conjecture 2.2.1, each curve in N is an exceptional curve, hence disjoint from H and D . In addition, these exceptional curves are pairwise orthogonal (since if C and C' both appear in N and have $C \cdot C' > 0$, then $h^0(X, C + C') \geq C \cdot C' + 1$ by Riemann-Roch, hence $C + C'$ cannot be part of the base locus of $|F|$). Also, none of the exceptional curves E appearing in N is the blow up of a point P_i (since if it were we would have $m_i = E \cdot F = E \cdot (H + D + N) = E \cdot N < 0$). Thus for some c_{j_k} we have $N = \sum_k c_{j_k} C_{j_k}$, where C_j is the proper transform of the plane curve defined by f_j in the statement of Conjecture 1.1.2. Applying Riemann-Roch gives

$$\begin{aligned} h_Z(t) &= h^0(X, F) = h^0(X, F - N) = ((F - N)^2 - K_X \cdot (F - N))/2 + 1 \\ &= (F^2 - K_X F)/2 + 1 + \sum_k \binom{c_{j_k}}{2} = \binom{t+2}{2} - \sum_i \binom{m_i+1}{2} + \sum_k \binom{c_{j_k}}{2}, \end{aligned}$$

as claimed in Conjecture 1.1.2.

Now consider degree $t + 1$, so $t + 1 > \alpha(Z)$. Then $h_Z(t + 1) = h^0(X, F + L)$, where $F + L = H + (L + D) + N$. Note that, as we saw above, $|L + D|$ is fixed component free. Thus the fixed part of $|F + L|$ consists at most of exceptional curves coming from $N = \sum_k c_{j_k} C_{j_k}$. In fact, by the first paragraph of the proof, $|F + D + L + \sum_k \min(d_{j_k}, c_{j_k}) C_{j_k}|$ is base curve free and $|F + D + L + N| = |F + D + L + \sum_k \min(d_{j_k}, c_{j_k}) C_{j_k}| + \sum_k \max(0, c_{j_k} - d_{j_k}) C_{j_k}$. I.e., $\sum_k \max(0, c_{j_k} - d_{j_k}) C_{j_k}$ is the divisorial part of the base locus of $|F + L|$. If we denote $\max(0, c_{j_k} - d_{j_k})$ by b_{j_k} and reindex, this becomes $\sum_l b_l C_l$, and applying Riemann-Roch as above gives

$$h_Z(t + 1) = h^0(X, F + L) = \binom{(t + 1) + 2}{2} - \sum_i \binom{m_i + 1}{2} + \sum_l \binom{b_l}{2},$$

as claimed in Conjecture 1.1.1. (Note that $t + 1$ here is the same as t in the statement of Conjecture 1.1.1, since there we assumed $t > \alpha(Z)$, but here, in order to handle Conjecture 1.1.2 simultaneously, we assumed only $t \geq \alpha(Z)$.) \blacksquare

2.3 Mumford's Snake Lemma

Mumford [Mu1] applied the snake lemma to questions related to μ_F . We recall that now. To do so we establish some notation that we will use here and throughout the paper. Let F , C and D be divisors on X with C effective; then we have the natural multiplication maps

$$\mu_{F,D} : H^0(X, F) \otimes H^0(X, D) \rightarrow H^0(X, F + D);$$

and

$$\mu_{C;F,D} : H^0(C, F|_C) \otimes H^0(X, D) \rightarrow H^0(C, (F + D)|_C).$$

In the particular case that $D = L$, which is almost always true in the present paper, we write μ_F and $\mu_{C;F}$ instead of $\mu_{F,L}$ and $\mu_{C;F,L}$.

Lemma 2.3.1 *Let $p : X \rightarrow \mathbf{P}^2$ be a blow up of \mathbf{P}^2 at n distinct points with L, E_1, \dots, E_n as usual. Let D be a divisor on X , let $V = H^0(X, D)$, and let F and C be divisors on X with C effective and with $h^1(X, F) = 0 = h^1(X, F + D)$. Then the following diagram is commutative with exact rows:*

$$\begin{array}{ccccccc} 0 & \rightarrow & H^0(X, F) \otimes V & \rightarrow & H^0(X, F + C) \otimes V & \rightarrow & H^0(C, (F + C)|_C) \otimes V & \rightarrow & 0 \\ & & \downarrow \mu_{F,D} & & \downarrow \mu_{F+C,D} & & \downarrow \mu_{C;F+C,D} & & (\circ) \\ 0 & \rightarrow & H^0(X, F + D) & \rightarrow & H^0(X, F + C + D) & \rightarrow & H^0(C, (F + C + D)|_C) & \rightarrow & 0 \end{array}$$

The snake lemma thus gives an exact sequence

$$\begin{aligned} 0 &\rightarrow \ker \mu_{F,D} \rightarrow \ker \mu_{F+C,D} \rightarrow \ker \mu_{C;F+C,D} \\ &\rightarrow \operatorname{cok} \mu_{F,D} \rightarrow \operatorname{cok} \mu_{F+C,D} \rightarrow \operatorname{cok} \mu_{C;F+C,D} \rightarrow 0 \end{aligned}$$

which we will refer to as $\mathcal{S}(F, C, D)$ (or $\mathcal{S}(F, C)$ if $D = L$).

Another useful fact is the Castelnuovo-Mumford Lemma [Mu2], which gives a criterion for μ_F to not only have maximal rank but to be surjective. The version we state, Corollary 2.3.2, follows easily from Lemma 2.3.1, using $\mathcal{S}(0, L, H)$. (The hypothesis $h^1(X, H - L) = 0$ is used to ensure that $H^0(X, H) \rightarrow H^0(L, H|_L)$ is surjective; $H \cdot L \geq 0$ then ensures that $\operatorname{cok} \mu_{L;L,H} = 0$.)

Corollary 2.3.2 *Let $p : X \rightarrow \mathbf{P}^2$ be obtained by blowing up n distinct points of \mathbf{P}^2 , with L the pullback of the class of a line. If H is a divisor on X with $h^1(X, H - L) = 0$ and $H \cdot L \geq 0$, then $\operatorname{cok} \mu_H = 0$.*

When $D = L$, explicit expressions for the kernels and cokernels in Lemma 2.3.1 can be given in terms of the cotangent bundle. Recall the Euler sequence defining $\Omega = \Omega_{\mathbf{P}^2}$:

$$0 \rightarrow \Omega(1) \rightarrow \mathcal{O}_{\mathbf{P}^2} \otimes H^0(\mathbf{P}^2, \mathcal{O}_{\mathbf{P}^2}(1)) \rightarrow \mathcal{O}_{\mathbf{P}^2}(1) \rightarrow 0.$$

Pulling the Euler sequence back to X , tensoring by $\mathcal{O}_X(F)$ and identifying $H^0(\mathbf{P}^2, \mathcal{O}_{\mathbf{P}^2}(1))$ with $H^0(X, L)$ gives an exact sequence

$$0 \rightarrow (p^*\Omega)(F + L) \rightarrow \mathcal{O}_X(F) \otimes H^0(X, L) \rightarrow \mathcal{O}_X(F + L) \rightarrow 0. \quad (\dagger)$$

If $h^1(X, F) = 0$, then taking cohomology gives an exact sequence

$$0 \rightarrow H^0(X, (p^*\Omega)(F + L)) \rightarrow H^0(X, F) \otimes H^0(X, L) \rightarrow H^0(X, F + L) \rightarrow H^1(X, (p^*\Omega)(F + L)) \rightarrow 0,$$

hence $\ker \mu_F = H^0(X, (p^*\Omega)(F + L))$ and $\operatorname{cok} \mu_F = H^1(X, (p^*\Omega)(F + L))$. Similarly, if $h^1(C, F|_C) = 0$, by restricting (\dagger) to C and taking cohomology we see $\ker \mu_{C;F} = H^0(C, ((p^*\Omega)(F + L))|_C)$ and $\operatorname{cok} \mu_{C;F} = H^1(C, ((p^*\Omega)(F + L))|_C)$. In case C is a smooth rational curve, taking $t = F \cdot C$ we have:

$$\begin{aligned} \ker \mu_{C;F} &= H^0(C, \mathcal{O}_C(t - a_C) \oplus \mathcal{O}_C(t - b_C)) \\ \operatorname{cok} \mu_{C;F} &= H^1(C, \mathcal{O}_C(t - a_C) \oplus \mathcal{O}_C(t - b_C)) \end{aligned} \quad (\ddagger)$$

with a_C and b_C as defined in section 1.2.

3 Main Results

Given a divisor F on a blow up X of \mathbf{P}^2 at n general points, the naive conjecture that $h^0(X, F)$ always equals $\max(0, (F^2 - K_X \cdot F)/2 + 1)$ is false. One way to salvage it, is to impose a niceness requirement on F , such as to require that F be nef, or even that $F \cdot C \geq 0$ for all exceptional C , which is weaker. In fact, the SHGH Conjecture is equivalent to the conjecture that $h^0(X, F) = \max(0, (F^2 - K_X \cdot F)/2 + 1)$ whenever $F \cdot C \geq 0$ for all exceptional C .

Likewise, the naive conjecture that μ_F should always have maximal rank is also false, as we saw above. Again we can try to salvage the naive conjecture by imposing a niceness requirement on F , but the necessary requirement is more subtle. Assuming we can compute $h^0(X, F)$ for an arbitrary divisor F , Lemma 2.1.1 reduces the problem of computing the rank of μ_F in general to

the case that F is effective and fixed component free. But, as we mentioned above, even if F is effective and fixed component free, or even very ample, μ_F can fail to have maximal rank.

Instead, we will consider all $F \in L + \text{EFF}(X)$, where $\text{EFF}(X)$ is the subsemigroup of the divisor class group of X of classes of all effective divisors. By Lemma 2.1.1 we need only consider such F which are also nef. In this section, refining ideas of Fitchett, we will give an upper bound on the dimension of the cokernel of μ_F for certain $F \in L + \text{EFF}(X)$ (for all of them if the SHGH Conjecture is true). We conjecture that this upper bound is in fact an equality.

Assuming the SHGH Conjecture, Fitchett reduced the problem of handling μ_F for an arbitrary $F \in L + \text{EFF}(X)$ to the case $F = L + mE$ where E is an exceptional curve and $0 \leq m \leq L \cdot E$. (See [F1], [F2], which give explicit bounds on $\dim \text{cok } \mu_F$ for $F \in L + \text{EFF}$ in the case of $n \leq 8$ general points, using a construction originally described in Fitchett's thesis.)

We now recall Fitchett's idea, assuming that X is obtained by blowing up n general points. Note that the SHGH Conjecture would make the assumption $h^1(X, H) = 0$ automatic.

Proposition 3.1 *Let $F \in L + \text{EFF}(X)$, so we have the decomposition $F - L = H + N$ given by Lemma A1.1.1(d,e), where H is effective with $H \cdot E \geq 0$ for all exceptional E and where either $N = 0$ or $N = c_1 C_1 + \cdots + c_r C_r$ for some mutually disjoint exceptional curves C_i and integers $c_i > 0$. Assume that $h^1(X, H) = 0$. Then, $\text{cok } \mu_F \cong \bigoplus_{i=1, \dots, r} \text{cok } \mu_{L+c_i C_i}$. If moreover $F \cdot C_i \geq 0$, then $0 < c_i \leq L \cdot C_i$.*

Proof. By Lemma A1.1.1(d,e), $F = L + H + N$ where either $N = 0$ or $N = c_1 C_1 + \cdots + c_r C_r$ for some mutually disjoint exceptional curves C_i and integers $c_i > 0$, and where H is effective and orthogonal to N . Then $h^1(X, H) = 0$ implies $\text{cok } \mu_{L+H} = 0$ by Lemma 2.3.2. Taking cohomology of

$$0 \rightarrow \mathcal{O}_X(H) \rightarrow \mathcal{O}_X(L+H) \rightarrow \mathcal{O}_L(H+L) \rightarrow 0$$

and using $h^1(X, H) = 0$ implies that $h^1(X, L+H) = 0$. Similarly, we also have $h^1(X, 2L+H) = 0$. So sequence $\mathcal{S}(F-N, N)$ of Lemma 2.3.1 holds (since $F-N = L+H$) and tells us that the cokernels for μ_F and $\mu_{N;F}$ are isomorphic. But $\mathcal{O}_N(F)$ is isomorphic to $\mathcal{O}_N(L+N)$, since $H \cdot N = 0$, and the C_i are disjoint, so $\mathcal{O}_N(L+N) \cong \bigoplus \mathcal{O}_{C_i}(L+c_i C_i)$; we finally get that the cokernel of $\mu_{N;F}$ is isomorphic to the direct sum of the cokernels of $\mu_{c_i C_i; L+c_i C_i}$. Moreover, $h^1(X, L) = h^1(X, 2L) = 0$, so sequence $\mathcal{S}(L, c_i C_i)$ of Lemma 2.3.1 gives $\text{cok } \mu_{c_i C_i; L+c_i C_i} \cong \text{cok } \mu_{L+c_i C_i}$ since $\text{cok } \mu_L = 0$. Hence, as Fitchett observed, the cokernel for μ_F is isomorphic to the direct sum of the cokernels for $\mu_{L+c_i C_i}$. ■

This and Lemma 2.1.1 motivate the following problem:

Problem 3.2 *Determine the rank of μ_F for each $F = L + iE$, where E is smooth and rational with $E^2 = -1$ and $0 \leq i \leq E \cdot L$.*

Concerning this problem, we prove Theorem 3.3, which gives explicit upper bounds for the dimension of $\text{cok } \mu_{L+iE}$. Similar but less precise results were given in [F1], [F2]. Theorem 3.3(c) seems to be entirely new, however.

For the purpose of stating the result, given $0 \leq a \leq b$, let us define a sequence $\theta = (\theta_j)$, where $\theta_j = \theta_j(a, b)$, as follows. Let $\theta_j = 0$ for $j < a+3$ and for $j > b$. For values $a+3 \leq j \leq (a+b+3)/2$, let $\theta_j = j - (a+2)$ and for values $(a+b+3)/2 \leq j \leq b$, let $\theta_j = b+1-j$. Note that θ is symmetric and unimodal, and the number of nonzero values is $(b-a-2)_+$, where for any integer q , $(q)_+$ is the maximum of q and 0. If $a = 2$ and $b = 9$, then the nonzero values of θ in order would be $(1, 2, 3, 2, 1)$, while $a = 3$ and $b = 11$ gives $(1, 2, 3, 3, 2, 1)$.

Theorem 3.3 *Let X be the blow up of \mathbf{P}^2 at n distinct points P_1, \dots, P_n , and take L, E_1, \dots, E_n as usual. Let $F = L + iE$, where $d = E \cdot L$, m is the maximum of $E \cdot E_j$ over $1 \leq j \leq n$, and where E is smooth and rational with $E^2 = -1$ and $0 \leq i \leq d$. Let $\theta_j = \theta_j(a_E, b_E)$. Then we always have*

$$\binom{i - b_E}{2} + \binom{i - a_E}{2} - \sum_{0 \leq j \leq i} \theta_j \leq \dim \operatorname{cok} \mu_F \leq \binom{i - b_E}{2} + \binom{i - a_E}{2},$$

and we have the equality

$$\dim \operatorname{cok} \mu_F = \binom{i - b_E}{2} + \binom{i - a_E}{2}$$

in the following cases:

- (a) $i \leq a_E + 2$;
- (b) $b_E - a_E \leq 2$; or
- (c) $a_E = d - m$.

Proof. Let $F_j = L + jE$ for $0 \leq j < i$; it is easy to check that $H^1(F_j) = 0 = H^1(F_j + L)$, so we can consider $\mathcal{S}(F_j, E)$. Since $(F_j + E) \cdot E = d - j \geq 0$, by $(\dagger\dagger)$ and Lemma 2.1.2 we have

$$\ker \mu_{E;F_j} = H^0(E, \mathcal{O}_E(b_E - j)) \oplus H^0(E, \mathcal{O}_E(a_E - j)),$$

$\operatorname{cok} \mu_{E;F_j} = H^1(E, \mathcal{O}_E(b_E - j)) \oplus H^1(E, \mathcal{O}_E(a_E - j))$, and $\dim \ker \mu_{E;F_j} = (b_E - j + 1)_+ + (a_E - j + 1)_+$, while $\dim \operatorname{cok} \mu_{E;F_j} = (j - b_E - 1)_+ + (j - a_E - 1)_+$.

Writing a and b for a_E and b_E , we have the following exact sequences:

$$\begin{array}{llll} \operatorname{cok} \mu_{F_{i-1}} & \rightarrow & \operatorname{cok} \mu_{F_i} & \rightarrow & H^1(\mathcal{O}_E(b - i) \oplus \mathcal{O}_E(a - i)) \rightarrow 0 \\ \operatorname{cok} \mu_{F_{i-2}} & \rightarrow & \operatorname{cok} \mu_{F_{i-1}} & \rightarrow & H^1(\mathcal{O}_E(b - i + 1) \oplus \mathcal{O}_E(a - i + 1)) \rightarrow 0 \\ \dots & & & & \\ \operatorname{cok} \mu_L & \rightarrow & \operatorname{cok} \mu_{L+E} & \rightarrow & H^1(\mathcal{O}_E(b - 1) \oplus \mathcal{O}_E(a - 1)) \rightarrow 0 \end{array} \quad (\ddagger)$$

Since $\operatorname{cok} \mu_L = 0$, $\dim \operatorname{cok} \mu_{F_i}$ is at most the sum of the dimensions of the column of H^1 's, which is $\sum_{j \geq 0} ((i - b - 1 - j)_+ + (i - a - 1 - j)_+) = \binom{i-b}{2} + \binom{i-a}{2}$, and equality holds if and only if the displayed sequences are all exact on the left. Moreover, $h^1(\mathcal{O}_E(b - j) \oplus \mathcal{O}_E(a - j)) = 0$ for all $j \leq a + 1$, so $\operatorname{cok} \mu_{F_j} = 0$ for all $0 \leq j \leq a + 1$. Thus each sequence is exact for which F_j in the middle column has index $j \leq a + 2$. This implies claim (a). Moreover, any of the sequences for which $\dim \ker \mu_{E;F_j} = 0$ will also be exact on cokernels, and $\dim \ker \mu_{E;F_j} = 0$ for all $j \geq b + 1$. It follows that equality holds if $j > a + 2$ implies $j > b$; i.e., if $a + 2 \geq b$. This shows (b).

By taking account of the contributions coming from $\ker \mu_{E;F_j}$, we have that $\dim \operatorname{cok} \mu_{F_j} \geq \dim \operatorname{cok} \mu_{E;F_j} + (\dim \operatorname{cok} \mu_{F_{j-1}} - \dim \ker \mu_{E;F_j})_+$. Induction using the fact that $\dim \operatorname{cok} \mu_{F_j} = 0$ for small j now gives the lower bound in the statement of the theorem. (It is perhaps more persuasive to check an example or two rather than submit oneself to reading a detailed proof.)

Finally, consider (c); thus $b = m$. It is enough to show that the maps $\ker \mu_{F_j} \rightarrow \ker \mu_{E;F_j}$ are onto for $a + 3 \leq j \leq b$; we already observed above that exactness holds on cokernels (and hence for kernels) for other values of j . We may assume, after reindexing if need be, that $E_1 \cdot E = m$.

From the exact sequence $0 \rightarrow \mathcal{O}_E(m - d) \rightarrow \mathcal{O}_E \otimes H^0(X, L - E_1) \rightarrow \mathcal{O}_E(d - m) \rightarrow 0$ we see that $\ker \mu_{E;F_j, L - E_1} \cong H^0(E, F_j \cdot E + m - d) = H^0(E, m - j)$. The inclusion $H^0(X, L - E_1) \subset H^0(X, L)$ induces an inclusion $\ker \mu_{E;F_j, L - E_1} \rightarrow \ker \mu_{E;F_j, L} = \ker \mu_{E;F_j}$. Unwinding definitions, we see that the induced map $H^0(E, m - j) \cong \ker \mu_{E;F_j, L - E_1} \hookrightarrow \ker \mu_{E;F_j}$ sends an element $\sigma \in H^0(E, m - j)$

to $x|_E\sigma \otimes y - y|_E\sigma \otimes x \in \ker \mu_{E;F_j}$, if we choose homogeneous coordinates x, y and z on \mathbf{P}^2 such that P_1 is the point where $x = 0 = y$.

Since $a + 3 \leq j \leq b = m$, the induced inclusion $H^0(E, m - j) \hookrightarrow \ker \mu_{E;F_j}$ is an isomorphism. Moreover, taking cohomology of $0 \rightarrow \mathcal{O}_X(E_1 + (j - 1)E) \rightarrow \mathcal{O}_X(E_1 + jE) \rightarrow \mathcal{O}_E(m - j) \rightarrow 0$ gives the map $H^0(X, E_1 + jE) \rightarrow H^0(E, \mathcal{O}_E(m - j))$, which by induction on j is surjective for $0 \leq j \leq m + 1$. Composing $H^0(X, E_1 + jE) \rightarrow H^0(E, \mathcal{O}_E(m - j))$ with the induced isomorphism $H^0(E, m - j) \rightarrow \ker \mu_{E;F_j}$ gives for each $f \in H^0(X, E_1 + jE)$ the map $f \mapsto f|_E \mapsto x|_E f|_E \otimes y - y|_E f|_E \otimes x \in \ker \mu_{E;F_j}$. Thus every element of $\ker \mu_{E;F_j}$ is of the form $f|_E \mapsto x|_E f|_E \otimes y - y|_E f|_E \otimes x$ where $f \in H^0(X, E_1 + jE)$. But $f|_E \mapsto x|_E f|_E \otimes y - y|_E f|_E \otimes x$ is the image of $xf \otimes y - yf \otimes x \in \ker \mu_{F_j}$ under the map $\ker \mu_{F_j} \rightarrow \ker \mu_{E;F_j}$, so the map is surjective. \blacksquare

In fact, we do not know any times that the equality in the theorem does not hold. This suggests the following conjecture:

Conjecture 3.4 *Let $F = L + iE$ be as in Theorem 3.3 for general points P_i . Then $\dim \text{cok } \mu_F = \binom{i - b_E}{2} + \binom{i - a_E}{2}$.*

Remark 3.5 (a) Conjecture 3.4 is equivalent to the first column of maps in (\ddagger) all being injective. Hence if Conjecture 3.4 holds for some i , then it holds for all $0 \leq j \leq i$. Moreover, Conjecture 3.4 holds for $i = L \cdot E$ if and only if it holds for all $a_E + 3 \leq i \leq b_E$, since the proof of Theorem 3.3 shows that the first column of maps in (\ddagger) are injective (i.e., $\theta_i = 0$) for $i \leq a_E + 2$ and for $i > b_E$.

(b) In the notations of Proposition 3.1, assuming the SHGH Conjecture and Conjecture 3.4, and assuming we can determine splitting types, we thus can determine the dimension of the cokernel of μ_F for any F as long as $|F - L|$ is not empty. The splitting type of an exceptional curve can be computed fairly efficiently, at least provisionally (that is, by Macaulay 2 [GS], say, in positive characteristic, using randomly chosen points; a Macaulay 2 script that does this is included in Section A2.3). We discuss this, and we give additional, computational, support for Conjecture 3.4, in Section A2 of the appendix.

(c) Translating in terms of fat points, this says that we can produce conjectural dimensions for the cokernels of μ_t for $I(Z)$ in every degree t but $t = \alpha(Z)$. We can even sometimes determine the dimension for the cokernels of μ_α , for example by applying Lemma A1.2.6 (see Example A1.2.7), or if $h_Z(\alpha) = 1$, or if $F_\alpha(Z)$ decomposes as $F_\alpha(Z) = L + H + N$ where $h^1(X, H) = 0$ and $H \cdot L \geq 0$, even if H is not effective (see Example A1.2.8).

(d) Conjecture 1.2.2 is equivalent to Conjecture 3.4, assuming the SHGH Conjecture. The first sum in the bound in Theorem 1.2.1 is exactly the difference term in Lemma 2.1.1, which accounts for the contribution to the cokernel owing to loss in fixed components in going from degree t to degree $t + 1$. (This term does not occur in Conjecture 3.4 since the F there is base curve free.) The second sum in the bound in Theorem 1.2.1 sums up exactly what each of the disjoint exceptional curves in the base locus of $F_t(Z) - L$ should contribute to the cokernel, according to Proposition 3.1 and Conjecture 3.4.

Proof of Theorem 1.2.1: Let $t > \alpha(Z)$ and let $F = F_t(Z)$. Thus $F - L$ is effective. By the SHGH Conjecture (2.2.1), as in the proof of Theorem 2.2.2, we have $F - L = H + N$, where H is nef and effective with $h^1(X, H) = 0$, and where $N = c_1 C_1 + \cdots + c_r C_r$ is a sum of pairwise orthogonal exceptional curves orthogonal to H with the curves C_j being the proper transforms of those curves in the base locus of $I(Z)_{t-1}$ which are negative for the points P_i . Let N' be that part of N which remains in the base locus for $|F|$ and let N'' be what remains in the base locus of $|F + L|$; thus $N' = c'_1 C_1 + \cdots + c'_r C_r$ and $N'' = c''_1 C_1 + \cdots + c''_r C_r$. Note that $c_j - c'_j \leq \deg(C_j)$,

because $-c_j = F_{t-1}(Z) \cdot C_j$ and $-c_j \leq -c'_j = \min(F_t(Z) \cdot C_j, 0) \leq L \cdot C_j + F_{t-1}(Z) \cdot C_j = \deg(C_j) - c_j$. By Lemma 2.1.1, $\dim \text{cok } \mu_t = \dim \text{cok } \mu_F = \dim \text{cok } \mu_{H+L+N-N'} + (h^0(X, F+L) - h^0(X, H+2L+N-N'))$. By Proposition 3.1 and Theorem 3.3, $\dim \text{cok } \mu_{H+L+N-N'} = \sum_j \dim \text{cok } \mu_{L+(c_j-c'_j)N_j} \leq \sum_j \left(\binom{c_j-c'_j-a_{C_j}}{2} + \binom{c_j-c'_j-b_{C_j}}{2} \right)$. Using Riemann-Roch and Conjecture 2.2.1 gives $h^0(X, F+L) - h^0(X, H+2L+N-N') = \sum_j d_j(c'_j - c''_j) - \sum_j \binom{c'_j - c''_j}{2}$. Putting these together gives the result. ■

References

- [AC] E. Arbarello and M. Cornalba, *Footnotes to a paper of B. Segre*, Math. Ann., 256 (1981), 341–362.
- [As] M.-G. Ascenzi, *The restricted tangent bundle of a rational curve in \mathbf{P}^2* , Comm. Algebra 16 (1988), no. 11, 2193–2208.
- [BI] E. Ballico and M. Idà, *On the minimal free resolution for fat point schemes of multiplicity at most 3 in \mathbf{P}^2* , preprint, 2006.
- [BZ] A. Buckley and M. Zompatori, *Linear systems of plane curves with a composite number of base points of equal multiplicity*, Trans. Amer. Math. Soc. 355 (2003), no. 2, 539–549.
- [Cas] G. Castelnuovo, *Ricerche generali sopra i sistemi lineari di curve piane*, Mem. Accad. Sci. Torino, II 42 (1891).
- [Cat] M. V. Catalisano, *“Fat” points on a conic*, Comm. Alg. 19(8) (1991), 2153–2168.
- [CM1] C. Ciliberto and R. Miranda, *Degenerations of planar linear systems*, J. Reine Angew. Math. 501 (1998), 191–220.
- [CM2] C. Ciliberto and R. Miranda, *Linear systems of plane curves with base points of equal multiplicity*, Trans. Amer. Math. Soc. 352 (2000), no. 9, 4037–4050.
- [CM3] C. Ciliberto and R. Miranda, *Nagata’s conjecture for a square or nearly-square number of points*, Ric. Mat. 55 (2006), no. 1, 71–78.
- [CCMO] C. Ciliberto, F. Cioffi, R. Miranda and F. Orecchia, *Bivariate Hermite interpolation and linear systems of plane curves with base fat points*, in: Computer mathematics, 87–102, Lecture Notes Ser. Comput., 10, World Sci. Publishing, River Edge, NJ, 2003.
- [E1] L. Evain, *La fonction de Hilbert de la réunion de 4^h gros points génériques de \mathbf{P}^2 de même multiplicité*, J. Algebraic Geom. 8 (1999), no. 4, 787–796.
- [E2] L. Evain, *Computing limit linear series with infinitesimal methods*, preprint 2004 (arXiv:math.AG/0407143).
- [F1] S. Fitchett, *On Bounding the Number of Generators for Fat Point Ideals on the Projective Plane*, J. Algebra, 236 (2001), 502–521.
- [F2] S. Fitchett, *Corrigendum to: “On bounding the number of generators for fat point ideals on the projective plane” [J. Algebra 236 (2001), no. 2, 502–521]*, J. Algebra 276 (2004), no. 1, 417–419.

- [F3] S. Fitchett, *Maps of linear systems on blow ups of the Projective Plane*, J. Pure Appl. Algebra 156 (2001), 1-14.
- [FHH] S. Fitchett, B. Harbourne and S. Holay, *Resolutions of Fat Point Ideals Involving Eight General Points of \mathbf{P}^2* , J. Algebra 244 (2001), 684–705.
- [GGR] A. V. Geramita, D. Gregory and L. and Roberts, *Monomial ideals and points in projective space*, J. Pure Appl. Alg. 40 (1986), 33–62.
- [G] A. Gimigliano, *On linear systems of plane curves*, Thesis, Queen’s University, Kingston (1987).
- [GHI1] A. Gimigliano, B. Harbourne and M. Idà, *On the ideal resolution of fat points in the plane*, in preparation.
- [GI] A. Gimigliano and M. Idà, *The ideal resolution for generic 3-fat points in \mathbf{P}^2* , J. Pure Appl. Algebra 187 (2004), no. 1-3, 99–128.
- [GS] D. Grayson, and M. Stillman, *Macaulay 2, a software system for research in algebraic geometry*, Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [Ha1] B. Harbourne, *An Algorithm for Fat Points on \mathbf{P}^2* , Can. J. Math. 52 (2000), 123–140.
- [Ha2] B. Harbourne, *The Ideal Generation Problem for Fat Points*, J. Pure Appl. Alg. 145(2), 165–182 (2000).
- [Ha3] B. Harbourne, *Free Resolutions of Fat Point Ideals on \mathbf{P}^2* , J. Pure Appl. Alg. 125, 213–234 (1998).
- [Ha4] B. Harbourne, *The Geometry of rational surfaces and Hilbert functions of points in the plane*. Can. Math. Soc. Conf. Proc., vol. 6 (1986), 95-111.
- [Ha5] B. Harbourne, *Very ample divisors on rational surfaces*, Math. Ann. 272, 139–153 (1985).
- [HHF] B. Harbourne, S. Holay and S. Fitchett, *Resolutions of ideals of quasiuniform fat point subschemes of \mathbf{P}^2* , Trans. Amer. Math. Soc. 355 (2003), no. 2, 593-608.
- [HR] B. Harbourne and J. Roé. *Linear systems with multiple base points in \mathbf{P}^2* , Adv. Geom. 4 (2004), 41–59.
- [Hi1] A. Hirschowitz, *Une conjecture pour la cohomologie des diviseurs sur les surfaces rationnelles génériques*, Journ. Reine Angew. Math. 397 (1989), 208–213.
- [Hi2] A. Hirschowitz. *La méthode d’Horace pour l’interpolation à plusieurs variables*, Manus. Math. 50 (1985), 337–388.
- [I] M. Idà, *The minimal free resolution for the first infinitesimal neighborhoods of n general points in the plane*, J. Alg. 216 (1999), 741–753.
- [Mi] T. Mignon, *Systèmes de courbes planes à singularités imposées: le cas des multiplicités inférieures ou égales à quatre*, J. Pure Appl. Algebra 151 (2000), no. 2, 173–195.
- [Mu1] D. Mumford, *Varieties defined by quadratic equations*, in: Questions on algebraic varieties, Corso C.I.M.E. 1969 Rome: Cremonese, 1970, 30–100.

- [Mu2] D. Mumford, *Lectures on curves on an algebraic surface*, Princeton 1966.
- [N1] M. Nagata, *On the 14-th problem of Hilbert*, Amer. J. Math. 81 (1959), 766–772.
- [N2] M. Nagata, *On rational surfaces, II*, Mem. Coll. Sci. Univ. Kyoto, Ser. A Math. 33 (1960), 271–293.
- [R] J. Roé, *Limit linear systems and applications*, preprint (math/0602213), 2006.
- [S] B. Segre, *Alcune questioni su insiemi finiti di punti in Geometria Algebrica*, Atti del Convegno Internaz. di Geom. Alg., Torino (1961).
- [Y] S. Yang, *Linear series in \mathbf{P}^2 with base points of bounded multiplicity*, preprint 2004 (arXiv:math.AG/0406591).

Additional Citations for the Appendix:

- [A.Ar] M. Artin, *Some numerical criteria for contractability of curves on algebraic surfaces*, Amer. J. Math. 84 (1962), 485–497.
- [A.H1] B. Harbourne, *Blowings-up of \mathbf{P}^2 and their blowings-down*, Duke Math. J. 52, 129–148 (1985).
- [A.H2] B. Harbourne, *Complete linear systems on rational surfaces*, Trans. Amer. Math. Soc. 289, 213–226 (1985).

Appendix

A1 Making the SHGH Conjecture Explicit

In this appendix we give another version of the SHGH Conjecture and show how to derive explicit predictions for values of Hilbert functions using it.

A1.1 The Weyl Group

We now recall the Weyl group $W = W_n$, which acts on $\text{Cl}(X)$ but which depends only on the number n of points P_i blown up. If $0 \leq n \leq 1$, then $W = \{id\}$ is trivial. If $n = 2$, then $W = \{id, s_1\}$, where for any divisor class F , $s_1(F) = F + (r_1 \cdot F)r_1$, where $r_1 = E_1 - E_2$. For $n > 2$, let $r_0 = L - E_1 - E_2 - E_3$ and for $1 \leq i < n$, let $r_i = E_i - E_{i+1}$. Then W is generated by the operators s_i , $0 \leq i < n$, where $s_i(F) = F + (r_i \cdot F)r_i$. It is now easy to check that W preserves the intersection form (i.e., $wF \cdot wG = F \cdot G$ for all $w \in W$ and all $F, G \in \text{Cl}(X)$), and that $wK_X = K_X$ for all $w \in W$. The subgroup generated by s_1, \dots, s_{n-1} is just the permutation group on E_1, \dots, E_n . The action of the element s_0 corresponds to that of the quadratic Cremona transformation centered at P_1, P_2, P_3 .

If $n = 0$, $X = \mathbf{P}^2$ has no exceptional curves. If $n = 1$, then E_1 is the only exceptional curve, and if $n = 2$, then E_1, E_2 and $L - E_1 - E_2$ are the only exceptional curves. For $n \geq 3$, Nagata [N2] has shown that if a class E is the class of an exceptional curve, then $E \in WE_n$; i.e., the classes of exceptional curves lie in a single W -orbit. Moreover, if $E \in WE_n$ and the points P_i are general, Nagata showed E is the class of an exceptional curve. (When $E \in WE_n$ but the points are not general, then although E is effective, it can fail to be reduced and irreducible, and thus need not be the class of an exceptional curve. For example, $2L - E_1 - \dots - E_5 \in WE_5$, but if P_1, P_2, P_3 are collinear, then the proper transform of the line through P_1, P_2, P_3 is a fixed component of $|2L - E_1 - \dots - E_5|$.)

If $n = 0$, let \mathcal{E}_n be the submonoid of $\text{Cl}(X)$ generated by L . If $n = 1$, let \mathcal{E}_n be the submonoid generated by $L - E_1$ and E_1 . If $n = 2$, let \mathcal{E}_n be generated by $L - E_1 - E_2, E_1$ and E_2 , while if $n \geq 3$, let \mathcal{E}_n be generated by $L - E_1 - E_2$ and E_1, \dots, E_n (or equivalently by the orbit WE_n of E_n under W). Thus if $n \geq 3$ every element $D \in \mathcal{E}_n$ is of the form $D = \sum c_i C_i$, where c_i is a nonnegative integer and C_i is an exceptional curve. Define \mathcal{E}_n^* to be the dual cone; thus $F \in \mathcal{E}_n^*$ means that $F \cdot D \geq 0$ for every $D \in \mathcal{E}_n$ (and thus that $F \cdot E \geq 0$ for every exceptional curve E).

Let $\text{EFF} = \text{EFF}(X) \subset \text{Cl}(X)$ denote the submonoid of classes of effective divisors, let $\text{NEF} = \text{NEF}(X) \subset \text{Cl}(X)$ denote the submonoid (indeed the cone, since a class is nef if a positive multiple is) of classes of nef divisors, let Ψ_n be the submonoid generated by the union of \mathcal{E}_n and the element $-K_X$ and let Δ_n be the submonoid of $\text{Cl}(X)$ generated by $H_0 = L, H_1 = L - E_1, H_2 = 2L - E_1 - E_2$, and $H_i = -K_X + E_{i+1} + \dots + E_n = 3L - E_1 - \dots - E_i$, for $3 \leq i \leq n$. Notice that $F = tL - m_1 E_1 - \dots - m_n E_n \in \Delta_n$ if and only if $t \geq m_1 + m_2 + m_3$ and $m_1 \geq m_2 \geq \dots \geq m_n \geq 0$, and that H_n is $-K_X$ and $H_i \cdot H_j \geq 0$ unless $i, j \geq 10$.

Since the next result is a statement for all classes on X , we need to state it in terms of a blowing up of generic points. However, when we are interested in a specific class F , it is enough to consider a blow up of general points (but the conditions of generality will depend on F). The following result is known but hard to cite.

Lemma A1.1.1 *Let X be the blow up of \mathbf{P}^2 at n generic points P_i .*

(a) *If $A \in \Delta_n$ and $w \in W_n(X)$, then $wA = A + a_0 r_0 + \dots + a_{n-1} r_{n-1}$ for some nonnegative integers a_i .*

- (b) $\text{NEF}(X) \subset \mathcal{E}_n^* = W_n \Delta_n \subset \Psi_n$
- (c) $h^j(X, F) = h^j(X, wF)$ for all j , all $w \in W_n$ and all $F \in \text{Cl}(X)$
- (d) $\text{EFF}(X) \subset \Psi_n$
- (e) If $F \in \Psi_n$, then there is a unique decomposition $F = H + N$ where $H \in \mathcal{E}_n^*$, $N \in \mathcal{E}_n$, $H \cdot N = 0$ and, if $N \neq 0$, then $N = c_1 C_1 + \cdots + c_r C_r$ where for each i , c_i is a positive integer and C_i is the class of an exceptional curve with $C_i \cdot C_j = 0$ for all $i \neq j$. Moreover, H is effective if F is.

Proof. (a) See, for example, Lemma 1.2 (1) of [Ha5]. (b) First we verify that $\mathcal{E}_n^* = W_n \Delta_n$. We leave the cases $0 \leq n < 3$ to the reader; assume $n \geq 3$. It is known that EFF and NEF are W_n invariant, and that the set of exceptional divisors is just the orbit $W_n E_n$; cf. [N2]. Since W_n preserves the set of exceptional curves, if we show $\Delta_n \subset \mathcal{E}_n^*$, then $W_n \Delta_n \subset \mathcal{E}_n^*$. But H_i is nef (hence in Δ_n) for $i \leq 9$. For $i > 9$, $H_i = -K_X + E_{i+1} + \cdots + E_n$, hence for any exceptional curve E we have $H_i \cdot E \geq 0$, since E meets $-K_X$ once and $E \cdot E_j \geq -1$ with equality if and only if $E = E_j$. Thus $\Delta_n \subset \mathcal{E}_n^*$.

Conversely, say $F \in \mathcal{E}_n^*$. Note that every element $D \in \mathcal{E}_n^*$ satisfies $D \cdot L \geq 0$, since $L = (L - E_1 - E_2) + E_1 + E_2$. Since W_n preserves \mathcal{E}_n^* , there must be some $w \in W_n$ such that $L \cdot wF$ is as small as possible. Thus, $wF \cdot r_0 \geq 0$, otherwise we would have $s_0 wF \cdot L < wF \cdot L$. We can also assume $m_1 \geq m_2 \geq \cdots \geq m_n$, where $m_i = wF \cdot E_i$, since each operator s_i , $i > 0$, merely transposes E_i and E_{i+1} , so we can in W_n permute the E_i without affecting $L \cdot wF$. Thus $wF \cdot r_i \geq 0$ for all $i \geq 1$. Finally, $wF \cdot (L - E_1 - E_2) \geq 0$ since $wF \in \mathcal{E}_n^*$, and $wF \cdot (L - E_1) \geq 0$ since $L - E_1 = (L - E_1 - E_2) + E_2$ is a sum of exceptional curves. By Lemma 1.4 of [A.H2], we thus have $wF \in \Delta_n$.

Finally, $\text{NEF} \subset \mathcal{E}_n^*$ since $\mathcal{E}_n \subset \text{EFF}$, and clearly $\Delta_n \subset \Psi_n$ but Ψ_n is W_n -invariant, so $W_n \Delta_n \subset \Psi_n$.

(c) This is, in somewhat different language, due to Nagata [N2]. The basic idea is this. Say $F = tL - m_1 E_1 - \cdots - m_n E_n$. Then $wF = t wL - m_1 wE_1 - \cdots - m_n wE_n$, where $L' = wL$, $E'_1 = wE_1, \dots, E'_n = wE_n$ is a basis of $\text{Cl}(X)$. This basis is, however, an *exceptional configuration*; i.e., there is a birational morphism $p' : X \rightarrow \mathbf{P}^2$ such that $E'_i = p'^{-1}(P'_i)$ for some points $P'_i \in \mathbf{P}^2$ and such that L' is the pullback via p' of the class of a line in \mathbf{P}^2 (see Theorem 0.1 of [A.H1]), but the points P'_i are themselves generic. Since P_i and P'_i both give generic sets of points, all that matters are the coefficients t and m_i , so we have $h^j(X, tL - m_1 E_1 - \cdots - m_n E_n) = h^j(X, tL' - m_1 E'_1 - \cdots - m_n E'_n)$; i.e., $h^j(X, F) = h^j(X, wF)$.

(d) Let $F \in \text{EFF}(X)$. Then there are at most finitely many exceptional curves E such that $F \cdot E < 0$. Let this finite set of distinct exceptional curves be C_1, \dots, C_r , let $c_i = -F \cdot C_i$, let $N = c_1 C_1 + \cdots + c_r C_r$ and let $H = F - N$. Note that $C_i \cdot C_j = 0$ for all $i \neq j$ (since $(C_i + C_j) \cdot F < 0$, but $C_i \cdot C_j > 0$ implies $(C_i + C_j)$ meets both C_i and C_j nonnegatively and hence is nef) and that $H \cdot C_i = 0$. Since F is effective, N is contained in the scheme theoretic base locus of $|F|$, hence H is effective. But if $H \cdot E < 0$, then E is not C_i for any i , hence $E \cdot F \geq 0$ so we get $E \cdot C_i > 0$ (implying $E + C_i$ is nef) for some i even though $H \cdot (E + C_i) < 0$. It follows that $H \in \mathcal{E}_n^*$. The result follows since $N \in \mathcal{E}_n \subset \Psi_n$ and $H \in \mathcal{E}_n^* \subset \Psi_n$.

(e) We leave the cases $0 \leq n < 3$ to the reader, so assume $n \geq 3$. If $F \in \Psi_n$, then $wF \cdot L \geq 0$ and $wF \cdot (L - E_1) \geq 0$ for all $w \in W_n$, since $wF \in \Psi_n$ but L and $L - E_1$ are nef and meet $-K_X$ nonnegatively. Choose w such that $wF \cdot L$ is as small as possible, and write $wF = tL - m_1 E_1 - \cdots - m_n E_n$. Since W_n includes the group of permutations of E_1, \dots, E_n , we may assume that $m_1 \geq \cdots \geq m_n$. Since t is as small as possible, we know that $t \geq m_1 + m_2 + m_3$ (otherwise $s_0 wF \cdot L < wF \cdot L$). If $m_3 \geq 0$ or $m_2 \leq 0$, let $H' = tL - \sum_{m_i > 0} m_i E_i$ and $N' = -\sum_{m_i < 0} m_i E_i$.

Using the definition of Δ and the facts that $t \geq m_1 + m_2 + m_3$ and $t \geq m_1$, it is now not hard to check that $H' \in \Delta$, and clearly $N' \in \mathcal{E}_n$. If $m_3 < 0$ and $m_2 > 0$, there are two cases. If $c = (tL - m_1E_1 - m_2E_2) \cdot (L - E_1 - E_2) < 0$, then let $H = (t + c)L - (m_1 + c)E_1 - (m_2 + c)E_2$, and let $N = (-c)(L - E_1 - E_2) - m_3E_3 - \cdots - m_nE_n$. If $(tL - m_1E_1 - m_2E_2) \cdot (L - E_1 - E_2) \geq 0$, let $H = tL - m_1E_1 - m_2E_2$, and let $N = -m_3E_3 - \cdots - m_nE_n$. Either way $H' \in \Delta_n$ and $N' \in \mathcal{E}_n$. In all cases we also have $H' \cdot N' = 0$ and that the components of N' are disjoint and orthogonal to H' . We now take $H = w^{-1}H'$ and $N = w^{-1}N'$, where the classes C_i are the w^{-1} translates of the components of N' . Uniqueness follows from the fact that $N = \sum_E \text{exceptional} -(F \cdot E)E$. Note that in case F is effective we found a decomposition $F = H + N$ in (d), with H effective. Uniqueness now shows that H is necessarily effective. \blacksquare

Remark A1.1.2 As an application of Lemma A1.1.1, *and assuming the SHGH Conjecture*, we will classify all smooth rational curves C on a blow up $p : X \rightarrow \mathbf{P}^2$ of general points. According to Conjecture 2.2.1, we must have $C^2 \geq -1$. If $C^2 = -1$, then C is an exceptional curve. If $C^2 > -1$, then C is nef, hence $wC \in \Delta$ for some $w \in W$ by Lemma A1.1.1(b), so it suffices if we find all smooth rational $C \in \Delta$. Since $C \in \Delta$, we have $C = \sum_i a_i H_i$ for some nonnegative integers a_i . Note that $C \cdot H_j \geq 0$ for $0 \leq j \leq 2$. By adjunction and $C^2 > -1$ we have $-C \cdot K_X = C^2 + 2 \geq 2$. Since C is nef, we have $C \cdot (E_{j+1} + \cdots + E_n) \geq 0$. Thus $C \cdot H_j = C \cdot (-K_X + E_{j+1} + \cdots + E_n) \geq 2$ for all $j > 2$. If $a_j > 0$ for some $j > 2$, let $C' = C - H_j$. Since C' is still a nonnegative integer combination of the H_i , we have $C \cdot C' \geq 0$. Now $C^2 = C \cdot C' + C \cdot H_j \geq C \cdot H_j \geq C \cdot H_n = -C \cdot K_X = C^2 + 2$. I.e., we must have $a_j = 0$ for all $j > 2$.

Thus $C = aH_0 + bH_1 + cH_2$. It is now an easy exercise using adjunction to show that the only solutions to $C^2 + C \cdot K_X = -2$ are $H_0, 2H_0, H_1, H_2, H_0 + bH_1$ and $H_2 + bH_1$. Thus W orbits of these and E_1 are the only possible smooth rational curves in X , assuming the SHGH Conjecture.

Each such C can be turned into an exceptional curve E by subtracting off additional E_i ; for example, if $C = H_2 + 2H_1 = 4L - 3E_1 - E_2$, then $E = 4L - 3E_1 - E_2 - \cdots - E_9$ is an exceptional curve. Moreover, $p^*\Omega(1)|_C$ has the same splitting as does $p^*\Omega(1)|_E$, since C and E both have the same image $p(C) = p(E)$ in \mathbf{P}^2 .

Thus an algorithm for computing the splitting for exceptional curves handles all known smooth rational C .

A1.2 The SHGH Conjecture

Given any class $F \in \text{Cl}(X)$, there is a geometrically defined quantity $e(h^0, F)$ such that $h^0(X, F) \geq e(h^0, F)$ holds for general points P_i . We now define this lower bound.

If $F \notin \Psi_n$, then $h^0(X, F) = 0$ by Lemma A1.1.1, and we set $e(h^0, F) = 0$. If $F \in \Psi_n$, then we have the decomposition $F = H + N$ given by Lemma A1.1.1(e), and we have $h^0(X, F) = h^0(X, H)$. Since $H \cdot L \geq 0$, we have $h^0(X, H) \geq \max(0, (H^2 - K_X \cdot H)/2 + 1)$ and we set $e(h^0, F) = \max(0, (H^2 - K_X \cdot H)/2 + 1)$. Clearly, $h^0(X, F) \geq e(h^0, F)$ holds. We can now state the SHGH Conjecture, which says that equality in fact holds:

Conjecture A1.2.1 *We have $h^0(X, F) = e(h^0, F)$, where $F \in \text{Cl}(X)$ and X is the blow up of \mathbf{P}^2 at general points P_i .*

A version of the SHGH Conjecture for fat points follows from this. Given a fat point subscheme $Z = m_1P_1 + \cdots + m_nP_n$ supported at general points P_i , we define $e(h_Z, t)$ to be $e(h^0, F_t(Z))$, where $F_t(Z) = tL - m_1E_1 - \cdots - m_nE_n$.

Conjecture A1.2.2 We have $h_Z(t) = e(h_Z, t)$, where $Z = m_1P_1 + \cdots + m_nP_n$ is a fat point scheme supported at general points P_i of \mathbf{P}^2 .

To apply these conjectures, one must be able to compute $e(h^0, F)$. We give two examples showing how to do so.

Example A1.2.3 Suppose $F = tL - (77(E_1 + \cdots + E_7) + 44E_8 + 11E_9 + 11E_{10} + 11E_{11})$. To compute $e(h^0, F)$ for any given t , just mimic the proof of Lemma A1.1.1(e). The idea is to find an element $w \in W$ such that either $wF \cdot L$ is as small as possible, or $wF \cdot L < 0$ or $wF \cdot (L - E_1) < 0$. For example, say $t = 208$. Apply s_0 to F to get $s_0F = 185L - 54E_1 - 54E_2 - 54E_3 - 77E_4 - 77E_5 - 77E_6 - 77E_7 - 44E_8 - 11E_9 - 11E_{10} - 11E_{11}$. Permute the E_i so that the coefficients are nondecreasing, which gives $F' = 185L - 77E_1 - 77E_2 - 77E_3 - 77E_4 - 54E_5 - 54E_6 - 54E_7 - 44E_8 - 11E_9 - 11E_{10} - 11E_{11}$. This operation, taking F to F' , is now repeated until we obtain a class F'' such that either $F'' \cdot L < 0$, or $F'' \cdot (L - E_1) < 0$, or until $F'' \cdot L \geq 0$, $F'' \cdot (L - E_1) \geq 0$ and $F'' \cdot r_0 \geq 0$. In this case the class F'' we eventually end up with is $-23L - 8E_1 + E_2 + 5E_3 + 5E_4 + 5E_5 + 5E_6 + 8E_7 + 8E_8 + 14E_9 + 17E_{10} + 17E_{11}$, hence $e(h^0, F) = 0$, since $F'' \cdot L < 0$ (so $F'' \notin \Psi_{11}$). If, for example, $t = 209$, then the class we end up with is $F'' = 11E_{11}$, so the decomposition of Lemma A1.1.1(d) is $H = 0$ and $N = F$, so $e(h^0, F) = 1$. And if $t = 210$, then we end up with $F'' = 27L - 8(E_1 + \cdots + E_4) - 5(E_5 + \cdots + E_{11})$, which is in Δ , so $F = H$, $N = 0$, and $e(h^0, F) = (H^2 - K_X \cdot H)/2 + 1$. In this example, the SHGH Conjecture, that $h^0(X, F) = e(h^0, F)$, in fact holds for all t . It holds for $t < 209$ since $F \notin \Psi_n$ for those t . It holds for $t = 209$, since $wF = 11E_{11}$ for some w , so $h^0(X, F) = h^0(X, 11E_{11}) = 1$. And it holds for $t > 209$ since for these cases $F \in \mathcal{E}_{11}^* = W_{11}\Delta_{11}$, so $F = H$, and $-K_X \cdot F \geq 0$, so $h^0(X, F) = (F^2 - K_X \cdot F)/2 + 1$ by Theorem 1.1 of [A.H2] and semicontinuity of h^0 . (Macaulay 2 scripts for carrying out both the Lemma A1.1.1(d) decomposition and the Weyl group calculations, and a sample Macaulay 2 session demonstrating their use, are included in Section A2.3.)

Example A1.2.4 Now consider $F = tL - 50E_1 - 50E_2 - 38E_3 - 38E_4 - 26E_5 - 26E_6 - 22E_7 - 18E_8 - 14E_9 - 14E_{10}$. As in Example A1.2.3, we have $e(h^0, F) = 0$ for $t < 102$, since $F \notin \Psi_{10}$. For $t = 102$, we find a w such that $wF = 6L - 2(E_2 + \cdots + E_8) + 2E_9 + 6E_{10}$. Thus the decomposition $F = H + N$ has $H = w^{-1}(6L - 2(E_2 + \cdots + E_8))$ and $N = w^{-1}(2E_9 + 6E_{10})$, where w^{-1} can be performed by simply reversing the operations which gave w . What we find is $H = 38L - 18E_1 - 18E_2 - 14E_3 - 14E_4 - 10E_5 - 10E_6 - 8E_7 - 8E_8 - 6E_9 - 6E_{10}$, and $N = 2C_1 + 6C_2$, where $C_1 = 8L - 4E_1 - 4E_2 - 3E_3 - 3E_4 - 2E_5 - 2E_6 - E_7 - 2E_8 - E_9 - E_{10}$ and $C_2 = 8L - 4E_1 - 4E_2 - 3E_3 - 3E_4 - 2E_5 - 2E_6 - 2E_7 - E_8 - E_9 - E_{10}$. Thus $h^0(X, F) = h^0(X, H)$, and it is known that $h^0(X, H) = 4$. For $t \geq 103$, we have $F = H$ and $N = 0$. In fact, for $t = 103$ we have $h^0(X, F) = 92$, and for $t = 104$ we have $h^0(X, F) = 197$. (For the same reasons as in Example A1.2.3, the SHGH Conjecture holds for F for all t .)

Given a fat point subscheme $Z \subset \mathbf{P}^2$ with general support, we can now define the expected value $e(g_\bullet(Z), i)$ of the Betti number $g_i(Z)$ for $i > \alpha + 1$:

Definition A1.2.5 Let $F = F_{i-2}(Z)$; note that F is effective, since $i > \alpha + 1$. Thus we have a decomposition $F = H + N$, with $N = c_1C_1 + \cdots + c_rC_r$, as in Lemma A1.1.1(e). Let $m_i = \min(c_i, L \cdot C_i)$. Let $M = m_1C_1 + \cdots + m_rC_r$, so $N - M$ is effective and $F = H + M + (N - M)$. Then $\dim \text{cok } \mu_{L+H+M} = \dim \text{cok } \mu_{L+M}$ by Lemma 2.3.2 assuming the SHGH Conjecture, so $g_i(Z) = \dim \text{cok } \mu_{L+F} = \dim \text{cok } \mu_{L+M} + (h^0(X, 2L + F) - h^0(X, 2L + H + M))$ by Lemma 2.1.1, and $\dim \text{cok } \mu_{L+M} \leq \sum_i \binom{m_i - bc_i}{2} + \binom{m_i - ac_i}{2}$ by Theorem 3.3 with equality assuming Conjecture 3.4. Thus we take $e(g_\bullet(Z), i)$ to be $(h^0(X, 2L + F) - h^0(X, 2L + H + M)) + \sum_i \binom{m_i - bc_i}{2} + \binom{m_i - ac_i}{2}$. (In the notation of the proof of Theorem 1.2.1, M is $N - N'$ and $m_i = c_i - c'_i$, so the upper bound in Theorem 1.2.1 is by the proof of Theorem 1.2.1 exactly $e(g_\bullet(Z), i)$.)

The following result will be useful in our examples. Versions of this result were proved in [Ha2] and [FHH] and were the basis for the results in [Ha1] and [FHH].

Lemma A1.2.6 *Let $Z = m_1P_1 + \dots + m_nP_n$ be a fat point subscheme of \mathbf{P}^2 . Assume $F = F_k(Z)$ is the class of an effective divisor on X , and define $h(F) = h^0(X, F)$, $l(F) = h^0(X, F - (L - E_1))$, $l^*(F) = h^1(X, F - (L - E_1))$, $q(F) = h^0(X, F - E_1)$, and $q^*(F) = h^1(X, F - E_1)$. Then*

$$l(F) \leq \dim \ker \mu_F \leq l(F) + q(F).$$

If moreover $h^1(X, F) = 0$, then

$$k + 2 - 2h(F) + l(F) \leq \dim \operatorname{cok} \mu_F \leq q^*(F) + l^*(F).$$

Here are two examples showing explicitly how to compute expected values of the graded Betti numbers.

Example A1.2.7 Suppose we want to determine the graded Betti numbers of the fat points subscheme $Z = m_1P_1 + \dots + m_nP_n \in \mathbf{P}^2$ where $n = 10$ here, the points P_i are general and the sequence of multiplicities m_i is $(50, 50, 38, 38, 26, 26, 22, 18, 14, 14)$. We found the Hilbert function in Example A1.2.4, from which it follows that $g_i = 0$ except for $i = 102$ (where we have $g_{102} = h_Z(102) = 4$) and possibly for $i = 103$ and $i = 104$. We do not have an expected value for g_{103} , since $103 = \alpha(Z) + 1$, but an ad hoc use of Lemma A1.2.6 gives $\dim \ker \mu_{F_{102}(Z)} \leq l(F_{102}(Z)) + q(F_{102}(Z))$, where $l(F_{102}(Z)) = h^0(X, F_{102}(Z) - (L - E_1))$ and $q(F_{102}(Z)) = h^0(X, F_{102}(Z) - E_1)$. Neither $F_{102}(Z) - (L - E_1)$ nor $F_{102}(Z) - E_1$ is in Ψ , so $l = q = 0$, so $\mu_{F_{102}(Z)}$ is injective, hence $g_{103} = \dim \operatorname{cok} \mu_{F_{102}(Z)} = h^0(X, F_{103}(Z)) - 3h^0(X, F_{102}(Z)) = 92 - 3(4) = 80$. To compute g_{104} , recall in Example A1.2.4 we found $F_{103}(Z) = H + 2C_1 + 6C_2$. From Lemma 2.1.2 we have $a_{C_i} = b_{C_i} = 4$ for $i = 1, 2$. Since $h^1(X, H) = 0$ holds here, we have $g_{104} = \dim \operatorname{cok} \mu_{F_{103}(Z)} = \dim \operatorname{cok} \mu_{L+2C_1} + \dim \operatorname{cok} \mu_{L+6C_2}$. By Theorem 3.3 we have $\dim \operatorname{cok} \mu_{L+2C_1} = 0$ and $\dim \operatorname{cok} \mu_{L+6C_2} = 2$, so $g_{104} = 2$. For $i \geq 104$, we have $h^1(X, F_i(Z)) = 0$, hence $g_{i+1} = 0$ by Lemma 2.3.2.

We can now write down a minimal free graded resolution for $I(Z)$. It is $0 \rightarrow M_1 \rightarrow M_0 \rightarrow I(Z) \rightarrow 0$, where $M_0 = R^2[-104] \oplus R^{80}[-103] \oplus R^4[-102]$, and from the Hilbert functions of $I(Z)$ and M_0 we now find $M_1 = R^{16}[-105] \oplus R^{69}[-104]$.

Example A1.2.8 Consider $Z = 48P_1 + 33P_2 + 33P_3 + 33P_4 + 32P_5 + 32P_6 + 32P_7 + 24P_8 + 16P_9$, where the points P_i are general. Then $h_Z(t) = 0$ for $t < 98$, since $F_t(Z) \notin \Psi$, and $h_Z(t) = \binom{t+2}{2} - 4879$ for $t \geq 98$, since then $h_Z(t) > 0$, $F_t(Z) \in \mathcal{E}^*$ and the SHGH Conjecture is known to hold for $n = 9$ general points (see [N2], or use the fact that any nef divisor F on a blow up of \mathbf{P}^2 at 9 general points has $-K_X \cdot F \geq 0$ and apply the results of [A.H2]). We also know that $g_t = 0$ for $t < 98$, and $g_{98} = h_Z(98) = 71$. By Lemma 2.3.2, we have $g_t = 0$ for $t > 99$. In this case $\alpha = 98$, and we do not in general have a conjectural value for $g_{\alpha+1}$, but in this case there is an element $w \in W$ such that $F_{97}(Z) = L + H + N$ where $H = L - E_1 - E_2 - E_3$, $N = 8E$, where E is the exceptional curve $96L - 48E_1 - 32E_2 - \dots - 32E_7 - 24E_8 - 16E_9$. Since $h^1(X, H) = 0$, reasoning as in the proof of Proposition 3.1, we see that μ_{L+H+8E} and μ_{L+8E} have isomorphic cokernels. By Lemma 2.1.2 we have $a_e = 6 = b_E$, so by Theorem 3.3 we have $g_{99} = \dim \operatorname{cok} \mu_{L+8E} = 2$. The minimal free graded resolution for $I(Z)$ is thus $0 \rightarrow R^{28}[-100] \oplus R^{44}[-99] \rightarrow R^2[-99] \oplus R^{71}[-98] \rightarrow I(Z) \rightarrow 0$.

A2 Computational Aspects

In this section we discuss various computational aspects of the problem of computing graded Betti numbers, partly to explain how our geometric approach can be used to make computer calculations more efficient, and partly to give additional evidence in support of Conjecture 3.4.

A2.1 Splitting Types

The first issue is the need to determine the splitting of the restriction of $p^*\Omega_{\mathbf{P}^2}(1)$ to a smooth rational curve $C \subset X$, where $p : X \rightarrow \mathbf{P}^2$ is a blow up of general points. We are mainly interested in doing this for exceptional curves, but it is of interest also to consider any smooth rational curve C . However, doing so for exceptional curves suffices to do it for all other smooth rational C known (see Remark A1.1.2).

So suppose E is an exceptional curve on X . The simplest approach conceptually is to find a Cremona transformation w of the plane that transforms $p(E)$ to a line A . Pick a general basis f_1, f_2, f_3 of the linear forms on \mathbf{P}^2 and find their images $g_i = f_i \circ w^{-1}$ under w . Given the equation of A , which is easy to get since $\deg(A) = 1$, we can find the ideal J' generated by the restrictions of the g_i to A . This ideal typically has base points; the ideal J residual to the base points is just the ideal generated by the restriction of the f_i to E , but regarding $A = E$ as \mathbf{P}^1 , we can find a minimal free resolution of the ideal over $K[\mathbf{P}^1]$. The degree of the syzygy of least degree is a_E ; then $b_E = d - a_E$ where d is the degree of $p(E)$.

It is not hard to convert this conceptual algorithm into code. For speed, all of the actual symbolic operations should be done in $K[\mathbf{P}^1]$ rather than in $K[\mathbf{P}^2]$. That is, one does not actually want to find the g_i first and then restrict to the line.

We now discuss how to push all of the computation down to $K[\mathbf{P}^1]$. An explicit script that implements the computation and which is very fast is included in Section A2.3. Here is how it works. Suppose we are given d and m_i such that $E = dL - m_1E_1 - \cdots - m_nE_n$ is the exceptional curve. Choose n generic points; in practice we choose n points of \mathbf{P}^2 at random, working over a field of finite but large characteristic. And instead of picking a general basis f_i , we pick three random linear forms. Instead of finding w , we employ a series of quadratic transformations, and find the result of each one as we go. (Of course, picking randomly means that sometimes the choices are bad, and the script gives a wrong result, such as would occur if we attempt to do a quadratic transformation, but the three points at which it is centered turn out to be on a line.)

Let $K[\mathbf{P}^2] = K[x, y, z]$. Now, for example, say $E = 3L - 2E_1 - E_2 - \cdots - E_7$ and assume that we have randomly chosen 7 points $P_i \in \mathbf{P}^2$ and three random linear forms f_i . Use an element of the projective general linear group to move P_1, P_2, P_3 to $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$. Replace the original P_i by their images under this change of coordinates, and replace the f_i by what they become in the new coordinate system. Now perform a quadratic transformation centered at P_1, P_2, P_3 (which are now coordinate vertices). I.e., replace the current points P_i by what they become under the quadratic transformation (i.e., the coordinate vertices are still P_1, P_2, P_3 , and for $i > 3$, $P_i = (a, b, c)$ becomes $P_i = (bc, ac, ab)$), and replace the linear forms f_i by what they become under the quadratic transformation (so $f_i(x, y, z)$ becomes $f_i(yz, xz, xy)$). Note that E also transforms, to $2L - E_1 - E_4 - \cdots - E_7$; i.e., its image in \mathbf{P}^2 is a conic through what are now the points P_1, P_4, \dots, P_7 . Now choose another three points at which to perform another quadratic transformation that will reduce $2L - E_1 - E_4 - \cdots - E_7$ further. For example, a quadratic transformation centered at P_4, P_5, P_6 transforms $2L - E_1 - E_4 - \cdots - E_7$ to $L - E_1 - E_7$. Repeating what we did before, we now have three forms f_i (of degree 4, in this example, since the degree doubles after each application of a quadratic transformation). By performing one last change of coordinates, we can assume that the image of $L - E_1 - E_7$ under p is the line $x = 0$. Thus the restrictions f'_i of the f_i to this line are obtained by removing all terms in which x appears.

There are a number of problems with the approach we just described. First, the degree of the f_i doubles at each step. If more than a few steps are needed, the degree gets unreasonably large. What turns out to be worse, is that the projective changes of coordinates, although they are just linear substitutions, seem to run very slowly: substituting into the f_i to find what they

transform to under the projective linear (and also under the quadratic transformations) is quite slow. Altogether, examples seem to show that computing splittings this way seems no faster than computing resolutions of ideals of fat points directly, in which case giving solutions to fat point problems in terms of splittings is not much help. But both problems can be overcome.

We can keep the degree of the f_i from growing so fast by removing extraneous factors as we go. For example, performing a quadratic transformation centered at coordinate vertices takes the line $y+z$ to the conic $xz+xy$. The image of the line $y+z=0$ is of course itself, which we see by removing the extraneous factor of x . If, at any step, performing a quadratic transformation introduces an extraneous factor, we should remove it. (This is in any case obligatory before restricting to the ultimate line $x=0$, since x itself may be an extraneous factor.)

The other issue, that this approach is slow, seems mostly due to working in $K[\mathbf{P}^2]$ until the end, where we restrict to $K[\mathbf{P}^1]$. This can be fixed essentially by performing the procedure described before in reverse order, thereby by working in $K[\mathbf{P}^1]$ throughout. In particular, first find a sequence χ_t of quadratic transformations that take $E = dL - m_1E_1 - \cdots - m_nE_n$ to some E' with $E' \cdot L = 1$. (All that you need to know is which three points at each stage the desired quadratic transformation will be centered at. So for $E = 3L - 2E_1 - E_2 - \cdots - E_7$, we need two transformations, the first can be taken to be centered at P_1, P_2, P_3 , the second at P_4, P_5, P_6 .) Choose n random points P_i ; since $E' = L - E_{i_1} - E_{i_2}$, E' corresponds to the line through the points P_{i_1} and P_{i_2} . Choose coordinates on E' ; i.e., variables u and v such that $K[E'] = K[u, v]$. The inclusion $p(E') \subset \mathbf{P}^2$ induces a linear homomorphism $K[\mathbf{P}^2] \rightarrow K[E'] = K[u, v]$. Let g_1, g_2, g_3 be the images of the variables x, y, z on \mathbf{P}^2 . The last quadratic transformation in our sequence induces a homomorphism $K[\mathbf{P}^2] \rightarrow K[\mathbf{P}^2]$ which composes to give $K[\mathbf{P}^2] \rightarrow K[\mathbf{P}^2] \rightarrow K[E']$. In terms of the g_i we want to find the images of x, y, z under this composition. These images, which we can express in terms of products of pairs of linear combinations of the g_i , replace the previous g_i . Eventually, after applying each quadratic transformation (in reverse order) in our sequence, we end up with the images $g_i \in K[u, v]$ of x, y and z under restriction to E . These generate an ideal in $K[u, v]$, from whose resolution over $K[u, v]$, as before, we obtain a_E and b_E .

It would be nice to be able to predict what a_E and b_E should be, based only on knowing t and the m_i , given a class $[E] = tL - m_1E_1 - \cdots - m_nE_n$. This seems to be a difficult problem, with Lemma 2.1.2 being the main result. However, computational data suggests a possible new constraint on the splitting type for a smooth rational curve C with $C^2 = 1$. Using sequences analogous to (‡) we get homomorphisms

$$0 \rightarrow \text{cok } \mu_{1C} \rightarrow \text{cok } \mu_{2C} \rightarrow \text{cok } \mu_{3C} \rightarrow \cdots \quad (*)$$

The cokernel of $\text{cok } \mu_{iC} \rightarrow \text{cok } \mu_{(i+1)C}$ is $\text{cok } \mu_{C; (i+1)C} \cong H^1(C, \mathcal{O}_C(i+1-a_C) \oplus \mathcal{O}_C(i+1-b_C))$. For $r \geq b_C - 2$, we thus get an upper bound $\dim \text{cok } \mu_{rC} \leq (a-1)(a-2)/2 + (b-1)(b-2)/2$ by adding up the dimensions of the cokernels of $\text{cok } \mu_{C; (i+1)C}$ for $i \leq r$.

Moreover, $C = wL$ for some $w \in W(X)$, hence C is part of an exceptional configuration; i.e., $C = wL, C_1 = wE_1, \dots, C_n = wE_n$. When r is big enough, $|rC - L|$ is nonempty and the fixed part of $|rC - L|$ is $d_1C_1 + \cdots + d_nC_n$, where $d_i = C_i \cdot L$. Thus, by Proposition 3.1, Conjecture 3.4 and the SHGH Conjecture, we have $\dim \text{cok } \mu_{rC} = ((a_1^2 - a_1)/2 + (b_1^2 - b_1)/2) + \cdots + ((a_n^2 - a_n)/2 + (b_n^2 - b_n)/2)$, where (a_i, b_i) is the splitting type for C_i . This gives the inequality

$$((a_1^2 - a_1)/2 + (b_1^2 - b_1)/2) + \cdots + ((a_n^2 - a_n)/2 + (b_n^2 - b_n)/2) \leq (a-1)(a-2)/2 + (b-1)(b-2)/2. \quad (**)$$

This can be enough to determine (a_C, b_C) . For example, let $C = 12L - 5E_1 - 5E_2 - 5E_3 - 4E_4 - 4E_5 - 4E_6 - 4E_7 - 2E_8$. So C comes from a plane curve of degree 12 with three points of multiplicity 5, four of multiplicity 4 and one of multiplicity 2. By Lemma 2.1.2, (a_C, b_C) is either

(6, 6) or (5, 7). Here are the C_i and their types (a_i, b_i) . For convenience we only give the coefficients of $L, -E_1, \dots, -E_8$ followed by the splitting type:

$$\begin{aligned} C_1 &= (5; 2, 2, 2, 2, 1, 2, 2, 1), (2, 3) \\ C_2 &= (5; 2, 2, 2, 2, 2, 1, 2, 1), (2, 3) \\ C_3 &= (5; 2, 2, 2, 1, 2, 2, 2, 1), (2, 3) \\ C_4 &= (5; 2, 2, 2, 2, 2, 2, 1, 1), (2, 3) \\ C_5 &= (4; 2, 2, 2, 1, 1, 1, 1, 1), (2, 2) \\ C_6 &= (3; 1, 1, 2, 1, 1, 1, 1, 0), (1, 2) \\ C_7 &= (3; 2, 1, 1, 1, 1, 1, 1, 0), (1, 2) \\ C_8 &= (3; 1, 2, 1, 1, 1, 1, 1, 0), (1, 2) \end{aligned}$$

Now $((a_1^2 - a_1)/2 + (b_1^2 - b_1)/2) + \dots + ((a_n^2 - a_n)/2 + (b_n^2 - b_n)/2)$ here is 21. But $(a - 1)(a - 2)/2 + (b - 1)(b - 2)/2 = 20$ if we use $(a_C, b_C) = (6, 6)$, so we see that $(a_C, b_C) = (5, 7)$; this is also what we get if we use Macaulay 2 [GS] to compute (a_C, b_C) , using randomly chosen points (so this is a check but not a proof that $(a, b) = (5, 7)$ here).

Also, it follows from (*) and (**) that $\dim \text{cok } \mu_{iC} = \dim \text{cok } \mu_{C;C} + \dots + \dim \text{cok } \mu_{C;iC}$ for all i if (**) is an equality.

Thus (**) can, conjecturally, sometimes tell us both what the splitting type is and what the cokernel is. Moreover, (**) sometimes also applies to exceptional curves. In the example above, $E = C - E_9 - E_{10}$ is exceptional and has the same splitting type as does C , so we get information about E via C .

We have applied our script for computing splitting types to numerous examples. In these examples, (a_C, b_C) always made $(a_C - 1)(a_C - 2)/2 + (b_C - 1)(b_C - 2)/2$ as small as possible subject to (**). (Moreover, in those cases where (**) was not an equality, it was off by exactly 1, and in those cases it always happened that $a_i = b_i$ for all i .) This and Lemma 2.1.2 lead us to make the following conjecture:

Conjecture A2.1.1 *Let $C = wL$ for some $w \in W(X)$, let $d = C \cdot L$, let m be the maximum of $C \cdot E_1, \dots, C \cdot E_n$ and let $C_1 = wE_1, \dots, C_n = wE_n$. Then (a_C, b_C) is the solution (a, b) to $a \leq b$, $\min(m, d - m) \leq a \leq d - m$ and $d = a + b$ which minimizes $(a - 1)(a - 2)/2 + (b - 1)(b - 2)/2$ subject to (**).*

A2.2 Computational Evidence for Conjecture 3.4

There are 2051 exceptional classes of the form $E = tL - m_1E_1 - \dots - m_nE_n$ with $1 \leq t \leq 20$ (taking n to be as large as necessary) and $m_1 \geq \dots \geq m_n \geq 0$. We have applied our splitting script (using randomly chosen points P_i and working in characteristic 31991) to determine the splitting types of all 2051. It turned out that Theorem 3.3(b, c) implies for all but 25 of the 2051 cases that Conjecture 3.4 holds for the given E .

Theorem 3.3(b, c) does not apply in the remaining 25 cases. Here we list these 25 cases, giving $a_E, b_E, E \cdot L$ and $E \cdot E_i$ for all i such that $E \cdot E_i > 0$:

5	8	13	5	5	5	5	5	5	4	1	1	1	1	8	11	19	8	8	7	7	7	6	6	3	2	1	1		
6	9	15	6	6	6	6	5	5	5	2	1	1	1	8	11	19	8	8	7	7	7	7	5	2	2	2	1		
6	9	15	6	6	6	6	6	5	4	1	1	1	1	8	11	19	8	8	8	7	6	6	6	2	2	2	1		
6	10	16	6	6	6	6	6	6	6	1	1	1	1	8	11	19	8	8	8	7	6	6	6	3	1	1	1	1	
7	10	17	7	7	6	6	6	6	6	2	2	2	2	8	11	19	8	8	8	7	7	6	5	2	2	1	1	1	
7	10	17	7	7	6	6	6	6	6	3	1	1	1	8	11	19	8	8	8	7	7	7	4	1	1	1	1	1	1
7	10	17	7	7	7	6	6	6	5	2	2	1	1	8	11	19	8	8	8	8	6	6	5	2	1	1	1	1	1

7 10 17 7 7 7 7 6 5 5 2 1 1 1 1	8 12 20 8 8 8 7 7 7 7 2 2 2 1
7 10 17 7 7 7 7 6 6 4 1 1 1 1 1	8 12 20 8 8 8 7 7 7 7 3 1 1 1 1
7 11 18 7 7 7 7 7 6 6 2 1 1 1 1	8 12 20 8 8 8 8 7 7 6 2 2 1 1 1
7 11 18 7 7 7 7 7 7 5 1 1 1 1 1	8 12 20 8 8 8 8 8 6 6 2 1 1 1 1
8 11 19 7 7 7 7 7 7 7 4 1 1 1	8 12 20 8 8 8 8 8 7 5 1 1 1 1 1
8 11 19 8 7 7 7 7 7 6 3 2 2	

These cases can be checked by directly computing $\dim \operatorname{cok} \mu_F$ for $F = L + iE$. The critical value of i is $i = b_E$ (if Conjecture 3.4 holds for $L + b_E E$, it follows from the sequences (§) that it holds for $L + iE$ for all $0 \leq i \leq E \cdot L$). In principle, one can compute $\dim \operatorname{cok} \mu_F$ by computing a resolution of $I(Z)$ for $Z = m_1 P_1 + \cdots + m_n P_n$ in the usual way, using Gröbner bases. But for all but one of the 25 examples above we found this computation too large to successfully complete using this usual approach. By taking a different geometrically inspired approach we have in fact been able to check that Conjecture 3.4 holds in these 24 other cases. We now discuss this alternate approach for computing the dimension of the cokernel of μ_{L+mE} , when E is an exceptional curve and $0 \leq m \leq L \cdot E$. Since μ_{L+mE} and $\mu_{mE; L+mE}$ have isomorphic cokernels, it is enough to compute the dimension of the cokernel of the latter, which it turns out one can do fairly efficiently, regarding the scheme structure of mE as Proj of a certain graded ring. So first we determine this scheme structure.

Let $E \subset X$ be smooth and rational with $E^2 = -1$. First, we would like to know $\operatorname{Pic}(mE)$, and to determine $h^i(mE, \mathcal{F})$ for both $i = 0$ and $i = 1$ for every line bundle \mathcal{F} on mE .

Using exact sequences of the form $0 \rightarrow \mathcal{O}_X(dL + (j-m)E) \rightarrow \mathcal{O}_X(dL + jE) \rightarrow \mathcal{O}_{mE}(dL + jE) \rightarrow 0$ and $0 \rightarrow \mathcal{O}_X((d-1)L + jE) \rightarrow \mathcal{O}_X(dL + jE) \rightarrow \mathcal{O}_L(dL + jE) \rightarrow 0$ and the cohomology of divisors of the form $dL + jE$, which is known for all d and j , we find that: $h^0(mE, \mathcal{O}_{mE}(dL + jE)) - h^1(mE, \mathcal{O}_{mE}(dL + jE)) = \binom{m+1}{2} + mt$ holds for all $t = E \cdot (dL + jE)$; $h^0(mE, \mathcal{O}_{mE}(dL + jE)) = \binom{m+1}{2} + mt$ (and hence $h^1(mE, \mathcal{O}_{mE}(dL + jE)) = 0$) and $h^0(mE, \mathcal{O}_{mE}(-dL - jE)) = \binom{m-t+1}{2}$ hold if $t \geq 0$; and it follows that $h^1(mE, \mathcal{O}_{mE}(-dL - jE)) = \binom{t}{2}$ holds if $0 \leq t \leq m$, and $h^1(mE, \mathcal{O}_{mE}(-dL - jE)) = tm - \binom{m+1}{2}$ holds if $t \geq m$.

In particular, $h^1(mE, \mathcal{O}_{mE}) = 0$, so it follows that the inclusion $E \subset mE$ induces an isomorphism $\operatorname{Pic}(mE) \rightarrow \operatorname{Pic}(E) = \mathbf{Z}$ (see [A.Ar], (1.3) and (1.4)). It now follows for any line bundle F on X that $h^i(mE, \mathcal{O}_{mE}(F))$ depends only on m , i and $E \cdot F$. In particular, $h^i(mE, \mathcal{O}_{mE}(F)) = h^i(mE, \mathcal{O}_{mE}(t))$, where $t = E \cdot F$ so we define $\mathcal{O}_{mE}(1)$ to be $\mathcal{O}_{mE}(-E)$, and write $h^i(mE, t)$ for $h^i(mE, \mathcal{O}_{mE}(t))$.

To get the scheme structure, we can pick any E which is convenient, since mE is isomorphic for all E on X . Let $\pi : S \rightarrow T$ be the blow up of the point P defined by $x = 0 = y$ in $T = \operatorname{Spec}(K[x, y])$, and take $E = \pi^{-1}(P)$. We can regard S as $\operatorname{Proj}_A(A[u, v]/(xv - uy))$, where $A = K[x, y]$, and E as $\operatorname{Proj}_A(A[u, v]/((xv - uy) + (x, y)))$, which is just $E = \operatorname{Proj}_K(K[u, v])$. Similarly, mE is the fiber $\pi^{-1}(mP)$, so $mE = \operatorname{Proj}_A(A[u, v]/((xv - uy) + (x, y)^m))$. Note that the ring $B = A[u, v]/((xv - uy) + (x, y)^m)$ is graded, with x and y of degree 0 and u and v of degree 1. The homogeneous component B_t for $t \geq 0$ is just the span of the monomials $x^i y^j u^r v^s$ with nonnegative exponents where $r + s = t$, and $i + j < m$ (since we have modded out by $(x, y)^m$). In fact B_t can be identified with $H^0(mE, t)$. However, B has no components of negative degree, whereas $h^0(mE, t) > 0$ for $t > -m$. But note that $x/u = y/v$ on the open set of mE where u and v are neither 0. Thus taking x/u where $u \neq 0$ and y/v where $v \neq 0$ defines an element $\epsilon \in H^1(mE, -1)$. Let $C = K[u, v, \epsilon]/(\epsilon^m)$. This is graded if we take ϵ to have degree -1 . We have a graded injective ring homomorphism $B \rightarrow C$ given by sending $x \mapsto \epsilon u$ and $y \mapsto \epsilon v$, and now we can identify C_t with $H^0(mE, t)$ for all t . In particular, the monomials of the form $\epsilon^i u^r v^s$ with nonnegative exponents satisfying $r + s = t + i$ and $i \leq m - 1$ give a basis for $H^0(mE, t)$, and we can regard mE as

$\text{Proj}_K(C)$.

The map $\mu_{mE;L+mE}$ factors via the restriction $H^0(X, L) \rightarrow H^0(mE, \mathcal{O}_{mE}(L))$ through

$$\mu : H^0(mE, \mathcal{O}_{mE}(L + mE)) \otimes H^0(mE, \mathcal{O}_{mE}(L)) \rightarrow H^0(mE, \mathcal{O}_{mE}(2L + mE)).$$

Thus $\text{Im } \mu_{mE;L+mE}$ is the K -span of the product $(H^0(mE, \mathcal{O}_{mE}(L + mE)))(\text{Im}(H^0(X, L) \rightarrow H^0(mE, \mathcal{O}_{mE}(L))))$ in $H^0(mE, \mathcal{O}_{mE}(2L + mE))$. It has the same dimension as does

$$(H^0(mE_1, \mathcal{O}_{mE_1}(L' + mE_1)))(\text{Im}(H^0(X, L') \rightarrow H^0(mE_1, \mathcal{O}_{mE_1}(L'))))$$

in $H^0(mE_1, \mathcal{O}_{mE_1}(2L' + mE_1))$, where $wL = L'$ for some appropriate Cremona transformation w with $wE = E_1$.

Now choose coordinates on \mathbf{P}^2 such that $K[\mathbf{P}^2] = K[a, b, c]$, where E_1 is the blow up of the point $a = b = 0$. Let $d = E_1 \cdot L' = E \cdot L$. Note that $L' - E_1$ is nef, so $0 \leq (L' - E_1) \cdot L = L' \cdot L - d$; thus $2d - 1 \leq d - 1 + L \cdot L'$. Now, $H^0(mE_1, \mathcal{O}_{mE_1}(2L' + mE_1)) = H^0(mE_1, \mathcal{O}_{mE_1}(2d - m))$, so the monomials of the form $u^i v^j \epsilon^k$ with $2d - m \leq i + j \leq 2d - 1$ and $0 \leq k = i + j - (2d - m)$ give a basis. Thus there is a surjective map of the homogeneous component $((a, b, c)^{2d-1}) \cap ((a, b)^{2d-m})_{d-1+L \cdot L'}$ onto $H^0(mE_1, \mathcal{O}_{mE_1}(2d - m))$, defined by sending $a^i b^j c^{d-1-i-j+L \cdot L'}$ to $u^i v^j \epsilon^{i+j-2d+m}$, where (a, b, c) denotes the ideal in $K[a, b, c]$ generated by a, b and c . Moreover, the kernel of this surjective map is spanned by the monomials $a^i b^j c^{d-1-i-j+L \cdot L'}$ with $i + j \geq 2d$.

Let $r = E_1 \cdot L' = E \cdot L$. Note that the elements of $H^0(X, L')$, regarded as homogeneous polynomials in $K[a, b, c]$, are certain polynomials $f(a, b, c)$ of degree $L \cdot L'$ such that the terms of $f(a, b, 1)$ of least degree have degree r . The image of $f(a, b, c)$ in $H^0(mE_1, \mathcal{O}_{mE_1}(L'))$ (i.e., the restriction of f to mE_1) is just what you get if you formally simplify $(\epsilon^r)(f(u/\epsilon^r, v/\epsilon^r, 1))$. Thus we have a surjection of $((a, b, c)^{d-1}) \cap ((a, b)^{d-m})$ onto $H^0(mE_1, \mathcal{O}_{mE_1}(L' + mE_1))$, defined by sending $a^i b^j c^{d-1-i-j}$ to $u^i v^j \epsilon^{i+j-d+m}$. This gives a surjective map of the homogeneous component $((a, b, c)^{d-1}) \cap ((a, b)^{d-m}) H^0(X, L') + ((a, b)^{2d})_{d-1+L \cdot L'}$ onto $\text{Im } \mu_{mE_1;L'+mE_1}$ with kernel $((a, b)^{2d})_{d-1+L \cdot L'}$. Thus $\dim \text{Im } \mu_{mE_1;L'+mE_1}$ equals $\dim (((a, b, c)^{d-1}) \cap ((a, b)^{d-m}) H^0(X, L') + ((a, b)^{2d})_{d-1+L \cdot L'}) - \dim ((a, b)^{2d})_{d-1+L \cdot L'}$. The hardest part of this calculation is finding $H^0(X, L')$. It can be done either using a Gröbner basis calculation (but one much smaller than what is needed to calculate the dimension of the image of $\mu_{mE;L+mE}$ directly) or by applying w to a basis for $H^0(X, L)$.

A2.3 Scripts with Examples

In this section we give Macaulay 2 scripts that implement the computations discussed in this paper, followed by examples of their use.

The Macaulay script file "BMSmacaulayscripts":

```
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
```

```
R=ZZ/31991[x,y, MonomialSize=>16, MonomialOrder => ProductOrder {1,1}];
S=ZZ/31991[a,b,c, MonomialSize=>16, MonomialOrder => ProductOrder {1,1,1}];

-- input: list of form {m1,...,mn} giving the multiplicities
-- of a fat point subscheme Z = m_ip_1+...m_np_n of P2, where
-- p_i are randomly chosen (and hence probably general) points.
-- output: the ideal of Z is returned and the Maculay formatted
-- graded Betti numbers are printed. This script is based on a script
-- provided to us by H. Schenck.

mixer = (l)->{(i:=0;
  bb:=ideal (matrix {{l}}**S);
  I:=ideal(a,b);
  scan(#l, i->(
    bb=intersect(bb,I^(l#i));
    f:=random(S^1,S^-1)});
```

```

        g:=random(S^1,S^{-1});
        I=(ideal (f | g)));
        print betti res coker gens bb;
bb)

--decomp: prints a decomposition F=H+N for any divisor class F in Psi
--as described in the print statements of the script.
--input: decomp(l), where l={d,{m1,...,mn}}, signifying the divisor
--class F = dE_0-(m_1E_1+...+m_nE_n).

decomp = (l) -> (
<< "Let Psi be the subsemigroup of divisor classes generated by " << endl;
<< "exceptional classes and by -K. For any divisor class F, this " << endl;
<< "script determines if F is in Psi, and if so gives a decomposition" << endl;
<< "F=H+N, where N is a sum of exceptionals orthogonal to each other and to H" << endl;
<< "and H is in Psi but H.E >= 0 for all exceptionals E. The point of this" << endl;
<< "is that dim |F| = dim |H|, and conjecturally dim |H| = (H^2-H.K)/2." << endl << endl;
i:=0;
j:=0;
w:=1;
v:={};
ww:={};
ex:={};
mult:=0;
tmp:=fundomfully(l);
if tmp#0<tmp#1#0 or tmp#0<0 then (<< "Your class is not in Psi." << endl) else (
  if #l#1<3 then w={l#0,join(l#1,{0,0,0})};
  d:=3*(#(w#1)); --define an element {d,v} in fundamental domain
  scan(#(w#1), i->(v=join(v,{#(w#1)} - i)));
  ww=fundombbothfully(w,{d,v});
  if ww#0#1 == zr(ww#0#1) then (<< "N = 0" << endl) else (
    << "N is a sum of the following fixed exceptional classes:" << endl;
    if (ww#0#0) - (ww#0#1#0) - (ww#0#1#1) < 0 then (
      scan(#(w#1), j->(if j<=1 then ex=join(ex,{1}) else ex=join(ex,{0})));
      ex={1,ex};
      ex=(fundombbothfully(ww#1,ex))#1;
      <<ex<<" is a fixed component of multiplicity ";
      <<(ww#0#1#0)+(ww#0#1#1)-(ww#0#0)<<endl;
      ex={(ww#0#0)-(ww#0#1#1),(ww#0#0)-(ww#0#1#0)};
      scan(#(w#1), j->(if j>1 then ex=join(ex,{ww#0#1#j})));
      ww={{2*(ww#0#0)-(ww#0#1#0)-(ww#0#1#1),ex},ww#1}};
    scan(#(w#1), i->(
      if (ww#0#1)#i<0 then (
        ex={};
        mult--(ww#0#1)#i;
        scan(#(w#1), j->(if j==i then ex=join(ex,{-1}) else ex=join(ex,{0})));
        ex={0,ex};
        ex=(fundombbothfully(ww#1,ex))#1;
        << ex << " is a fixed component of multiplicity " << mult << endl));
      << endl << "and H = " << (fundombbothfully(ww#1,{ww#0#0,zr(ww#0#1)}))#1 << endl));
  -- zr: replaces negative values in a list l by zeroes.
  -- Call it as zr(l) where l is a list of integers.

zr = (l) -> (
v:={};
i:=0;
scan(#l, i->(
  if l#i<0 then v=join(v,{0}) else v=join(v,{l#i}));
v)

-- homcompdim: computes e(F_t(Z)), the expected dimension of a component I_d
-- of a fat points ideal I corresponding to a fat point subscheme Z of general
-- points taken with multiplicities m_1, ..., m_n. Call it as
-- homcompdim({d,{m_1,...,m_n}}); the output is the SHGH conjectural
-- dimension of I_d, which is the actual dimension if n < 10.

homcompdim = (l) -> (
h:=0;
i:=0;
w:=1;
if #l#1<3 then w={l#0,join(l#1,{0,0,0})};
w=fundomfully(w);
d:=w#0;
w=fundomfully({d,zr(w#1)});
d=w#0;
v:=zr(w#1);
if d<0 then h=0 else (
  tmp:=0;
  scan(#v, i->(tmp = v#1*v#i+v#i+tmp));
  h=floor((d*d+3*d+2-tmp)/2);
  if h < 0 then h=0);
h)

-- input: list of form l = {d,{m1,...,mn}} which reduces by
-- quadratic transformations to {1,{1,1,0,...,0}}. Thus there is a
-- birational morphism f: P1 -> P2 parametrizing the curve {d,{m1,...,mn}}.
-- output: returns the result of restrictionIdeal(l,0)
-- and prints the degrees d1 and d2 of the splitting of the bundle f*\Omega_{P2}.
-- Note that d1+d2=3d. The script uses the fact
-- that the splitting is known if d-m1 <= m1 + 1 to avoid doing any
-- computation in this case.

```

```

bundledecomp = (l) -> (
  if #(l#1) < 3 then (
    << "number of points is " << #(l#1) << " but must be 3 or more; please try again" << endl) else
  (
    v:=l#1;
    ev:={1,1};
    scan(#v-2, i->(ev=join(ev,{0})));
    ev={1,ev}; -- ev is the exceptional curve L-E_1-E_2
    if ev != fundom(l) then (
      << "Input class is not an exceptional class; ignore output" << endl) else (
        d1:=l#0;
        m1:=l#1#0;
        if d1-m1<m1+1 then << l << " cotangent splitting: " << 2*d1-m1 << " " << d1+m1 << endl;
        if d1-m1==m1+1 then << l << " cotangent splitting: " << d1+m1 << " " << 2*d1-m1 << endl;
        if d1-m1>m1+1 then (
          ii:=restrictionIdeal(l,0);
          I:=(ii)_0;
          J:=(ii)_1;
          z:=res I;
          d1=(degree (z_2)_0)_0;
          d2=(degree (z_2)_1)_0;
          << l << " cotangent splitting: " << d1 << " " << d2 << endl;
          ii))))
    -- lstprmt: arranges the elements of the list l1 in descending order,
    -- and applies the same permutation to l2

    lstprmt = (l1,l2) -> (
      tmpv1:=l1;
      tmpv2:=l2;
      v1:=l1;
      v2:=l2;
      i:=0;
      j:=0;
      k:=0;
      scan(#l1, i->(scan(#l1, j->(
        if tmpv1#i < tmpv1#j then (
          if i < j then (
            k=-1;
            v1={};
            v2={};
            while(k<#l1-1) do (k=k+1;
              if k==i then (v1=join(v1,{tmpv1#j});
                v2=join(v2,{tmpv2#j});) else (
                  if k==j then (v1=join(v1,{tmpv1#i});
                    v2=join(v2,{tmpv2#i});) else (v1=join(v1,{tmpv1#k});
                      v2=join(v2,{tmpv2#k})))));
            tmpv1=v1;
            tmpv2=v2))))));
      {v1,v2})

    -- quad: performs a quadratic transform on a divisor class dE_0-(m_1E_1+...+m_nE_n).
    -- Call it as quad({d,{m1,...,mn}}). The output is
    -- {2d-m1-m2-m3,{d-m2-m3,d-m1-m3,d-m1-m2,m4,...,mn}}.

    quad = (l) -> (
      i:=0;
      w:=1;
      if #l#1<3 then w={l#0,join(l#1,{0,0,0})};
      v:={w#0 - w#1#1 - w#1#2,w#0 - w#1#0 - w#1#2,w#0 - w#1#0 - w#1#1};
      scan(#w#1, i->(if i>2 then v=join(v,{w#1#i})));
      v:={2*(w#0) - w#1#0 - w#1#1 - w#1#2,v};
      v)

    -- fundomboth: applies quad iteratively to l1 to reduce l1 to standard form
    -- (this being l1 = {x,{y1,y2,...,yn}}, where either x <= 1 and y1>y2>= ...
    -- or y1>y2>= ... and x >= y1 + y2 + y3; when l1 starts out being an exceptional
    -- class with x >= 1, it ends up as {1,{1,1,0,...,0}}).
    -- Note: fundomboth applies the same operations to l2 as it does to l1
    -- (note: l1 and l2 should have the same number of entries).
    -- If l2 starts out in the fundamental domain,
    -- and {l1',l2'}=fundomboth(l1,l2), then {l2,g^{-1}l}=fundomboth(l2',l).
    -- This allows one to compute the action of g^{-1}.

    fundomboth = (l1,l2) -> (
      w1:=l1;
      w2:=l2;
      v:={};
      if #l1#1<3 then (
        w1={l1#0,join(l1#1,{0,0,0})};
        w2={l2#0,join(l2#1,{0,0,0})};
      );
      v=lstprmt(w1#1,w2#1);
      w1={w1#0,v#0};
      w2={w2#0,v#1};
      while ((w1#0 < w1#1#0 + w1#1#1 + w1#1#2) and (w1#0 > 1)) do (
        w1=quad(w1);
        w2=quad(w2);
        v=lstprmt(w1#1,w2#1);

```

```

w1={w1#0,v#0};
w2={w2#0,v#1};
);
{w1,w2}

-- fundombothfully: this is the same as fundomboth except
-- l1 = {x,{y1,y2,...,yn}} is reduced to the standard form
-- where either x < 0 and y1>y2>= ... , or y1>y2>= ... and x >= y1 + y2 + y3;
-- thus an exceptional class always reduces to {0,{0,...,0,-1}}.

fundombothfully = (l1,l2) -> (
w1:=l1;
w2:=l2;
v:={};
if #l1#1<3 then w1={l1#0,join(l1#1,{0,0,0})};
if #l2#1<3 then w2={l2#0,join(l2#1,{0,0,0})};
v=prmtboth(w1#1,w2#1);
w1={w1#0,v#0};
w2={w2#0,v#1};
while ((w1#0 < w1#1#0 + w1#1#1 + w1#1#2) and (w1#0 >= 0)) do (
w1=quad(w1);
w2=quad(w2);
v=prmtboth(w1#1,w2#1);
w1={w1#0,v#0};
w2={w2#0,v#1};
{w1,w2}

-- prmtboth: arranges the elements of the list l1 in descending order,
-- and applies the same permutation to l2
-- Call it as prmt(l1,l2) where l1 and l2 are lists of integers.

prmtboth = (l1,l2) -> (
tmpv1:=l1;
tmpv2:=l2;
v1:=l1;
v2:=l2;
i:=0;
j:=0;
k:=0;
scan(#l1, i->(scan(#l1, j->(
if tmpv1#i < tmpv1#j then (
if i < j then (
k=-1;
v1={};
v2={};
while(k<#l1-1) do (k=k+1;
if k=i then (v1=join(v1,{tmpv1#j});
v2=join(v2,{tmpv2#j})) else (
if k=j then (v1=join(v1,{tmpv1#i});
v2=join(v2,{tmpv2#i})) else (v1=join(v1,{tmpv1#k});
v2=join(v2,{tmpv2#k})))));
tmpv1=v1;
tmpv2=v2))))));
{v1,v2}

-- fundom: Call it as fundom({d,{m1,...,mn}}). The output is a new
-- list {d',{m1',...,mn'}}; the class d'E_0-(m_1E_1+...+m_nE_n) is
-- equivalent via Cremona transformations to d'E_0-(m_1'E_1+...+m_n'E_n),
-- where d' is the smallest positive integer possible.

fundom = (l) -> (
(fundomboth(l,l))#0)

-- fundomfully: Same as fundom except uses fundombothfully instead of fundomboth.

fundomfully = (l) -> (
(fundombothfully(l,l))#0)

-- input: list {m1,...,mn} of integers and a 3 by n matrix M
-- output: list of integers in descending order, with matrix N whose columns
-- are those of M permuted by the same permutation that arranged the integers

matprmt = (l,M) -> (
i:=0;
tmpprm:={};
scan(#l, i->(tmpprm=join(tmpprm,{i})));
myoutput:=lstprmt(l,tmpprm);
N:=M_(myoutput#1);
{myoutput#0,N})

-- input: l is an exceptional class, MM is a matrix of n=#l#1 points,
-- and flag is an integer; if flag=1, then extra steps are printed out.
-- Making MM an input allows use of a given set of points, if desired.
-- Output: the parametrization of the curve l

myloop = (l, MM, flag) -> (
mymaps:={};
tmpM:=id(S^3);
npts:=#l#1;
M:=MM;
ev:={1,1};

```

```

i:=0;
scan(npts-2, i->(ev=join(ev,{0})));
ev={1,ev}; -- ev is the exceptional curve L-E_1-E_2
fe:={};
scan(npts, i->(fe=join(fe,{npts-i})));
fe={3*npts-2,fe}; -- element of fundamental domain
tfe:=(fundombboth(1,fe))_1; -- this element is a translate of previous fe;
-- translating fe back to fund dom takes ev to 1

v:={};
f:=(x,y,0*x); -- this is the parametrization of ev, where ev is z=0 in P2
if flag==1 then << "test elt: " << tfe << " exc class: " << ev << endl;
if flag==1 then << "the points in P2: " << M << endl;
if flag==1 then << "Parametrization of exc class: " << toString f << endl;
line0:=det(submatrix(M,{1,2})|matrix{{a},{b},{c}});
line1:=det(submatrix(M,{0,2})|matrix{{a},{b},{c}});
line2:=det(submatrix(M,{0,1})|matrix{{a},{b},{c}});
g:=map(R,S,f); -- this is the map S=k[P2]->R=k[P1] parametrizing ev
K:=kernel(g); -- compute equation of image in P2 of ev
I:=ideal(f#0,f#1,f#2);
J:=saturate I;
if flag==1 then << "Equation of image of exc class: " << toString K << endl;
while(tfe != fe) do (
v=lstprmt(tfe#1,ev#1);
M=(matprmt(tfe#1,M))#1;
tfe={tfe#0,v#0};
ev={ev#0,v#1};
if (tfe#0 < tfe#1#0 + tfe#1#1 + tfe#1#2) then (
line0:=det(submatrix(M,{1,2})|matrix{{a},{b},{c}});
-- << "line0: " << line0 << endl;
line1:=det(submatrix(M,{0,2})|matrix{{a},{b},{c}});
-- << "line1: " << line1 << endl;
line2:=det(submatrix(M,{0,1})|matrix{{a},{b},{c}});
-- << "line2: " << line2 << endl;
-- << "param before sat: " << {g(line1*line2),g(line0*line2),g(line1*line0)} << endl;
I={g(line1*line2),g(line0*line2),g(line1*line0)};
J=gcd(gcd(I_0,I_1),I_2); -- I typically has base points that need to be removed; J defines them
-- << "J: " << J << endl;
f={I_0/J,I_1/J,I_2/J}; -- this removes the base points
-- << "param after sat: " << f << endl;
g:=map(R,S,f);
if flag==1 then K=kernel(g); -- to see equations of the images of the curves, uncomment out: THIS IS SLOW
tfe=quad(tfe);
ev=quad(ev);
-- << "After perm: " << M << endl;
mymaps={}; -- for each point, define map given by evaluation at that point
scan(npts, i->(mymaps=join(mymaps,{map(S,S,{M_(0,i), M_(1,i), M_(2,i)}))}));
tmpM=id_(S^3);
scan(npts, i->(if i>2 then tmpM=tmpM|matrix{{mymaps_i(line1*line2)},{mymaps_i(line0*line2)},{mymaps_i(line1*line0)}}));
M=tmpM;
-- uncomment the next line to see which points are on the given curve:
-- scan(npts, i-><< "pt " << i << " : " << mymaps_i(K_0) << endl ));
);
if flag==1 then << "test elt: " << tfe << " exc class: " << ev << endl;
if flag==1 then << "the points in P2: " << M << endl;
if flag==1 then << "Parametrization of exc class: " << toString f << endl;
if flag==1 then << "Equation of image of exc class: " << toString K << endl; -- to see equations of the images of the curves, uncomment out
);
f)

```

```

-- input: list of form l = {d,{m1,...,mn}} which reduces by
-- quadratic transformations to {1,{1,1,0,...,0}}. Thus there is a
-- birational morphism f: P1 -> P2 parametrizing the curve {d,{m1,...,mn}}.
-- output: prints res of ideal on P1 of the linear forms in P2 pulled back via f
-- to the curve {d,{m1,...,mn}}.
-- Note: flag is an integer; if flag=1, then extra steps are printed out.

```

```

restrictionIdeal = (1,flag) -> (
I:=ideal(0*x);
J:=ideal(0*x);
M:=matrix(table(3,#1#1-3,(i,j)->random(0,S)));
M=id_(S^3)|M;
f:=myloop(1,M,0);
I:=ideal(f_0,f_1,f_2);
if flag==1 then << "Resolution of ideal of restrictions" << endl << betti res coker gens I << endl;
myM:=syz gens I;
J:=ideal(myM_(0,0),myM_(1,0),myM_(2,0));
if flag==1 then << "Resolution of least degree relation ideal" << endl << betti res coker gens J << endl;
{I,J})

```

```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

```

Here is sample Macaulay session demonstrating use of the scripts (interspersed comments which are not part of the session are indented)

```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

```

```

load "BMSmacaulayscripts"
decomp({101,{50,50,38,38,26,26,22,18,14,14}})
homcompdim({101,{50,50,38,38,26,26,22,18,14,14}})
decomp({102,{50,50,38,38,26,26,22,18,14,14}})
homcompdim({102,{50,50,38,38,26,26,22,18,14,14}})
homcompdim({38, {18, 18, 14, 14, 10, 10, 8, 8, 6, 6}})
bundledcomp({8, {4, 4, 3, 3, 2, 2, 1, 2, 1, 1}})
decomp({103,{50,50,38,38,26,26,22,18,14,14}})
homcompdim({103,{50,50,38,38,26,26,22,18,14,14}})
homcompdim({104,{50,50,38,38,26,26,22,18,14,14}})
l=homcompdim({37, {17, 18, 14, 14, 10, 10, 8, 8, 6, 6}})
Proof: decomp({37, {17, 18, 14, 14, 10, 10, 8, 8, 6, 6}})
q=homcompdim({38, {19, 18, 14, 14, 10, 10, 8, 8, 6, 6}})
Proof: decomp({38, {19, 18, 14, 14, 10, 10, 8, 8, 6, 6}})
homcompdim({39, {18, 18, 14, 14, 10, 10, 8, 8, 6, 6}})
time mixer({8, 8, 6, 6, 4, 4, 2, 4, 2, 2});
time mixer({24, 24, 18, 18, 12, 12, 12, 6, 6, 6});
fundombothfully({8,{4,4,3,3,2,2,2,1,1,1}},{1,{0,0,0,0,0,0,0,0,0}})
time II=mixer({8, 5, 5, 4, 4, 3, 2, 2, 2, 1});
I=ideal(II_0,II_1,II_2);
J=intersect((ideal(a,b,c))^7,(ideal(a,b))^2);
K=I*J+(ideal(a,b))^16;
time hilbertFunction(20,K)
L=(ideal(a,b))^16;
time hilbertFunction(20,L)

```

In this example, we want to find the graded Betti numbers for $Z=Z(\{50,50,38,38,26,26,22,18,14,14\})$. Here's how we can do so using the M2 scripts "BMSmacaulayscripts". This shows an actual M2 session; our comments are interspersed and indented.

```

[G4desktop:~/Desktop] bh% M2
Macaulay 2, version 0.9.2
--Copyright 1993-2001, D. R. Grayson and M. E. Stillman
--Singular-Factory 1.3c, copyright 1993-2001, G.-M. Greuel, et al.
--Singular-Libfac 0.3.2, copyright 1996-2001, M. Messollen

i1 : load "BMSmacaulayscripts"
--loaded BMSmacaulayscripts

First find the least degree t such that F_t=
{t,{50,50,38,38,26,26,22,18,14,14}} is in Psi.
The answer is t = 102. (If F_t is not in Psi
then it is known that h^0(X, F_t)=0.)

i2 : decomp({101,{50,50,38,38,26,26,22,18,14,14}})
Let Psi be the subsemigroup of divisor classes generated by
exceptional classes and by -K. For any divisor class F, this
script determines if F is in Psi, and if so gives a decomposition
F=H+N, where N is a sum of exceptionals orthogonal to each other and to H
and H is in Psi but H.E >= 0 for all exceptionals E. The point of this
is that dim |F| = dim |H|, and conjecturally dim |H| = (H^2-H.K)/2.

Your class is not in Psi.

o2 = stdio
o2 : File

-- the standard input output file

The next command computes h^0(X, F_t) for the given F_t,
assuming the SHGH conjecture. As expected, we get 0 for t<102.

i3 : homcompdim({101,{50,50,38,38,26,26,22,18,14,14}})
o3 = 0

We now see that F_t is in Psi for t=102, and we get
what is essentially a Zariski decomposition of F_t;
we get F_t = H + N, where N is a sum of orthogonal exceptional
curves, and H is a class such that H.E >= 0 for all exceptional
curves E and such that H.N=0. Thus h^0(X, F_t) = h^0(X, H).

i4 : decomp({102,{50,50,38,38,26,26,22,18,14,14}})
Let Psi be the subsemigroup of divisor classes generated by
exceptional classes and by -K. For any divisor class F, this
script determines if F is in Psi, and if so gives a decomposition
F=H+N, where N is a sum of exceptionals orthogonal to each other and to H
and H is in Psi but H.E >= 0 for all exceptionals E. The point of this
is that dim |F| = dim |H|, and conjecturally dim |H| = (H^2-H.K)/2.

N is a sum of the following fixed exceptional classes:
{8, {4, 4, 3, 3, 2, 2, 1, 2, 1, 1}} is a fixed component of multiplicity 2
{8, {4, 4, 3, 3, 2, 2, 2, 1, 1, 1}} is a fixed component of multiplicity 6
and H = {38, {18, 18, 14, 14, 10, 10, 8, 8, 6, 6}}

o4 = stdio

```

```

o4 : File
-- the standard input output file

Here we assume the SHGH conjecture to check that
the conjecture does in fact agree that  $h^0(X, F_t) = h^0(X, H)$ .
[In fact, the SHGH conjecture holds for H since the command
fundomfully({38, {18, 18, 14, 14, 10, 10, 8, 8, 6, 6}}) gives
outputs {6, {2, 2, 2, 2, 2, 2, 2, 0, 0}}; i.e., H reduces by Cremona
transformations to sextics through 8 general points with multiplicity 2,
and it is known that this reduction preserves  $h^0$ , and that
8 general points of multiplicity 2 impose independent conditions on sextics,
hence that there are 4, in agreement with the SHGH conjecture.]

i5 : homcompdim({102, {50, 50, 38, 38, 26, 26, 22, 18, 14, 14}})
o5 = 4

i6 : homcompdim({38, {18, 18, 14, 14, 10, 10, 8, 8, 6, 6}})
o6 = 4

Next we determine how the bundle  $\Omega(1)|_E$  splits for
 $E = \{8, \{4, 4, 3, 3, 2, 2, 1, 2, 1, 1\}\}$ ; i.e., we find
 $a_E$  and  $b_E$  where  $\Omega(1)|_E = \mathcal{O}_E(-a_E) \oplus \mathcal{O}_E(-b_E)$ .
Computationally bundledecomp gives  $\Omega|_E = \mathcal{O}_E(-12) \oplus \mathcal{O}_E(-12)$ ,
hence  $a_E = 12 - L.E = 4$  and  $b_E = 12 - L.E = 4$ . In this case
we didn't need to rely on the M2 computation since we know
 $a_E + b_E = d$ , where  $d = L.E$  and we know  $\min(d-m, m) \leq a_E \leq d-m$ ,
where  $m$  is the maximum multiplicity. Thus  $a_E = 4 = b_E$ ,
in agreement with the computation.

i7 : bundledecomp({8, {4, 4, 3, 3, 2, 2, 1, 2, 1, 1}})
{8, {4, 4, 3, 3, 2, 2, 1, 2, 1, 1}} cotangent splitting: 12 12

Here we check that  $F_t$  is fixed component free for  $t=103$ .
This already follows from  $F_{102} = H + N$ , since both
H and (it is easy to see)  $L + N$  are fixed component free.

i8 : decomp({103, {50, 50, 38, 38, 26, 26, 22, 18, 14, 14}})
Let Psi be the subsemigroup of divisor classes generated by
exceptional classes and by  $-K$ . For any divisor class F, this
script determines if F is in Psi, and if so gives a decomposition
 $F = H + N$ , where N is a sum of exceptionals orthogonal to each other and to H
and H is in Psi but  $H.E \geq 0$  for all exceptionals E. The point of this
is that  $\dim |F| = \dim |H|$ , and conjecturally  $\dim |H| = (H^2 - H.K)/2$ .

N = 0

and H = {103, {50, 50, 38, 38, 26, 26, 22, 18, 14, 14}}

o8 = stdio

o8 : File
-- the standard input output file

Here we check what the SHGH conjecture says about  $h^0(X, F_t)$ 
for  $t=103$ . Since  $F_{103}$  is effective but fixed component free
we expect that the points impose independent conditions
on 103-ics, which would give  $h^0(X, F_{103})=92$ .
In fact, given that  $h^1(X, H)=0$ , it is easy to see that
 $h^1(X, F_{103})$  is also 0, and hence that  $h^0(X, F_{103})=92$ 
is indeed correct.

i9 : homcompdim({103, {50, 50, 38, 38, 26, 26, 22, 18, 14, 14}})
o9 = 92

Likewise it is now easy to see that the SHGH conjecture
gives the correct value for  $h^0(X, F_{104})$ , and for all  $t > 104$ .

i10 : homcompdim({104, {50, 50, 38, 38, 26, 26, 22, 18, 14, 14}})
o10 = 197

We now start computing the graded Betti numbers  $g_{t+1} = \dim \text{cok } \mu_{F_t}$ .
It is easy to see that  $g_{102} = 4$ , since
 $h^0(X, F_{102})=4$  and since  $h^0(X, F_t)=0$  for  $t < 102$ .
So now we consider  $g_{t+1} = \dim \text{cok } \mu_{F_t}$  for  $t=102$ .
If  $F = H + N$ , where H and N are effective and N is in the fixed part of  $|F|$ ,
we have  $\text{Im } \mu_F \subset \text{Im } \mu_H$ , hence
 $\dim \text{cok } \mu_F = (\dim \text{cok } \mu_H) + (h^0(X, F+L) - h^0(X, H+L))$ .
We also have the bound  $\dim \ker \mu_H \leq h^0(X, H-L) + h^0(X, H-(L-E_1))$ .
The next three commands check that  $h^0(X, H-L) = 0$  and  $h^0(X, H-(L-E_1)) = 0$ ,
hence that  $\ker \mu_H = 0$ , hence that
 $\dim \text{cok } \mu_H = h^0(X, H+L) - 3h^0(X, H) = 44 - 3*4 = 32$ ,
so that  $g_{103} = 32 + (92-44) = 80$ .

i11 : homcompdim({37, {17, 18, 14, 14, 10, 10, 8, 8, 6, 6}})
o11 = 0

```

```

i12 : decomp({37, {17, 18, 14, 14, 10, 10, 8, 8, 6}})
Let Psi be the subsemigroup of divisor classes generated by
exceptional classes and by -K. For any divisor class F, this
script determines if F is in Psi, and if so gives a decomposition
F=H+N, where N is a sum of exceptionals orthogonal to each other and to H
and H is in Psi but H.E >= 0 for all exceptionals E. The point of this
is that dim |F| = dim |H|, and conjecturally dim |H| = (H^2-H.K)/2.

Your class is not in Psi.

o12 = stdio

o12 : File
-- the standard input output file

i13 : homcompdim({38, {19, 18, 14, 14, 10, 10, 8, 8, 6}})

o13 = 0

i14 : decomp({38, {19, 18, 14, 14, 10, 10, 8, 8, 6}})
Let Psi be the subsemigroup of divisor classes generated by
exceptional classes and by -K. For any divisor class F, this
script determines if F is in Psi, and if so gives a decomposition
F=H+N, where N is a sum of exceptionals orthogonal to each other and to H
and H is in Psi but H.E >= 0 for all exceptionals E. The point of this
is that dim |F| = dim |H|, and conjecturally dim |H| = (H^2-H.K)/2.

Your class is not in Psi.

o14 = stdio

o14 : File
-- the standard input output file

i15 : homcompdim({39, {18, 18, 14, 14, 10, 10, 8, 8, 6}})

o15 = 44

We now want to compute g_{104}. Regularity considerations
(i.e., the fact that h^{-1}(X,F_{103})=0) ensure that g_t=0 for t>104.
To find g_{104} we use that fact that
dim cok \mu_F_{103} = (dim cok \mu_N) = (dim cok \mu_{L+2E'}) + (dim cok \mu_{L+6E''}).
Computing the graded Betti numbers for I(Z) directly is at the edge of or
perhaps just beyond what we can do computationally, but we can salvage
(dim cok \mu_{L+2E'}) and (dim cok \mu_{L+6E''}) from computing
the graded Betti numbers for Z' = Z({8, 8, 6, 6, 4, 4, 2, 4, 2, 2})
and Z'' = Z({24, 24, 18, 18, 12, 12, 12, 6, 6, 6}), since each of these
is a much smaller computation than doing I(Z). [Note that we get
Z' from 2E' = {16, {8, 8, 6, 6, 4, 4, 2, 4, 2, 2}} and
Z'' from 6E'' = {48, {24, 24, 18, 18, 12, 12, 12, 6, 6, 6}}.]
What we find computationally (using characteristic p = 31991 and
randomly chosen points is that
(dim cok \mu_{L+2E'}) + (dim cok \mu_{L+6E''}) = 0 + 2, where
the 0 is the entry in row 17 and column 2 of the output
of mixer({8, 8, 6, 6, 4, 4, 2, 4, 2, 2}); and the 2 is the entry
in row 49 and column 2 of mixer({24, 24, 18, 18, 12, 12, 12, 6, 6, 6}); .

i16 : time mixer({8, 8, 6, 6, 4, 4, 2, 4, 2, 2});
total: 1 16 15
0: 1 . .
1: . . .
2: . . .
3: . . .
4: . . .
5: . . .
6: . . .
7: . . .
8: . . .
9: . . .
10: . . .
11: . . .
12: . . .
13: . . .
14: . . .
15: . 1 .
16: . 15 14
17: . . 1
-- used 1.12 seconds

o16 : Ideal of S

i17 : time mixer({24, 24, 18, 18, 12, 12, 12, 6, 6, 6});
total: 1 36 35
0: 1 . .
1: . . .
2: . . .
3: . . .
4: . . .
5: . . .

```



```

6: . . .
7: . . .
8: . . .
9: . . .
10: . . .
11: . . .
12: . . .
13: . . .
14: . . .
15: . . .
16: . . .
17: . . .
18: . . .
19: . . .
20: . . .
21: . . .
22: . . .
23: . . .
24: . . .
25: . . .
26: . . .
27: . . .
28: . . .
29: . . .
30: . . .
31: . . .
32: . . .
33: . . .
34: . . .
35: . . .
36: . . .
37: . . .
38: . . .
39: . . .
40: . . .
41: . . .
42: . . .
43: . . .
44: . . .
45: . . .
46: . . .
47: . 1 .
48: . 33 20
49: . 2 15
-- used 70.78 seconds

o17 : Ideal of S

Thus  $g_{104} = 2$  so we are now done. Unfortunately, although computing
dim cok  $\mu_N$  component by component is much easier than computing
dim cok  $\mu_{F,t}$  directly, it is not hard to give an  $m$  and an  $E$  such
that computing dim cok  $\mu_{L+mE}$  is itself beyond what current equipment
can do. So we now give a more efficient approach. For comparison,
we apply it to  $6E$ . Using the direct approach above using mixer
took 70 seconds. We now present a method that is an order of magnitude
or two faster.

This method depends on using the isomorphism  $\text{cok } \mu_{L+mE}$  with
 $\text{cok } \mu_{L+mE, mE}$  where  $\mu_{L+mE, mE}$  is the map
 $H^0(mE, \mathcal{O}_{mE}(L+mE)) \otimes H^0(X, L) \rightarrow H^0(mE, \mathcal{O}_{mE}(2L+mE))$ .

Assume  $L \cdot E = d$  and that  $0 < m \leq d$ . The inclusion  $E \rightarrow mE$  of schemes
induces an isomorphism  $\mathbb{Z} = \text{Pic}(E) = \text{Pic}(mE)$ . Let  $\mathcal{O}_{mE}(1)$  be the ample
generator of  $\text{Pic}(mE)$ . Then we can regard  $mE$  as the scheme  $\text{Proj}(k[u,v,e]/(e^m))$ 
under the grading such that  $\deg u = \deg v = -\deg e = 1$ .
In particular,  $H^0(mE, \mathcal{O}_{mE}(t))$  is the  $k$ -span of the monomials
 $u^i v^j e^s$ , such that  $i, j$  and  $s$  are nonnegative,  $i+j+s = t$  and  $i < m$ .
It follows that  $h^0(mE, \mathcal{O}_{mE}(t)) = \binom{m+1}{2} + mt$  for  $t \geq 0$ 
and  $h^0(mE, \mathcal{O}_{mE}(t)) = \binom{m+1+t}{2}$  for  $t < 0$ .

The map  $\mu_{L+mE, mE}$  factors via the restriction  $H^0(X, L) \rightarrow H^0(mE, \mathcal{O}_{mE}(L))$ 
through  $H^0(mE, \mathcal{O}_{mE}(L+mE)) \otimes H^0(mE, \mathcal{O}_{mE}(L)) \rightarrow H^0(mE, \mathcal{O}_{mE}(2L+mE))$ .
Thus to determine  $\text{Im } \mu_{L+mE, mE}$  we need to determine
 $\text{Im}(H^0(X, L) \rightarrow H^0(mE, \mathcal{O}_{mE}(L)))$ . After applying an appropriate Cremona
transformation  $w$  we have  $wE = E_1$ ; let  $wL = L'$ . We may assume that
 $E_1$  is the blow up of the point  $a=b=0$  with respect to  $k[P_2]=S=k[a,b,c]$ .
Let  $r = E_1 \cdot L'$ . Then the elements of  $H^0(X, L')$ , regarded as homogeneous
elements polynomials in  $S$ , are polynomials  $f(a,b,c)$  of degree  $L \cdot L'$ 
such that the terms of  $f(a,b,1)$  of least degree have degree  $r$ .
The image of  $f(a,b,c)$  in  $H^0(mE, \mathcal{O}_{mE}(L))$  (i.e., the restriction
of  $f$  to  $mE$ ) is just what you get if you formally simplify  $(e^r)(f(u/e^r, v/e^r, 1))$ .
To compute  $\text{dim Im } \mu_{L+mE, mE}$ , we just need to find the  $k$ -span
of the product  $(H^0(mE, \mathcal{O}_{mE}(L+mE))) (\text{Im}(H^0(X, L) \rightarrow H^0(mE, \mathcal{O}_{mE}(L))))$  in  $S$ .
Alternatively, it is easier to implement in Macaulay the following way.
We have a surjection of  $(\text{ideal}(a,b,c)^{\{d-1\}} \cap \text{ideal}(a,b)^{\{d-m\}})$ 
onto  $H^0(mE, \mathcal{O}_{mE}(L+mE))$  defined by sending  $a^i b^j c^{d-1-i-j} \rightarrow u^i v^j e^{i+j-d+m}$ .
Thus there is a surjective map of the homogeneous component
 $((\text{ideal}(a,b,c)^{\{d-1\}} \cap \text{ideal}(a,b)^{\{d-m\}}) H^0(X, L') + (\text{ideal}(a,b)^{\{2d\}})_{d-1+L \cdot L'})$ 
onto  $\text{Im } \mu_{L+mE, mE}$  with kernel  $(\text{ideal}(a,b)^{\{2d\}})_{d-1+L \cdot L'}$ .
Thus  $\text{dim Im } \mu_{L+mE, mE}$  equals
 $\text{dim } ((\text{ideal}(a,b,c)^{\{d-1\}} \cap \text{ideal}(a,b)^{\{d-m\}}) H^0(X, L') + (\text{ideal}(a,b)^{\{2d\}})_{d-1+L \cdot L'})$ 
-  $\text{dim } (\text{ideal}(a,b)^{\{2d\}})_{d-1+L \cdot L'}$ . Thus  $\text{dim Im } \mu_{L+6E, 6E} = 174 - 95 = 79$ ,

```

computed with Macaulay, so $\dim \text{cok } \mu_{[L+6E, 6E]} = h^0(6E, 10) - 79 = 81 - 79 = 2$.
 In this case we know that the bound $\dim \text{cok } \mu_{[L+6E, 6E]} \leq \binom{m-a_E}{2} + \binom{m-b_E}{2} = \binom{6-4}{2} + \binom{6-4}{2} = 2$
 is in fact an equality. In other cases, where we only conjecture equality,
 we can use Macaulay to carry out the necessary computations, as shown below.
 A direct approach, where we find the graded Betti number of $I(Z)$ for
 $Z=Z(\{50,50,38,38,26,26,22,18,14,14\})$ in degree 104 requires working
 with homogeneous polynomials of degree 104 in three variables, whereas
 the approach using mE here involves homogeneous polynomials of degree at most 20,
 and the corresponding ideals are simpler (mostly being monomial ideals).

```
i18 : fundombothfully({8,{4,4,3,3,2,2,1,1}},{1,{0,0,0,0,0,0,0,0}})
o18 = {{0, {0, 0, 0, 0, 0, 0, 0, 0, -1}}, {13, {5, 5, 4, 4, 3, 2, 2, 1, 8}}
o18 : List
```

```
i19 : time II=mixer({8, 5, 5, 4, 4, 3, 2, 2, 1});
total: 1 12 11
0: 1 . .
1: . . .
2: . . .
3: . . .
4: . . .
5: . . .
6: . . .
7: . . .
8: . . .
9: . . .
10: . . .
11: . . .
12: . 3 .
13: . 9 11
-- used 0.57 seconds
```

```
o19 : Ideal of S
i20 : I=ideal(II_0,II_1,II_2);
o20 : Ideal of S
i21 : J=intersect((ideal(a,b,c))^7,(ideal(a,b))^2);
o21 : Ideal of S
i22 : K=I*J+(ideal(a,b))^16;
o22 : Ideal of S
i23 : time hilbertFunction(20,K)
-- used 1.99 seconds
o23 = 174
i24 : L=(ideal(a,b))^16;
o24 : Ideal of S
i25 : time hilbertFunction(20,L)
-- used 0.05 seconds
o25 = 95
```

```
XXXXXXXXXXXXXXXXXXXXXXXXXXXX
XXXXXXXXXXXXXXXXXXXXXXXXXXXX
```