

Directional Slope

The equation of a linear function $f(x, y)$ can be written as

$$f(x, y) = c + mx + ny,$$

where c , m , and n are constants.

Significance of m and n Consider two points, (x, y) and $(x + \Delta x, y + \Delta y)$. Let $\Delta f = f(x + \Delta x, y + \Delta y) - f(x, y)$. We may then calculate the change in f as


$$\Delta f = [c + m(x + \Delta x) + n(y + \Delta y)] - [c + mx + ny] = m\Delta x + n\Delta y.$$

This formula shows us the significance of the parameters m and n . Suppose we consider two points having the same y coordinate. Then $\Delta y = 0$. We can rearrange the last formula to get

$$m = \frac{\Delta f}{\Delta x}.$$

Thus, m is the slope of the function in the x -direction. Similarly, n is the slope of the function in the y -direction. Another way to state this is that m is the slope of the function in the direction given by the unit vector \vec{i} and n is the slope of the function in the direction given by the unit vector \vec{j} .

Slope in the direction of a unit vector \vec{u} Consider a specific example. We are interested in knowing the slope at the point $(2, 1)$ in the direction of the point $(6, 4)$.



Note that we have two different labels on the ray that begins at $(2, 1)$ and passes through the point $(6, 4)$. There is a coordinate label s and a unit vector label \vec{u} . This is analogous to the standard coordinate plane, in which the axes can be labeled with the coordinates x and y and also the unit vectors \vec{i} and \vec{j} .

The slope of f in the horizontal direction can be thought of as the rate of change of f with respect to x . Similarly, the slope of f along the path from $(2, 1)$ to $(6, 4)$ can be thought of as the rate of change of f with respect to s , where s is a coordinate that goes in the given direction. The slope in the s direction is $\Delta f / \Delta s$. Here, the change in f is $m\Delta x + n\Delta y = 4m + 3n$. The change in s is the distance between the points, or 5. So the slope is

$$\text{slope} = \frac{\Delta f}{\Delta s} = \frac{4m + 3n}{5} = 0.8m + 0.6n.$$

The slope of f in the horizontal direction can also be thought of as the slope of f in the \vec{i} direction. Similarly, the slope of f along the path from $(2, 1)$ to $(6, 4)$ can be thought of as the slope of f in the \vec{u} direction. The unit vector \vec{u} is obtained by dividing the displacement vector $4\vec{i} + 3\vec{j}$ by its length 5. Thus,

$$\vec{u} = 0.8\vec{i} + 0.6\vec{j}.$$

Now we define the **gradient vector**, which we will call g for now, as

$$\vec{g} = m\vec{i} + n\vec{j}.$$

The components of the gradient vector are the slopes in the corresponding directions. Is there a way to calculate the slope of f in the \vec{u} direction using the gradient and direction vectors? Clearly the answer is that the slope is the dot product of the two vectors:

$$\text{slope} = \vec{g} \cdot \vec{u}.$$

Summary

There are two ways to think of the slope of a linear function in a given direction.

- We can use a symbol, such as s , to represent distance in the given direction. Then we compute the slope in the given direction as the change in f divided by the change in s .

$$\text{slope} = \frac{\Delta f}{\Delta s}.$$

- We can describe the direction using the unit vector \vec{u} that points in the given direction. We can define the gradient vector to be the vector in which each component is the slope in the corresponding direction; thus, the \vec{i} component is the slope in the \vec{i} direction and so on. Then we compute the slope in the given direction as the dot product of the gradient and direction vectors.

$$\text{slope} = \vec{g} \cdot \vec{u}.$$

In 1-variable calculus, we use the concept of the slope of a line to define the derivative, which represents the slope of a curve. Similarly, in multi-variable calculus, we will use the concept of the slope of a plane to define the directional derivative, which represents the slope of a surface. The material here on slope of a linear function thus provides the precalculus foundation for the derivative in multi-dimensional calculus.

Exercises

1. Let $f = 2 + 3x + 2y$. Use both definitions of the slope to calculate the slope of f in the direction of the vector $2\vec{i} - \vec{j}$.
2. Let $f = 5 - x + 2y$. Use both definitions of the slope to calculate the slope of f in the direction of the vector $\vec{i} + 3\vec{j}$.
3. Let $f = 3 - 2x - 4y$. Use both definitions of the slope to calculate the slope of f in the direction of the vector $-2\vec{i} - 3\vec{j}$.
4. Let f be defined by the contour diagram in Figure 11.73. Determine the slope of f in the direction indicated by the ray from the point $(-1,-1)$ through the point $(1,2)$.