

## Math 107H Exam 1 Practice Problems

**Show all work.** How you get your answer is just as important, if not more important, than the answer itself.

Find each of the following integrals.

Note that “  $\int_3^x f(t) dt + C$  ” is not a sufficient computation of an antiderivative!

Some formulas of potential use can be found at the bottom of the last page of the exam.

1. (10 pts.)  $\int (x+2)^{3/2} dx$

2. (15 pts.)  $\int_0^{\pi/2} \sin^3 x dx$

3. (10 pts.)  $\int \frac{x^2 + x - 3}{x^{1/2}} dx$

4. (15 pts.)  $\int_0^1 e^{\sqrt{x}} dx$

5. (15 pts.)  $\int \frac{dx}{(x+1)^2(x+4)}$

6. (15 pts.)  $\int e^{-x} \sin(3x) dx$

7. (20 pts.)  $\int (x^2 + 1)^{3/2} dx = \int (\sqrt{x^2 + 1})^3 dx$

**Some formulas:**

$$\int \sec^n x dx = \frac{1}{n-1} \sec^{n-2} x \tan x + \frac{n-2}{n-1} \int \sec^{n-2} x dx$$

$$c^2 \int \frac{dy}{(y^2 + c^2)^k} = \frac{1}{(2k-2)} \cdot \frac{y}{(y^2 + c^2)^{k-1}} + \frac{(2k-3)}{(2k-2)} \int \frac{dy}{(y^2 + c^2)^{k-1}}$$

1. (10 pts. each) Find the following integrals:

(a):  $\int_1^4 x^2 \ln x dx$

(b):  $\int \sin^2 x \cos^3 x dx$

2. (10 pts. each) When you apply the appropriate trigonometric substitutions, what do the following integrals become?

(a):  $\int \frac{\sqrt{4-x^2}}{x^2} dx$

(b):  $\int \frac{x^2}{\sqrt{4x^2+9}} dx$

In problems 1 through 4, find each of the following integrals.

1. (10 pts.)  $\int_2^3 (3x-2)^{\frac{5}{4}} dx$

2. (15 pts.)  $\int x^{\frac{1}{3}} \ln x dx$

3. (10 pts.)  $\int_0^{\pi/2} \cos^3 x dx$

4. (15 pts.)  $\int \frac{dx}{1 + \sqrt{x}}$

6. (15 pts.) Use the method of partial fractions to express the function  $f(x) = \frac{1}{(x+2)(x+5)}$  as a sum of more elementary rational functions. Then **with no more work** use this answer to express  $g(x) = \frac{1}{(x^2+2)(x^2+5)}$  as a sum of elementary rational functions, and find the integral

$$\int \frac{1}{(x^2+2)(x^2+5)} dx .$$

6. (15 pts.) Recall that if a function  $f$  has second derivative satisfying  $|f''(x)| \leq M$  for every  $x$  in the interval  $[a, b]$ , then the error  $E_n$  in approximating the integral  $\int_a^b f(x) dx$  using the trapezoidal rule using  $n$  equal subintervals is at most

$$M \frac{(b-a)^3}{12n^2}$$

Based on this, how many subintervals should we divide the interval  $[2, 5]$  into in order to be sure to approximate the integral  $\int_2^5 x \ln x dx$  with an error of less than  $\frac{1}{100}$  ?