

We have introduced two homologies; simplicial, H_*^Δ , which is computationally straightforward, and singular, H_* , whose formal properties we have explored. For Δ -complexes, these homology groups are the same, $H_n^\Delta(X) \cong H_n(X)$ for every X . In fact, the isomorphism is induced by the inclusion $C_n^\Delta(X) \subseteq C_n(X)$. Note that most of the edifice we built for singular homology holds for simplicial homology, including relative homology (for a sub- Δ -complex A of X), and a SES of chain groups, giving a LES sequence for the pair, $\cdots \rightarrow H_n^\Delta(A) \rightarrow H_n^\Delta(X) \rightarrow H_n^\Delta(X, A) \rightarrow H_{n-1}^\Delta(A) \rightarrow \cdots$.

We proceed first by showing that the inclusion induces an isomorphism on k -skeleta, $H_n^\Delta(X^{(k)}) \cong H_n(X^{(k)})$, and this goes by induction on k using the

$$H_{n+1}^\Delta(X^{(k)}, X^{(k-1)}) \rightarrow H_n^\Delta(X^{(k-1)}) \rightarrow H_n^\Delta(X^{(k)}) \rightarrow H_n^\Delta(X^{(k)}, X^{(k-1)}) \rightarrow H_{n-1}^\Delta(X^{(k-1)})$$

Five Lemma applied to the diagram

$$\begin{array}{ccccccccc} & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H_{n+1}^\Delta(X^{(k)}, X^{(k-1)}) & \rightarrow & H_n^\Delta(X^{(k-1)}) & \rightarrow & H_n^\Delta(X^{(k)}) & \rightarrow & H_n^\Delta(X^{(k)}, X^{(k-1)}) & \rightarrow & H_{n-1}^\Delta(X^{(k-1)}) \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H_{n+1}(X^{(k)}, X^{(k-1)}) & \rightarrow & H_n(X^{(k-1)}) & \rightarrow & H_n(X^{(k)}) & \rightarrow & H_n(X^{(k)}, X^{(k-1)}) & \rightarrow & H_{n-1}(X^{(k-1)}) \end{array}$$

The second and fifth vertical arrows are, by an inductive hypothesis, isos. The first and fourth vertical arrows are isos because, essentially, we can, in each case, identify these groups. $H_n(X^{(k)}, X^{(k-1)}) \cong H_n(X^{(k)}/X^{(k-1)}) \cong \tilde{H}_n(\vee S^k)$ are either 0 (for $n \neq k$) or $\oplus \mathbb{Z}$ (for $n = k$), one summand for each n -simplex in X . But the same is true for $H_n^\Delta(X^{(k)}, X^{(k-1)})$; and for $n = k$ the generators are precisely the n -simplices of X . The inclusion-induced map takes generators to generators, so is an iso. So by the Five Lemma, the middle column is an iso, completing our inductive proof.

Returning to $H_n^\Delta(X) \xrightarrow{I_*} H_n(X)$, we wish now to show that this map is an isomorphism. Any $[z] \in H_n(X)$ is represented by a cycle $z = \sum a_i \sigma_i$ for $\sigma_i : \Delta^n \rightarrow X$. But each $\sigma_i(\Delta^n)$ is a compact subset of X , and so meets only finitely-many cells of X . This is true for every singular simplex, and so there is a k for which all of the simplices map into $X^{(k)}$, and so we may treat $z \in C_n(X^{(k)})$. Thought of in this way, it is still a cycle, and so $[z] \in H_n(X^{(k)}) \cong H_n^\Delta(X^{(k)})$ so there is a $z' \in C_n^\Delta(X^{(k)})$ and a $w \in C_{n+1}(X^{(k)})$ with $i_{\#} z' - z = \partial w$. But thinking of $z' \in C_n^\Delta(X)$ and $w \in C_{n+1}(X)$, we have the same equality, so $[z'] \in H_n^\Delta(X)$ and $i_*[z'] = [z]$. So i_* is surjective. If $i_*([z]) = 0$, then the cycle $z = \sum a_i \sigma_i$ is a sum of characteristic maps of n -simplices of X , and so can be thought of as an element of $C_n^\Delta(X^{(n)})$. Being 0 in $H_n(X)$, $z = \partial w$ for some $w \in C_{n+1}(X)$. But as before, $w \in C_n(X^{(r)})$ for some r , and so thought of as an element of the image of the isomorphism $i_* : H_n^\Delta(X^{(r)}) \rightarrow H_n(X^{(r)})$, $i_*([z]) = 0$, so $[z] = 0$. So $z = \partial u$ for some $u \in C_{n+1}^\Delta(X^{(r)}) \subseteq C_{n+1}^\Delta(X)$. So $[z] = 0$ in $H_n^\Delta(X)$. Consequently, simplicial and singular homology groups are isomorphic.

This implies the **topological invariance of the Euler characteristic of a space X** . If X is a Δ -complex made up of a finite number of simplices, then we can count the number m_i of i -simplices in the Δ -complex structure of X . The Euler characteristic of X is then defined to be the alternating sum $\chi(X) = \sum_{i=0}^\infty (-1)^i m_i$. Now, as a topological space, X can be given many different Δ -complex structures, and $\chi(X)$ is a priori a number which depends on the structure, not just on X . But once we note that m_i = the rank of the (simplicial) chain group $C_i^\Delta(X)$ (there is one generator for each i -simplex), we find that $\chi(X) = \sum_{i=0}^N (-1)^i \text{rank}(C_i(X))$, and then the following result from homological algebra establishes the topological invariance of this number:

Proposition: If $\cdots 0 \rightarrow C_n \rightarrow \cdots \rightarrow C_1 \rightarrow C_0 \rightarrow 0$ in a chain complex with all finite rank, then $\sum_{i=0}^n (-1)^i \text{rank}(C_i) = \sum_{i=0}^n (-1)^i \text{rank}(H_i(C))$.

The proof follows from the facts that since $H_i(C) = \ker \partial_i / \text{im} \partial_{i+1}$, $z_i = \text{rank}(\ker \partial_i) = \text{rank}(H_i(C)) + \text{rank}(\text{im} \partial_{i+1}) = h_i + b_{i+1}$, so $h_i = z_i - b_{i+1}$, and that (by Noether) $\text{im}(\partial_i) \cong C_i / \ker(\partial_i)$, so $c_i = \text{rank}(C_i) = z_i + b_i$. We therefore have

$$\sum_{i=0}^n (-1)^i \text{rank}(H_i(C)) = \sum_{i=0}^n (-1)^i h_i = \sum_{i=0}^n (-1)^i (z_i - b_{i+1}) = \sum_{i=0}^n (-1)^i z_i - \sum_{i=0}^n (-1)^i b_{i+1} = \sum_{i=0}^n (-1)^i z_i + \sum_{i=0}^n (-1)^i b_i = \sum_{i=0}^n (-1)^i (z_i + b_i) = \sum_{i=0}^n (-1)^i \text{rank}(C_i) \quad \text{as desired.}$$

Consequently, $\chi(X) = \sum_{i=0}^N (-1)^i \text{rank}(C_i^\Delta(X)) = \sum_{i=0}^N (-1)^i \text{rank}(H_i^\Delta(X)) = \sum_{i=0}^N (-1)^i \text{rank}(H_i(X))$, which is an invariant of X , since the singular homology groups are!

The fact that this number has two different interpretations leads to some non-trivial results. First, it tells us that the Euler characteristic calculation is independent of how we express a space X as a Δ -complex. χ is also actually invariant under homotopy equivalence, since the homology groups are; so **homotopy equivalent spaces have the same Euler χ** . Consequently, all contractible spaces, for example, must have Euler characteristic = 1.

Next, by the lifting criterion, if $p : \tilde{X} \rightarrow X$ is a k -fold covering space of a Δ -complex X , then \tilde{X} can be given a Δ -complex structure with k times as many i -simplices as X , for every i (lift the characteristic maps of the simplices of X ...). So $\chi(\tilde{X}) = k \cdot \chi(X)$. This gives a necessary condition for one space to be a covering of another; it's Euler χ must be a multiple of the other. For example, from our homology calculations, it follows that **for a closed orientable surface F_g of genus g , $\chi(F_g) = 2 - 2g$. So a k -fold covering of F_g will have Euler χ equal to $k(2 - 2g) = 2k - 2kg = 2 - 2(kg - k + 1)$, and so is a surface of genus $kg - k + 1$** . [The converse, that a surface with this genus k -fold covers F_g , can be established by building the coverings directly.] Consequently, F_5 is a 2-fold covering of F_3 , so there is a subgroup of index 2 of $\pi_1(F_3)$ isomorphic to $\pi_1(F_5)$, but F_6 is not a finite-sheeted cover of F_3 , because $-4 \nmid -10$. [It is also not an infinite-sheeted covering, because their total spaces are non-compact...] Consequently, $\pi_1(F_6)$ is not isomorphic to a subgroup of $\pi_1(F_3)$.