

Name: _____

Score: _____

Instructions: Show your work in the spaces provided below for full credit. Use the reverse side for additional space, *but clearly so indicate*. You must clearly identify answers and show supporting work to receive any credit. Exact answers (e.g., π) are preferred to inexact (e.g., 3.14). Make all obvious simplifications, e.g., 0 rather than $\sin \pi$. Point values of problems are given in parentheses. Point values of problems are given in parentheses. Notes or text in *any* form not allowed. The only electronic equipment allowed is a calculator.

(14) **1.** Evaluate the integral $I = \int_0^1 \int_0^{3-3x} \int_0^{3-3x-y} dz dy dx$.

SOLUTION. (Exercise 13.5.10) We have, with substitution $u = (3 - 3x)$, $du = -3dx$, $dx/2 = -du/6$, $u(0) = 3$, $u(1) = 0$, (or just observing $\int \frac{(3-3x)^2}{2} dx = \int \frac{9(x-1)^2}{2} dx = 3 \frac{(x-1)^3}{2}$)

$$\begin{aligned} I &= \int_0^1 \int_0^{3-3x} z \Big|_{z=0}^{3-3x-y} dy dx = \int_0^1 \int_0^{3-3x} (3-3x-y) dy dx \\ &= \int_0^1 \left((3-3x)y - \frac{y^2}{2} \right) \Big|_{y=0}^{3-3x} dx = \int_0^1 \left((3-3x)^2 - \frac{(3-3x)^2}{2} \right) dx \\ &= \int_0^1 \frac{(3-3x)^2}{2} dx = - \int_3^0 \frac{u^2}{6} du = \frac{u^3}{18} \Big|_0^3 = \frac{3}{2}. \end{aligned}$$

(18) **2.** Sketch the region D over which the iterated integral I below is calculated. Then express the integral in the order $dy dz dx$ and write a formula for the average value of $f(x, y, z)$ over D in terms of iterated integrals. Do NOT evaluate any integrals.

$$I = \int_0^4 \int_0^1 \int_{2y}^2 f(x, y, z) dx dy dz.$$

SOLUTION. (Exercise 13.5.41) Region Q is sketched to the right. From it we see that


$$I = \int_0^2 \int_0^4 \int_0^{x/2} f(x, y, z) dy dz dx.$$

Thus, we have that the area of the region is

$$A = \int_0^2 \int_0^4 \int_0^{x/2} dy dz dx$$

and that the average value of $f(x, y, z)$ over Q is

$$\bar{f}_Q = I/A.$$



(18) **3.** Sketch the region R in the plane bounded by the curves $y = 0$, $y^2 = 2x$, and $x + y = 4$ and use iterated integrals to write formulas for the area and the first moment of R about the x -axis (assume density $\delta = y^2$) in terms of iterated integrals. Do NOT evaluate any integrals. (Arithmetic check: second and third curves intersect at $(2, 2)$.)


SOLUTION.

(Exercise 13.6.3) The quadratic and straight line intersect where $y^2 = 2(4 - y)$, i.e., where $0 = y^2 + 2y - 8 = (y + 4)(y - 2)$, and at $y = 2$ we have $x = 2^2/2 = 2$. Region R is sketched in the graph to the right. We have that

$$\text{Area}(R) = \iint_R dA = \int_0^2 \int_{y^2/2}^{4-y} dx dy$$

(or $\int_0^2 \int_0^{\sqrt{2x}} dy dx + \int_2^4 \int_0^{4-x} dy dx$) and that the first moment of R about the x -axis, assuming that $\delta(x, y) = y^2$ is given by

$$M_x = \iint_R y \delta dA = \int_0^2 \int_{y^2/2}^{4-y} y^3 dx dy.$$



(20) **4.** A solid D is bounded by the surfaces $z = 1$ and $z = \sqrt{x^2 + y^2}$. Sketch it and express the integral $\iiint_D f(x, y, z) dV$ as an iterated integral in both cylindrical and spherical coordinates. Use one of these to express the moment of inertia I_z about the z -axis of a solid occupying D with density function $\delta = z$ as an iterated integral. Do NOT evaluate it.

SOLUTION.

(Exercise 13.7.77) The region is sketched in the figure to the right. The plane $z = 1$ and cone $z = \sqrt{x^2 + y^2}$ intersect in the circle $1 = x^2 + y^2$, which is the shadow R of the region D in the xy -plane. Also, the cone makes an angle of $\pi/4$ with the vertical and the plane $z = 1$ gives $\rho \cos \phi = 1$, i.e., $\rho = 1/\cos \phi = \sec \phi$. Hence, the integral in spherical coordinates is


Also in cylindrical coordinates

$$\begin{aligned} I_z &= \iiint_D (x^2 + y^2) \delta dV \\ &= \int_0^{2\pi} \int_0^1 \int_r^1 r^3 z dz dr d\theta. \end{aligned}$$

$$\int_0^{2\pi} \int_0^{\pi/4} \int_0^{\sec \phi} f(\rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi) \rho^2 \sin \phi d\rho d\phi d\theta.$$

In cylindrical coordinates the integral is

$$\int_0^{2\pi} \int_0^1 \int_r^1 f(r \cos \theta, r \sin \theta, z) dz r dr d\theta.$$



(16) **5.** Evaluate the line integral $\int_C 1 ds$ where C has position vector $\mathbf{r}(t) = \langle \cos t, \sin t, \frac{2}{3}t^{3/2} \rangle$, $0 \leq t \leq 1$.

SOLUTION. (Exercise 14.1.30) The position vector $\mathbf{r}(t)$ gives us a parametrization of C , namely

$$\begin{aligned}x &= \cos t \\y &= \sin t \\z &= \frac{2}{3}t^{3/2},\end{aligned}$$

from which we obtain differential formulas

$$\begin{aligned}dx &= -\sin t dt \\dy &= \cos t dt \\dz &= \frac{2}{3}t^{1/2} dt = t^{1/2} dt,\end{aligned}$$

from which it follows that $ds = \sqrt{dx^2 + dy^2 + dz^2} = \sqrt{(-\sin t)^2 + (\cos t)^2 + (t^{1/2})^2} = \sqrt{1+t} dt$. (Or use $ds = |d\mathbf{r}| = \left| \frac{d\mathbf{r}}{dt} \right| dt$.) Thus, with substitution $u = 1+t$, $du = dt$, $u(0) = 1$, $u(1) = 2$ (or just observing $\int \sqrt{1+t} dt = \frac{2}{3}(1+t)^{3/2}$)

$$\int_C 1 ds = \int_0^1 \sqrt{1+t} dt = \int_1^2 u^{1/2} du = \frac{2}{3}u^{3/2} \Big|_{u=1}^2 = \frac{2}{3}(2\sqrt{2} - 1) = \frac{4}{3}\sqrt{2} - \frac{2}{3}.$$

(14) **6.** Let C be the curve $y = x^2$ traversed from $(0,0)$ to $(1,1)$, and $\mathbf{F} = \langle x, y \rangle$ a vector field. Express the flow (i.e., work) of \mathbf{F} along C as a line integral and evaluate it.

SOLUTION. (Exercise 14.2.17) We have $\langle M, N \rangle = \mathbf{F}(x, y) = \langle x, y \rangle$ and that the flow of \mathbf{F} along C is

$$W = \int_C \mathbf{F} \cdot \mathbf{T} ds = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_C M dx + N dy = \int_C x dx + y dy.$$

Now parametrize C with position vector $\mathbf{r}(t) = \langle t, t^2 \rangle$ (or $\mathbf{r}(t) = \langle x, x^2 \rangle$) and get

$$\begin{aligned}x &= t \\y &= t^2, 0 \leq t \leq 1\end{aligned}$$

so that $dx = dt$, $dy = 2t dt$, and obtain that

$$W = \int_0^1 (t dt + t^2 2t dt) = \int_0^1 (t + 2t^3) dt = \left(\frac{t^2}{2} + \frac{2}{4}t^4 \right) \Big|_{t=0}^1 = \frac{1}{2} + \frac{1}{2} - 0 = 1.$$