

COFINITE MODULES AND LOCAL COHOMOLOGY

DONATELLA DELFINO AND THOMAS MARLEY¹

ABSTRACT. We show that if M is a finitely generated module over a commutative Noetherian local ring R and I is a dimension one ideal of R (i.e., $\dim R/I = 1$), then the local cohomology modules $H_I^i(M)$ are I -cofinite; that is, $\text{Ext}_R^j(R/I, H_I^i(M))$ is finitely generated for all i, j . We also show that if R is a complete local ring and P is a dimension one prime ideal of R , then the set of P -cofinite modules form an abelian subcategory of the category of all R -modules. Finally, we prove that if M is an n -dimensional finitely generated module over a Noetherian local ring R and I is any ideal of R , then $H_I^n(M)$ is I -cofinite.

Let R be a commutative Noetherian local ring with maximal ideal m and let I be an ideal of R . An R -module N is said to be I -cofinite if $\text{Supp } N \subseteq V(I)$ and $\text{Ext}_R^i(R/I, N)$ is finitely generated for all $i \geq 0$. Using Matlis duality one can show that a module is m -cofinite if and only if it is Artinian. As a consequence, the local cohomology modules $H_m^i(M)$ are m -cofinite for any finitely generated R -module M . In [6], Hartshorne posed the question of whether this statement still holds when m is replaced by an arbitrary ideal I ; i.e., is $H_I^i(M)$ I -cofinite for all i ? In general, the answer is no, even if R is a regular local ring. Let $R = k[[x, y, u, v]]$ be the formal power series ring in four variables over a field k , m the maximal ideal of R , $P = (x, u)R$ and $M = R/(xy - uv)$. Hartshorne showed that $\text{Hom}_R(R/m, H_P^2(M))$ is not finitely generated, and hence $\text{Hom}_R(R/P, H_P^2(M))$ cannot be finitely generated. In the positive direction, Hartshorne proved that if R is a complete regular local ring, P a dimension one prime ideal of R , and M a finitely generated R -module, then $H_P^i(M)$ is P -cofinite for all i . In 1991, Huneke and Koh proved that if R is a complete local Gorenstein domain, I a dimension one ideal of R , and M a finitely generated R -module, then $H_I^i(M)$ is I -cofinite for all i ([7, Theorem 4.1]). Recently, Delfino proved that the Gorenstein hypothesis in the Huneke-Koh theorem may be weakened to include all complete local domains R which satisfy one of the following conditions: (1) R contains a field; (2) if q is a uniformizing parameter for a coefficient ring for R then either $q \in \sqrt{I}$ or q is not in any prime minimal over I ; or (3) R is Cohen-Macaulay ([3, Theorem 3] and [4, Theorem 2.21]). In this paper, we eliminate the complete domain hypothesis entirely by proving the following:

Theorem 1. *Let R be a Noetherian local ring, I a dimension one ideal of R , and M a finitely generated R -module. Then $H_I^i(M)$ is I -cofinite for all i .*

We prove this by establishing a change of ring principle for cofiniteness (Proposition 2) and then applying it to the Huneke-Koh result. Using this change of ring principle,

¹Both authors were partially supported by the National Science Foundation.

we are also able to generalize Hartshorne's theorem that over a regular local ring, the P -cofinite modules (P a dimension one prime) form an abelian subcategory of the category of R -modules (Theorem 2).

We also prove a cofiniteness result about $H_I^n(M)$, where M is a finitely generated R -module and $n = \dim M$. In [12], Sharp proved that if R is a Noetherian local ring of dimension d and I is any ideal of R , then $H_I^d(R)$ is Artinian. From this it follows easily that if M is a finitely generated R -module of dimension n then $H_I^n(M)$ is Artinian. (See also [10, Theorem 2.2].) Thus, $H_I^n(M)$ is m -cofinite. We prove that $H_I^n(M)$ is in fact I -cofinite (Theorem 3).

We begin the proof of Theorem 1 by proving the following generalization of [7, Lemma 4.2] and [3, Lemma 2]:

Proposition 1. *Let R be a Noetherian ring, M a finitely generated R -module and N an arbitrary R -module. Suppose that for some $p \geq 0$, $\text{Ext}_R^i(M, N)$ is finitely generated for all $i \leq p$. Then for any finitely generated R -module L with $\text{Supp } L \subseteq \text{Supp } M$, $\text{Ext}_R^i(L, N)$ is finitely generated for all $i \leq p$.*

proof. Using induction on p , we may assume that $\text{Ext}_R^i(L, N)$ is finitely generated for all $i < p$ and all finitely generated modules L with $\text{Supp } L \subseteq \text{Supp } M$. (This is satisfied vacuously if $p = 0$.) By Gruson's Theorem ([13, Theorem 4.1]), given any finitely generated R -module L with $\text{Supp } L \subseteq \text{Supp } M$ there exists a finite filtration

$$0 = L_0 \subset L_1 \subset \cdots \subset L_n = L$$

such that the factors L_i/L_{i-1} are homomorphic images of a direct sum of finitely many copies of M . By using short exact sequences and induction on n , it suffices to prove the case when $n = 1$. Thus, we have an exact sequence of the form

$$0 \longrightarrow K \longrightarrow M^n \longrightarrow L \longrightarrow 0$$

for some positive integer n and some finitely generated module K . This gives the long exact sequence

$$\cdots \longrightarrow \text{Ext}_R^{p-1}(K, N) \longrightarrow \text{Ext}_R^p(L, N) \longrightarrow \text{Ext}_R^p(M^n, N) \longrightarrow \cdots$$

Since $\text{Supp } K \subseteq \text{Supp } M$ we have that $\text{Ext}_R^{p-1}(K, N)$ is finitely generated (by the induction on p). As $\text{Ext}_R^p(M^n, N) \cong \text{Ext}_R^p(M, N)^n$ is finitely generated, the result follows.

As a consequence, we have the following:

Corollary 1. *Let R be a Noetherian ring, I an ideal of R and N an R -module. The following are equivalent:*

- (a) $\text{Ext}_R^i(R/I, N)$ is finitely generated for all $i \geq 0$,
- (b) $\text{Ext}_R^i(R/J, N)$ is finitely generated for all $i \geq 0$ and ideals $J \supseteq I$,
- (c) $\text{Ext}_R^i(R/P, N)$ is finitely generated for all $i \geq 0$ and all primes P minimal over I .

proof. We show that (c) implies (a). Let P_1, \dots, P_n be the minimal primes of I and $M = R/P_1 \oplus \cdots \oplus R/P_n$. Then $\text{Ext}_R^i(M, N)$ is finitely generated for all i . As $\text{Supp } R/I = \text{Supp } M$, $\text{Ext}_R^i(R/I, N)$ is finitely generated for all i by the proposition.

The next result concerns spectral sequences, for which we use the notation from chapter 5 of [14]. The essential idea for this lemma can be found in the proof of [3, Theorem 3].

Lemma 1. *Let R be a Noetherian ring and $\{E_r^{pq}\}$ a first quadrant cohomology spectral sequence (starting with E_a , for some $a \geq 1$) converging to H^* in the category of R -modules. For a fixed integer n , suppose H^n is finitely generated and $E_a^{p,q}$ is finitely generated for all $p < n$ and $q \geq 0$. Then $E_a^{n,0}$ is finitely generated.*

proof. If $n = 0$ then $E_a^{00} = H^0$ is finitely generated. Suppose $n > 0$. First note that E_r^{pq} is finitely generated for any $p < n$, $q \geq 0$, and $r \geq 1$, since E_r^{pq} is a subquotient of E_a^{pq} . Also, as $E_\infty^{n,0}$ is isomorphic to a submodule of H^n , $E_\infty^{n,0}$ is finitely generated. Now since $\{E_r^{pq}\}$ is a first quadrant spectral sequence (in particular, since there are no nonzero terms below the p -axis), there is an exact sequence

$$E_r^{n-r,r-1} \longrightarrow E_r^{n,0} \longrightarrow E_{r+1}^{n,0} \longrightarrow 0$$

for all $r \geq a$. As $E_r^{n,0} = E_\infty^{n,0}$ for sufficiently large r (and thus is finitely generated), we can work backwards to see that $E_r^{n,0}$ is finitely generated for all $r \geq a$.

We now prove the change of ring principle for cofiniteness:

Proposition 2. *Let R be a Noetherian ring and S a module finite R -algebra. Let I be an ideal of R and M an S -module. Then M is I -cofinite (as an R -module) if and only if M is IS -cofinite (as an S -module).*

proof. First note that $\text{Supp}_R M \subseteq V(I)$ if and only if $\text{Supp}_S M \subseteq V(IS)$. Now consider the Grothendieck spectral sequence (see [11, Theorem 11.65], for example)

$$E_2^{pq} = \text{Ext}_S^p(\text{Tor}_q^R(S, R/I), M) \implies \text{Ext}_R^{p+q}(R/I, M).$$

Suppose first that M is IS -cofinite. Then $E_2^{p,0} = \text{Ext}_S^p(S/IS, M)$ is finitely generated for all p . Since $\text{Supp Tor}_q^R(S, R/I) \subseteq \text{Supp } S/IS$ for all q , E_2^{pq} is finitely generated for all p and q by Proposition 1. Since the spectral sequence is bounded, it follows that $\text{Ext}_R^n(R/I, M)$ is finitely generated for all n .

Conversely, suppose that M is I -cofinite. We use induction on n to show $E_2^{n,0} = \text{Ext}_S^n(S/IS, M)$ is finitely generated. Now $E_2^{00} = \text{Hom}_S(S/IS, M) \cong \text{Hom}_R(R/I, M)$ is finitely generated. Suppose that $n > 0$ and $E_2^{p,0}$ is finitely generated for all $p < n$. By Proposition 1, E_2^{pq} is finitely generated for all $p < n$ and $q \geq 0$. Since $H^n = \text{Ext}_R^n(R/I, M)$ is finitely generated, $E_2^{n,0}$ is finitely generated by Lemma 1.

As a final preparation for the proof of Theorem 1, we need the following fact:

Lemma 2. *Let (R, m) be a local ring and S the m -adic completion of R . Let I be an ideal of R and M an R -module. Then $H_I^i(M)$ is I -cofinite if and only if $H_{IS}^i(M \otimes_R S)$ is IS -cofinite.*

proof. Since $\text{Ext}_R^i(R/I, H_I^i(M)) \otimes_R S \cong \text{Ext}_S^i(S/IS, H_{IS}^i(M \otimes_R S))$, it is enough to see that an R -module N is finitely generated if and only if $N \otimes_R S$ is finitely generated as an S -module. If N is finitely generated, the implication is obvious. If $N \otimes_R S$ is finitely generated then, using the faithful flatness of S , one can see that any ascending chain of submodules of N must stabilize.

Theorem 1 now follows readily:

proof of Theorem 1. By Lemma 2 we may assume R is complete. Thus, R is the homomorphic image of a regular local ring T . Let J be a dimension one ideal of T such that $JR = I$. Then $H_J^j(M)$ is J -cofinite by [7, Theorem 4.1] for all j . By Proposition 2, $H_I^j(M) \cong H_J^j(M)$ is I -cofinite for all j .

If N is an R -module then the i th Bass number of N with respect to p is defined by $\mu_i(p, N) = \dim_{k(p)} \text{Ext}_{R_p}^i(k(p), N_p)$, where $k(p) = (R/p)_p$. If M is finitely generated and I is a zero-dimensional ideal then the Bass numbers of $H_I^i(M)$ are finite since $H_I^i(M)$ is Artinian. However, as Hartshorne's example shows, this does not hold for arbitrary ideals and modules, even over a complete regular local ring. In the special case that $M = R$, Huneke and Sharp proved that if R is a regular local ring of characteristic p and I is an ideal of R , then the Bass numbers of $H_I^i(R)$ are finite for all i ([8, Theorem 2.1]). Lyubeznik proved this same result in the case R is a regular local ring containing a field of characteristic 0 ([9, Corollary 3.6]). In [1], it is proved that if R is a complete local Gorenstein domain, I is a dimension one ideal and M is a Matlis reflexive R -module (i.e., $\text{Hom}_R(\text{Hom}_R(M, E), E) \cong M$ where $E = E_R(R/m)$), then the Bass numbers of $H_I^i(M)$ are finite. Using Theorem 1, we can prove the following:

Corollary 2. *Let R be a Noetherian ring, I a dimension one ideal of R , and M a finitely generated R -module. Then $\mu_i(p, H_I^j(M)) < \infty$ for all integers i, j and $p \in \text{Spec}(R)$.*

proof. If $p \not\supseteq I$, then $\mu_i(p, H_I^j(M)) = 0$. If $p \supseteq I$ we can localize and assume $p = m$. By Theorem 1, $\text{Ext}_R^i(R/I, H_I^j(M))$ is finitely generated for all i, j . Thus, $\text{Ext}_R^i(R/m, H_I^j(M))$ is finitely generated for all i, j by Corollary 1.

Another question Hartshorne addressed in [6] was the following: if R is a complete regular local ring and P is a prime ideal, do the P -cofinite modules form an abelian subcategory of the category of all R -modules? That is, if $f: A \rightarrow B$ is an R -module map of P -cofinite modules, are $\ker f$ and $\text{coker } f$ P -cofinite? Hartshorne gave the following counterexample: let $R = k[[x, y, u, v]]$, $P = (x, u)$ and $M = R/(xy - uv)$ as in the example mentioned above. Applying the functor $H_P^0(-)$ to the exact sequence

$$0 \rightarrow R \xrightarrow{xy-uv} R \rightarrow M \rightarrow 0$$

we get the exact sequence

$$\dots \rightarrow H_P^2(R) \xrightarrow{f} H_P^2(R) \rightarrow H_P^2(M) \rightarrow 0.$$

Since $H_P^j(R) = 0$ for all $j \neq 2$, one can show (using a collapsing spectral sequence) that $\text{Ext}_R^i(R/P, H_P^2(R)) \cong \text{Ext}_R^{i+2}(R/P, R)$ for all i . Thus, $H_P^2(R)$ is P -cofinite. However, as mentioned above, $\text{coker } f = H_P^2(M)$ is not P -cofinite. On the positive side, Hartshorne proved that if P is a dimension one prime ideal of a complete regular local ring then the answer to his question is yes. Using Proposition 2, we can extend this result to arbitrary complete local rings:

Theorem 2. *Let R be a complete local ring and P a dimension one prime ideal of R . Then the P -cofinite modules form an abelian subcategory of the category of all R -modules.*

proof. Let $f: M \rightarrow N$ be a map of P -cofinite modules. Since R is complete there exists a regular local ring T and a dimension one prime ideal Q of T such that R is a quotient of T and $QT = P$. Since M and N are Q -cofinite T -modules, $\ker f$ and $\operatorname{coker} f$ are Q -cofinite by Hartshorne's theorem ([6, Proposition 7.6]). Therefore, $\ker f$ and $\operatorname{coker} f$ are P -cofinite by Proposition 2.

We strongly believe Theorem 2 holds for dimension one ideals of an arbitrary local ring.

We now turn our attention to proving Theorem 3. The techniques are essentially those of Sharp ([12]) and Yassemi ([15]). Let (R, m) be a local ring, M an R -module, and $E = E_R(R/m)$ the injective hull of R/m . Following [15], we define a prime p to be a *coassociated prime* of M if p is an associated prime of $M^\vee = \operatorname{Hom}_R(M, E)$. We denote the set of coassociated primes of M by $\operatorname{Coass}_R M$ (or simply $\operatorname{Coass} M$ if there is no ambiguity about the underlying ring). Note that $\operatorname{Coass} M = \emptyset$ if and only if $M = 0$. We first make a couple of preliminary remarks:

Remark 1. ([15, Theorem 1.22]) *Let (R, m) be a Noetherian local ring, M a finitely generated R -module and N an arbitrary R -module. Then $\operatorname{Coass}(M \otimes_R N) = \operatorname{Supp} M \cap \operatorname{Coass} N$.*

proof. Note that $(M \otimes_R N)^\vee \cong \operatorname{Hom}_R(M, N^\vee)$. Therefore

$$\begin{aligned} \operatorname{Coass}(M \otimes_R N) &= \operatorname{Ass}(\operatorname{Hom}_R(M, N^\vee)) \\ &= \operatorname{Supp} M \cap \operatorname{Ass} N^\vee \quad (\text{e.g., [2, IV.1.4, Prop. 10]}) \\ &= \operatorname{Supp} M \cap \operatorname{Coass} N. \end{aligned}$$

Remark 2. *Let R be a local ring of dimension d , I an ideal of R and M an R -module. Then $H_I^d(M) \cong M \otimes_R H_I^d(R)$.*

proof. Since $H_I^d(-)$ is a right exact functor, this remark is an immediate consequence of Watts' Theorem [11, Theorem 3.33]. Here is a more direct proof: since R is local, there exist elements $\underline{x} = x_1, \dots, x_d \in I$ which generate I up to radical. Then $H_I^i(M) = H_{(\underline{x})}^i(M)$ for all i . Using the Čech complex to compute $H_{(\underline{x})}^d(R)$, we see there is an exact sequence

$$\bigoplus_i R_{x_1 \dots \hat{x}_i \dots x_d} \rightarrow R_{x_1 \dots x_d} \rightarrow H_{(\underline{x})}^d(R) \rightarrow 0.$$

Tensoring this sequence with M , we get the exact sequence

$$\bigoplus_i M_{x_1 \dots \hat{x}_i \dots x_d} \xrightarrow{f} M_{x_1 \dots x_d} \rightarrow M \otimes_R H_{(\underline{x})}^d(R) \rightarrow 0.$$

Since $\operatorname{coker} f = H_{(\underline{x})}^d(M)$, we see that $H_I^d(M) \cong M \otimes_R H_I^d(R)$.

The next result is essentially a module version of [12, Theorem 3.4] combined with [15, Theorem 1.16]. As in [12], we make repeated use of the Hartshorne-Lichtenbaum vanishing theorem (HLVT): if (R, m) is a complete local ring of dimension d and I is an ideal of R , then $H_I^d(R) \neq 0$ if and only if $I + p$ is m -primary for some prime ideal p such that $\dim R/p = d$ ([5, Theorem 3.1]).

Lemma 3. *Let (R, m) be a complete Noetherian local ring, I an ideal of R and M a finitely generated R -module of dimension n . Then*

$$\text{Coass } H_I^n(M) = \{p \in V(\text{Ann}_R M) \mid \dim R/p = n \text{ and } \sqrt{I+p} = m\}.$$

proof. Let $S = R/\text{Ann}_R M$ and $E_S = \text{Hom}_R(S, E)$ the injective hull of the residue field of S . Observe that

$$\begin{aligned} \text{Hom}_S(H_{IS}^n(M), E_S) &\cong \text{Hom}_R(H_I^n(M), E_S) \\ &\cong \text{Hom}_R(H_I^n(M) \otimes_R S, E) \\ &\cong \text{Hom}_R(H_I^n(M), E). \end{aligned}$$

Consequently, $\text{Coass}_R H_I^n(M) = \pi(\text{Coass}_S H_{IS}^n(M))$ where $\pi: \text{Spec } S \rightarrow \text{Spec } R$. Thus, we may assume that $\text{Ann}_R M = 0$ and $n = \dim R$. By Remarks 1 and 2, we have $\text{Coass } H_I^n(M) = \text{Coass}(M \otimes_R H_I^n(R)) = \text{Coass } H_I^n(R)$, so it is enough to prove the result in the case $M = R$. By HLVT, both sets are empty if $H_I^n(R) = 0$, so assume that $H_I^n(R) \neq 0$. Let $q \in \text{Coass } H_I^n(R)$. By the remark, $q \in \text{Coass}(R/q \otimes_R H_I^n(R))$. In particular, $R/q \otimes_R H_I^n(R) \cong H_I^n(R/q) \neq 0$. Thus, $n = \dim R/q$ and $I+q$ is m -primary (by HLVT). Now suppose $\dim R/q = n$ and $\sqrt{I+q} = m$. By reversing the above argument we get that $R/q \otimes_R H_I^n(R) \neq 0$. Let $p \in \text{Coass}(R/q \otimes_R H_I^n(R))$. By Remark 1, $p \supseteq q$ and $p \in \text{Coass } H_I^n(R)$. We've already shown that every coassociated prime of $H_I^n(R)$ is a minimal prime of R . Hence $p = q$ and $q \in \text{Coass } H_I^n(R)$, which completes the proof.

We now show that $H_I^{\dim M}(M)$ is I -cofinite:

Theorem 3. *Let (R, m) be Noetherian local ring, I an ideal of R and M a finitely generated R -module of dimension n . Then $H_I^n(M)$ is I -cofinite. In fact, $\text{Ext}_R^i(R/I, H_I^n(M))$ has finite length for all i .*

proof. By Lemma 2 we may assume that R is complete. Let $\text{Coass } H_I^n(M) = \{p_1, \dots, p_k\}$. Since $H_I^n(M)$ is Artinian (see [12, Theorem 3.3], $H_I^n(M)^\vee$ is finitely generated. Hence $\text{Supp } H_I^n(M)^\vee = V(p_1 \cap \dots \cap p_k)$. By Matlis duality, $\text{Ext}_R^i(R/I, H_I^n(M))$ has finite length if and only if $\text{Ext}_R^i(R/I, H_I^n(M))^\vee \cong \text{Tor}_i^R(R/I, H_I^n(M)^\vee)$ ([11, Theorem 11.57]) has finite length. Since $\text{Tor}_i^R(R/I, H_I^n(M)^\vee)$ is finitely generated, it is enough to show its support is contained in $\{m\}$. But

$$\begin{aligned} \text{Supp } \text{Tor}_i^R(R/I, H_I^n(M)^\vee) &\subseteq V(I) \cap \text{Supp } H_I^n(M)^\vee \\ &= V(I) \cap V(p_1 \cap \dots \cap p_k) \\ &= V(I + (p_1 \cap \dots \cap p_k)) \\ &= \{m\} \quad (\text{by Lemma 3}). \end{aligned}$$

REFERENCES

- [1] R. Belshoff, S. Slattery and C. Wickham, *The local cohomology modules of Matlis reflexive modules are almost cofinite*, Proc. Amer. Math. Soc. (to appear).

- [2] N. Bourbaki, *Algèbre Commutative*, Hermann, Paris, 1961.
- [3] D. Delfino, *On the cofiniteness of local cohomology modules*, Math. Proc. Camb. Phil. Soc. **115** (1994), 79–84.
- [4] D. Delfino, *Cofiniteness and vanishing of local cohomology modules, and colength of conductor ideals*, Ph.D. Dissertation, Purdue University (1994).
- [5] R. Hartshorne, *Cohomological dimension of algebraic varieties*, Annals of Math. **88** (1968), 403–450.
- [6] R. Hartshorne, *Affine duality and cofiniteness*, Inv. Math. **9** (1970), 145–164.
- [7] C. Huneke and J. Koh, *Cofiniteness and vanishing of local cohomology modules*, Math. Proc. Cambridge Phil. Soc. **110** (1991), 421–429.
- [8] C. Huneke and R. Sharp, *Bass numbers of local cohomology modules*, Trans. American Math. Soc. **339** (1993), 765–779.
- [9] G. Lyubeznik, *Finiteness properties of local cohomology modules (an application of D -modules to commutative algebra)*, Inv. Math. **113**, 41–55.
- [10] L. Melkersson, *Some applications of a criterion for artinianness of a module*, J. Pure Appl. Algebra **101 (3)** (1995), 291–303.
- [11] J. Rotman, *An Introduction to Homological Algebra*, Academic Press, Orlando, FL, 1979.
- [12] R. Sharp, *On the attached prime ideals of certain Artinian local cohomology modules*, Proc. Edinburgh Math. Soc. **24(2)** (1981).
- [13] W. Vasconcelos, *Divisor Theory in Module Categories*, North-Holland Publishing Company, Amsterdam, 1974.
- [14] C. Weibel, *An Introduction to Homological Algebra*, Cambridge University Press, Cambridge, 1994.
- [15] S. Yassemi, *Coassociated primes*, preprint.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI, 48109
E-mail address: delfino@math.lsa.umich.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEBRASKA, LINCOLN, NE 68588
E-mail address: tmarley@math.unl.edu