

Craig Huneke · Roger Wiegand

Correction to “Tensor Products of Modules and the Rigidity of Tor”, Math. Annalen, 299 (1994), 449–476.

Received: July 30, 2006

Abstract This note makes a correction to the paper “Tensor products of modules and the rigidity of Tor”, a correction which is needed due to an incorrect convention for the depth of the zero module.

Mathematics Subject Classification (2000) 13C12, 13C14 · 13D02, 13H10

1 Correction

In the paper [4], a convention for the depth of the zero module was made which is incorrect and affects several statements and proofs in the paper. The depth of the zero module was set to be -1 , instead of the correct convention, which is ∞ . This has unexpected consequences in the statements of some of the results. Recall from [2] that a finitely generated module M over a Noetherian ring R is said to satisfy condition (S_n) if

$$\text{depth}(M_P) \geq \min\{n, \dim(R_P)\} \text{ for every } P \in \text{Spec}(R).$$

Using the convention that the depth of the module 0 is -1 implies that a module M can satisfy (S_n) only if the support of M is all of $\text{Spec}(R)$, which is not what is needed for various statements in the text. In particular, if R

Craig Huneke
Department of Mathematics
University of Kansas
Lawrence, KS 66045
E-mail: huneke@math.ku.edu

Roger Wiegand
Department of Mathematics
University of Nebraska
Lincoln, NE 68588
E-mail: rwiegand@unl.edu

is Gorenstein, then [2, 3.6] gives that a module satisfies S_2 if and only if it is reflexive, and this characterization requires the convention that the depth of the zero module be ∞ .

Another example is in application of the depth lemma. If

$$0 \rightarrow M \rightarrow N \rightarrow K \rightarrow 0$$

is a short exact sequence of finitely generated modules over a Noetherian local ring R , and $\text{depth}(N) > \text{depth}(K)$, then $\text{depth}(M) = \text{depth}(K) + 1$. This is clearly false under the convention that the depth of 0 be -1 , e.g., if $K = 0$. This necessitates additional arguments in some cases, e.g., when localizing a short exact sequence at a prime not in the support of one of the modules.

Changing the depth of the zero module to ∞ means that some statements need to be revised, and one proof in particular needs some more comment. The two results needing change are Corollary 2.6 and Theorem 2.7. Corollary 2.6 should be modified to the following statement:

Corollary 1 *Let R be a complete intersection, and let M and N be nonzero R -modules such that $\text{Tor}_i^R(M, N) = 0$ for all $i \geq 1$. If $M \otimes_R N$ is maximal Cohen-Macaulay, then so are M and N .*

The proof of this statement follows directly from Proposition 2.5, as in the text. Proposition 2.5 states:

Proposition 1 *Let R be a complete intersection. Let M and N be non-zero finitely generated R -modules such that $\text{Tor}_i^R(M, N) = 0$ for all $i \geq 1$. Then*

$$\text{depth}(M) + \text{depth}(N) = \dim(R) + \text{depth}(M \otimes_R N).$$

One needs to note here that $M \otimes_R N$ cannot be zero since both M and N are nonzero, and R is local.

The only change in Corollary 2.6 is the deletion of the additional assertion that if $M \otimes_R N$ satisfies Serre's property (S_n) , then so do M and N . This is true if the support of both modules is entire spectrum, but could fail otherwise.

Likewise, Theorem 2.7 should be contracted to the following statement:

Theorem 1 (Second Rigidity Theorem) *Let $R = S/(f)$ be a hypersurface, and let M and N be non-zero R -modules, at least one of which has constant rank. If $M \otimes_R N$ is reflexive, then $\text{Tor}_i^R(M, N) = 0$ for all $i \geq 1$.*

The additional conclusion that both M and N are reflexive is never used in the text (except in low dimension, in which case the fact that they are maximal Cohen-Macaulay follows from (2.6)). The problem with this additional conclusion is that if $N_P = 0$, then the hypothesis is satisfied, but one cannot necessarily conclude anything about M_P . One can only conclude results about *both* modules on the intersection of their supports.

The proof of (2.7) must be modified in a few spots to take into account localization at a prime not in the support of one of the modules, but this is

easy to do. One main place this problem arises is in the argument that from the short exact sequence

$$0 \rightarrow M \otimes_R N \rightarrow N^{(n)} \rightarrow M_1 \otimes_R N \rightarrow 0$$

one can conclude that the associated primes of $M_1 \otimes_R N$ are all of height at most one. The proof given has a gap in case a prime P is not in the support of the tensor product $M \otimes_R N$. One must argue separately that if P is associated to $M_1 \otimes_R N$, and if $(M \otimes_R N)_P = 0$, then P is also associated to N , and hence has height zero as N is torsion-free.

Corollary 2.6 and Theorem 2.7 are used in the paper [5] and similar modifications must be made, e.g., in [5] pgs. 169–172.

Acknowledgements We thank Hailong Dao for alerting us to this error, finding places where it was a problem, and for also pointing out that there is not an established convention for what (S_n) means for modules. We use the convention from [2] as stated above. In [3, Def. 5.7.2] and in [1, Sec. 2.1], a module is defined to satisfy (S_n) if

$$\text{depth}_{R_P}(M_P) \geq \min\{n, \dim(M_P)\} \text{ for every } P \in \text{Spec}(R).$$

This is weaker than the version we use.

References

1. Bruns, W. and Herzog, J.: Cohen-Macaulay Rings, Cambridge Studies in Advanced Mathematics, **39**. Cambridge University Press, Cambridge, (1993).
2. Evans, E.G. and Griffith, P.: Syzygies. Cambridge University Press, Cambridge. London Math. Soc. Lect. Notes Ser. **106** (1985)
3. Grothendieck, A. and Dieudonné, J.: Éléments de Géométrie Algébrique IV, Partie 2. Publ. Math. I.H.E.S. **24** (1967).
4. Huneke, C. and Wiegand, R.: Tensor products of modules and the rigidity of Tor. Math. Ann. **299** (1994), 449–476.
5. Huneke, C. and Wiegand, R.: Tensor products of modules, rigidity and local cohomology. Math. Scan. **81** (1997), 161–183.