

## Proof of Perron-Frobenius Theorem

Let  $x = (x_1, x_2, \dots, x_n)^T \in \mathbb{R}^n$  with  $y^T$  denoting exclusively the transpose of vector  $y$ . Let  $\|x\| = \max_i \{|x_i|\}$  be the norm. Then the induced operator norm for matrix  $A = [a_{ij}]$  is  $\|A\| = \max_i \{\sum_j |a_{ij}|\}$ .

Consider a Markov's chain on  $n$  states with transition probabilities  $p_{ij} = \Pr(X_{k+1} = i | X_k = j)$ , independent of  $k$ , and  $P = [p_{ij}]$  the transition matrix. Then  $\sum_{i=1}^n p_{ij} = 1$  for all  $j$ . Let  $p_{ij}^{(t)} = \Pr(X_{k+t} = i | X_k = j)$  and  $P^{(t)} = [p_{ij}^{(t)}]$  be the  $t$ -step transition probability matrix. Then we have  $p_{ij}^{(t)} = \sum_{\ell} p_{i\ell}^{(t-1)} p_{\ell j}$  for all  $i, j$ . In matrix,  $P^{(t)} = P^{(t-1)}P = \dots = P^t$  which is the  $t$ -step transition matrix. If  $q = (q_1, \dots, q_n)^T$  is a probability distribution for the Markovian states at a given iterate with  $q_i \geq 0, \sum q_i = 1$ , then  $Pq$  is again a probability distribution for the states at the next iterate. A probability distribution  $w$  is said to be a steady state distribution if it is invariant under the transition, i.e.  $Pw = w$ . Such a distribution must be an eigenvector of  $P$  and  $\lambda = 1$  must be the corresponding eigenvalue. The existence as well as the uniqueness of the steady state distribution is guaranteed for a class of Markovian chains by the following theorem due to Perron and Frobenius.

**Theorem 1.** *Let  $P = [p_{ij}]$  be a probability transition matrix, i.e.  $p_{ij} \geq 0$  and  $\sum_{i=1}^n p_{ij} = 1$  for every  $j = 1, 2, \dots, n$ . Assume  $P$  is irreducible and transitive in the sense that there is a  $t \geq 1$  so that  $p_{ij}^{(t)} > 0$  for all  $i, j$ . Then 1 is a simple eigenvalue of  $P$  and all other eigenvalues  $\lambda$  satisfy  $|\lambda| < 1$ . Moreover, the unique eigenvector can be chosen to be a probability vector  $w$  and it satisfies  $\lim_{t \rightarrow \infty} P^t = [w, w, \dots, w]$ . Furthermore, for any probability vector  $q$  we have  $P^t q \rightarrow w$  as  $t \rightarrow \infty$ .*

*Proof.* We first prove a claim that  $\lim_{t \rightarrow \infty} p_{ij}^{(t)}$  exist for all  $i, j$  and the limit is independent of  $j$ ,  $\lim_{t \rightarrow \infty} p_{ij}^{(t)} = w_i$ .

Because  $P = [p_{ij}]$  (is irreducible and transitive) has non-zero entries, we have

$$\delta = \min_{ij} p_{ij} > 0.$$

Consider the equation of the  $ij$ th entry of  $P^{t+1} = [p_{ij}^{(t+1)}] = P^t P$ ,

$$p_{ij}^{(t+1)} = \sum_k p_{ik}^{(t)} p_{kj}.$$

Let

$$0 < m_i^{(t)} := \min_j p_{ij}^{(t)} \leq \max_j p_{ij}^{(t)} := M_i^{(t)} < 1.$$

Then, we have

$$m_i^{(t+1)} = \min_j \sum_k p_{ik}^{(t)} p_{kj} \geq m_i^{(t)} \sum_k p_{kj} = m_i^{(t)}.$$

i.e., the sequence  $\{m_i^{(1)}, m_i^{(2)}, \dots\}$  is non-decreasing. Similarly, the upper bound sequence  $\{M_i^{(1)}, M_i^{(2)}, \dots\}$  is non-increasing. As a result, both limits  $\lim_{t \rightarrow \infty} m_i^{(t)} = m_i \leq M_i = \lim_{t \rightarrow \infty} M_i^{(t)}$  exist. We now prove they are equal  $m_i = M_i$ .

To this end, we consider the difference  $M_i^{(t+1)} - m_i^{(t+1)}$ :

$$\begin{aligned}
M_i^{(t+1)} - m_i^{(t+1)} &= \max_j \sum_k p_{ik}^{(t)} p_{kj} - \min_\ell \sum_k p_{ik}^{(t)} p_{k\ell} \\
&= \max_{j,\ell} \sum_k p_{ik}^{(t)} (p_{kj} - p_{k\ell}) \\
&= \max_{j,\ell} [\sum_k^+ p_{ik}^{(t)} (p_{kj} - p_{k\ell}) + \sum_k^- p_{ik}^{(t)} (p_{kj} - p_{k\ell})] \\
&\leq \max_{j,\ell} [M_i^{(t)} \sum_k^+ (p_{kj} - p_{k\ell}) + m_i^{(t)} \sum_k^- (p_{kj} - p_{k\ell})]
\end{aligned} \tag{1}$$

where  $\sum_k^+ p_{ik}^{(t)} (p_{kj} - p_{k\ell})$  means the summation of all non-negative terms  $p_{kj} - p_{k\ell} \geq 0$  and similarly  $\sum_k^- p_{ik}^{(t)} (p_{kj} - p_{k\ell})$  means the summation of all negative terms  $p_{kj} - p_{k\ell} < 0$ .

It is critical to notice the following unexpected equality:

$$\begin{aligned}
\sum_k^- (p_{kj} - p_{k\ell}) &= \sum_k^- p_{kj} - \sum_k^- p_{k\ell} \\
&= 1 - \sum_k^+ p_{kj} - (1 - \sum_k^+ p_{k\ell}) \\
&= \sum_k^+ (p_{k\ell} - p_{kj}) \\
&= -\sum_k^+ (p_{kj} - p_{k\ell}).
\end{aligned}$$

Hence, the inequality (1) becomes

$$M_i^{(t+1)} - m_i^{(t+1)} \leq (M_i^{(t)} - m_i^{(t)}) \max_{j,\ell} \sum_k^+ (p_{kj} - p_{k\ell}).$$

If  $\max_{j,\ell} \sum_k^+ (p_{kj} - p_{k\ell}) = 0$ , which is independent of all  $t$ , it is done that  $M_i^{(t)} = m_i^{(t)}$ . Otherwise, for the pair  $j, \ell$  that gives the maximum let  $r$  be the number of terms in  $k$  for which  $p_{kj} - p_{k\ell} > 0$ , and  $s$  be the number of terms for which  $p_{kj} - p_{k\ell} < 0$ . Then  $r \geq 1$ , and  $\tilde{n} := r + s \geq 1$  as well as  $\tilde{n} \leq n$ . More importantly

$$\begin{aligned}
\sum_k^+ (p_{kj} - p_{k\ell}) &= \sum_k^+ p_{kj} - \sum_k^+ p_{k\ell} \\
&= 1 - \sum_k^- p_{kj} - \sum_k^+ p_{k\ell} \\
&\leq 1 - s\delta - r\delta = 1 - \tilde{n}\delta \\
&\leq 1 - \delta < 1.
\end{aligned}$$

The estimate for the difference  $M_i^{(t+1)} - m_i^{(t+1)}$  at last reduces to

$$M_i^{(t+1)} - m_i^{(t+1)} \leq (1 - \delta)(M_i^{(t)} - m_i^{(t)}) \leq (1 - \delta)^t (M_i^{(1)} - m_i^{(1)}) \rightarrow 0,$$

as  $t \rightarrow \infty$ , showing  $M_i = m_i := w_i$ . As a consequence to the inequality  $m_i^{(t)} \leq p_{ij}^{(t)} \leq M_i^{(t)}$ , we have  $\lim_{t \rightarrow \infty} p_{ij}^{(t)} = w_i$  for all  $j$ . In matrix notation,  $\lim_{t \rightarrow \infty} P^t = [w, w, \dots, w] := W$ , a matrix of equal column vectors.

Next, assume there is a  $k \geq 1$  so that only  $p_{ij}^{(k)} > 0$  for all  $i, j$ . Then the result above implies  $P^{kt} \rightarrow W$  as  $t \rightarrow \infty$ . We need to show that  $P^t \rightarrow W$  as  $t \rightarrow \infty$  as well. This is left as an exercise.

Next, we show that  $\lambda = 1$  is an eigenvalue with eigenvector  $w$ . In fact from the definition of  $w$  above  $\lim_{t \rightarrow \infty} P^t = [w, w, \dots, w]$  and thus  $[w, w, \dots, w] = \lim_{t \rightarrow \infty} P^t = P \lim_{t \rightarrow \infty} P^{t-1} = P[w, w, \dots, w] = [Pw, Pw, \dots, Pw]$  showing  $Pw = w$ .

To show that the eigenvalue  $\lambda = 1$  is simple, two cases are considered. First, let  $x \neq 0$  be an eigenvector  $Px = x$  of the eigenvalue 1. Apply  $P$  to the identity repeatedly to have  $P^t x = x$ . In limit,  $\lim_{t \rightarrow \infty} P^t x = Wx = [w, w, \dots, w]x = wx_1 + wx_2 + \dots + wx_n = (\sum x_j)w = x$ . Denote it by  $\bar{x} = \sum_j x_j$ . Then  $x_i = \bar{x}w_i$  for all  $i$ . Because  $x \neq 0$ , we must have  $\bar{x} = \sum_j x_j \neq 0$ , and that  $x = \bar{x}w$  for some constant  $\bar{x} \neq 0$ , showing that the eigenvector  $w$  is unique up to a constant multiple. Second, let  $x \neq 0$  be a generalized eigenvector of  $\lambda = 1$ . Then there is a constant  $c \neq 0$  so that  $Px = x + cw$  which implies  $P^t x = x + ctw$ . Since  $\lim_{t \rightarrow \infty} P^t x = [w, w, \dots, w]x$  exists on the left, we must have  $c = 0$  on the right, a contradiction. Together we can conclude that the dimension of the generalized eigenspace for  $\lambda = 1$  is 1, i.e., the eigenvalue 1 is simple. In addition, for any probability vector  $q$ , the result above shows  $\lim_{t \rightarrow \infty} P^t q = Wq = \sum_j q_j w = w$  as  $Wx = \sum_j x_j w$  always holds.

Next, let  $\lambda$  be an eigenvalue of  $P$ . Then it is also an eigenvalue for the transpose  $P^T$ . Let  $x$  be an eigenvector of  $\lambda$  of  $P^T$ . Then  $P^T x = \lambda x$  and  $\|\lambda x\| = |\lambda| \|x\| = \|P^T x\| \leq \|P^T\| \|x\|$ . Since  $\|P^T\| = 1$  because  $\sum_{i=1}^n p_{ij} = 1$  we have  $|\lambda| \leq 1$ .

Next, let  $x$  be an eigenvector of an eigenvalue  $\lambda$ . Then we have  $\lim_{t \rightarrow \infty} P^t x = Wx = (\sum x_j)w$  on one hand and  $\lim_{t \rightarrow \infty} P^t x = \lim_{t \rightarrow \infty} \lambda^t x$  on the other hand. So either  $|\lambda| < 1$  in which case  $\lim_{t \rightarrow \infty} \lambda^t x = 0$  and then  $\sum x_j = 0$ , or  $|\lambda| = 1$  in which case  $\lambda = e^{i\theta}$  for some  $\theta$  and the limit  $\lim_{t \rightarrow \infty} \lambda^t = \lim_{t \rightarrow \infty} e^{i\theta t}$  exists since  $\lim_{t \rightarrow \infty} e^{i\theta t} x = \lim_{t \rightarrow \infty} \lambda^t x = \lim_{t \rightarrow \infty} P^t x = (\sum x_j)w$ . The latter case holds if and only if  $\sum x_j \neq 0$  and  $\theta = 0$ , i.e.,  $\lambda = 1$ . This shows that all eigenvalues that is not  $\lambda = 1$  are inside the unit circle and the corresponding eigenspace is  $\{x : \sum_j x_j = 0\}$ , which  $n - 1$  dimensional.  $\square$

References: Bellman(1997); Berman & Plemmons(1994); Frobenius(1908, 1912); Lancaster & Tismenetsky(1985); Marcus & Minc(1984); Perron(1907); Petersen(1983); Seneta(1973).

Ethier and Kurtz, Markov Processes – Characterization and Convergence.