

Name: \_\_\_\_\_

Score: \_\_\_\_\_

---

**Instructions:** You must show supporting work to receive full and partial credits. No text book, notes, formula sheets allowed.

---

---

**1(15 pts)** Verify that  $\ln y + xy = 5$  defines an implicit solution to the equation  $\frac{dy}{dx} = -y^2/(1 + xy)$ .

---

**2(18 pts)** Find the solution to the initial value problem  $\frac{dy}{dx} = \frac{x}{1+y}$ ,  $y(0) = 1$ .

---

**3(18 pts)** (a) Use the Euler method to approximate the solution to the initial value problem  $y' = (x + 1)(y^4 + 1)$ ,  $y(0) = 0$  at these points  $x = 0, 0.25, 0.5$ .  
(b) Plot your approximating solution in the  $xy$  plane.

---

**4(16 pts)** Find the general solution to the equation  $y' - \frac{1}{(1+t)}y = 1$ .

---

**5(18 pts)** Consider the equation

$$\frac{dy}{dt} = f(y) = (1 - y)(y - k),$$

where  $k$  is a parameter.

- (a) Sketch a phase line for the equation for some value of  $k$  between 0 and 1. Classify each equilibrium point as sink, source, or node.
  - (b) Sketch a few typical solutions qualitatively with  $y$  v.s. the time  $t$ , for the same value of  $k$  you used in (a) above.
  - (c) For what value of  $k$  does the equation has only one equilibrium point? Classify that point as sink, source, or node.
- 
- 6(15 pts)**
- Blood carries a drug into an organ at a rate of
- $3 \text{ cm}^3/\text{sec}$
- and leaves at the same rate. The organ has a liquid volume of
- $125 \text{ cm}^3$
- which has no trace of the drug initially. If the concentration of the drug in the blood entering the organ is
- $0.2 \text{ g/cm}^3$
- , write down an initial value problem for the amount of drug in gram at any time
- $t$
- . (
- Remark:*
- Don't think about by-passing differential equations to solve this problem.)
- 

**END**