

## Math 970 Homework

**Due:** Sept. 18

1. If  $(X, \mathcal{T})$  is a topological space,  $Y$  is a set, and  $f : X \rightarrow Y$  is a function, show that

$$\mathcal{T}' = \{\mathcal{U}' \subseteq Y : f^{-1}(\mathcal{U}') \in \mathcal{T}\}$$

is the finest topology on  $Y$  for which  $f : (X, \mathcal{T}) \rightarrow (Y, \mathcal{T}')$  is continuous.

(Note that this problem is actually asking you to show three things...)

2. If  $(X, \mathcal{T})$  is a topological space, and  $A \subseteq X$ , then  $A \in \mathcal{T}$  if and only if

for all  $x \in A$ , there is a  $U \in \mathcal{T}$  so that  $x \in U \subseteq A$

3. Show that  $\mathcal{B} = \{(a, \infty) \times (b, \infty) : a, b \in \mathbb{R}\}$  is a basis for a topology  $\mathcal{T}$  on  $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$ , which is coarser than the usual Euclidean topology on  $\mathbb{R}^2$ . Show that  $\mathcal{B}' = \{[a, \infty) \times [b, \infty) : a, b \in \mathbb{R}\}$  is a basis for a topology  $\mathcal{T}'$  which is strictly finer than  $\mathcal{T}$ , and not comparable to the usual Euclidean topology.
4. Show that, in general, if  $\mathcal{B}$  and  $\mathcal{B}'$  are both bases for topologies on  $X$ , that  $\mathcal{B} \cap \mathcal{B}'$  and  $\mathcal{B} \cup \mathcal{B}'$  need not be. Show, however, that  $\mathcal{B}'' = \{B \cap B' : B \in \mathcal{B}, B' \in \mathcal{B}'\}$  is a basis for a topology, and  $\mathcal{T}(\mathcal{B}'')$  is the coarsest topology containing both  $\mathcal{B}$  and  $\mathcal{B}'$ .
5. Show that the topology generated by a basis  $\mathcal{B}$  is the coarsest topology containing  $\mathcal{B}$  (i.e., it is the intersection of all such topologies).