

Chain rule

October 2, 2013

Composite function

Definition

The function $f(g(t))$ is a function of function, or **composite function**, in which there is an **inside function** and an **outside function**.

- Use a new variable u for the inside function to express each of the following as a composite functions:

Composite function

Definition

The function $f(g(t))$ is a function of function, or **composite function**, in which there is an **inside function** and an **outside function**.

- Use a new variable u for the inside function to express each of the following as a composite functions:
- $f(x) = \ln(3t + 1)$

Composite function

Definition

The function $f(g(t))$ is a function of function, or **composite function**, in which there is an **inside function** and an **outside function**.

- Use a new variable u for the inside function to express each of the following as a composite functions:
- $f(x) = \ln(3t + 1)$
- $y = 5(2x^2 + 4)^2$

Composite function

Definition

The function $f(g(t))$ is a function of function, or **composite function**, in which there is an **inside function** and an **outside function**.

- Use a new variable u for the inside function to express each of the following as a composite functions:
- $f(x) = \ln(3t + 1)$
- $y = 5(2x^2 + 4)^2$
- $P = (3x + 1) + 4e^{9x+3}$

Composite function

Definition

The function $f(g(t))$ is a function of function, or **composite function**, in which there is an **inside function** and an **outside function**.

- Use a new variable u for the inside function to express each of the following as a composite functions:
- $f(t) = \ln(3t + 1) \rightarrow f(u) = \ln u$
- $y = 5(2x^2 + 4)^2 \rightarrow y = 5u^2$
- $P = (3x + 1) + 4e^{9x+3} \rightarrow P = u + 4e^{3u}$

Derivative of a composite functions

- Suppose $y = f(z)$ and $z = g(t)$ for some outside function f and inside function g , where f and g are *differentiable*. A small change in t , Δt , generates a small change in z , Δz . In turn, Δz generates a small change in y , Δy . Provided Δt and Δz are not zero, we have

$$\frac{\Delta y}{\Delta t} = \frac{\Delta y}{\Delta z} \cdot \frac{\Delta z}{\Delta t}$$

Derivative of a composite functions

- Suppose $y = f(z)$ and $z = g(t)$ for some outside function f and inside function g , where f and g are *differentiable*. A small change in t , Δt , generates a small change in z , Δz . In turn, Δz generates a small change in y , Δy . Provided Δt and Δz are not zero, we have

$$\frac{\Delta y}{\Delta t} = \frac{\Delta y}{\Delta z} \cdot \frac{\Delta z}{\Delta t}$$

- The derivative $\frac{dy}{dt}$ is the limit of the quotient $\frac{\Delta y}{\Delta t}$ as Δt gets smaller and smaller.

Derivative of a composite functions

- Suppose $y = f(z)$ and $z = g(t)$ for some outside function f and inside function g , where f and g are *differentiable*. A small change in t , Δt , generates a small change in z , Δz . In turn, Δz generates a small change in y , Δy . Provided Δt and Δz are not zero, we have

$$\frac{\Delta y}{\Delta t} = \frac{\Delta y}{\Delta z} \cdot \frac{\Delta z}{\Delta t}$$

- The derivative $\frac{dy}{dt}$ is the limit of the quotient $\frac{\Delta y}{\Delta t}$ as Δt gets smaller and smaller.
- The derivative $\frac{dz}{dt}$ is the limit of the quotient $\frac{\Delta z}{\Delta t}$ as Δt gets smaller and smaller.

Derivative of a composite functions

- Suppose $y = f(z)$ and $z = g(t)$ for some outside function f and inside function g , where f and g are *differentiable*. A small change in t , Δt , generates a small change in z , Δz . In turn, Δz generates a small change in y , Δy . Provided Δt and Δz are not zero, we have

$$\frac{\Delta y}{\Delta t} = \frac{\Delta y}{\Delta z} \cdot \frac{\Delta z}{\Delta t}$$

- The derivative $\frac{dy}{dt}$ is the limit of the quotient $\frac{\Delta y}{\Delta t}$ as Δt gets smaller and smaller.
- The derivative $\frac{dz}{dt}$ is the limit of the quotient $\frac{\Delta z}{\Delta t}$ as Δt gets smaller and smaller.
- The derivative $\frac{dy}{dz}$ is the limit of the quotient $\frac{\Delta y}{\Delta z}$ as Δz gets smaller and smaller.

Derivative of a composite functions

- Suppose $y = f(z)$ and $z = g(t)$ for some outside function f and inside function g , where f and g are *differentiable*. A small change in t , Δt , generates a small change in z , Δz . In turn, Δz generates a small change in y , Δy . Provided Δt and Δz are not zero, we have

$$\frac{\Delta y}{\Delta t} = \frac{\Delta y}{\Delta z} \cdot \frac{\Delta z}{\Delta t}$$

- The derivative $\frac{dy}{dt}$ is the limit of the quotient $\frac{\Delta y}{\Delta t}$ as Δt gets smaller and smaller.
- The derivative $\frac{dz}{dt}$ is the limit of the quotient $\frac{\Delta z}{\Delta t}$ as Δt gets smaller and smaller.
- The derivative $\frac{dy}{dz}$ is the limit of the quotient $\frac{\Delta y}{\Delta z}$ as Δz gets smaller and smaller.
- $\frac{dy}{dt} = \frac{dy}{dz} \cdot \frac{dz}{dt}$.

Theorem

Let $y = f(z)$ and $z = g(t)$ be two differentiable functions, then the derivative of the composite function $y = f(g(t))$ is given by

$$\frac{dy}{dt} = \frac{dy}{dz} \cdot \frac{dz}{dt}.$$

In other word, the derivative of a composite function is the derivative of the outside function times the derivative of the inside function:

$$\frac{d}{dt}(f(g(t))) = f'(g(t)) \cdot g'(t).$$

Example

- The amount of gas G , in gallons, consumed by a car depends on the distance traveled, s , in miles, and s depends on the time t , in hours. If 0.05 gallons of gas is consumed for each mile traveled, and the car is traveling at 30 miles/hour, **how fast is gas being consumed?**

Example

- The amount of gas G , in gallons, consumed by a car depends on the distance traveled, s , in miles, and s depends on the time t , in hours. If 0.05 gallons of gas is consumed for each mile traveled, and the car is traveling at 30 miles/hour, **how fast is gas being consumed?**
- Unit= gallon/hour

Example

- The amount of gas G , in gallons, consumed by a car depends on the distance traveled, s , in miles, and s depends on the time t , in hours. If 0.05 gallons of gas is consumed for each mile traveled, and the car is traveling at 30 miles/hour, **how fast is gas being consumed?**
- Unit= gallon/hour
- Rate of gas consumed with respect to distance = $\frac{dG}{ds} = 0.05$ gallons/mile.

Example

- The amount of gas G , in gallons, consumed by a car depends on the distance traveled, s , in miles, and s depends on the time t , in hours. If 0.05 gallons of gas is consumed for each mile traveled, and the car is traveling at 30 miles/hour, **how fast is gas being consumed?**
- Unit= gallon/hour
- Rate of gas consumed with respect to distance = $\frac{dG}{ds} = 0.05$ gallons/mile.
- Rate of distance increasing with respect to time = $\frac{ds}{dt} = 30$ miles/hour.

Example

- The amount of gas G , in gallons, consumed by a car depends on the distance traveled, s , in miles, and s depends on the time t , in hours. If 0.05 gallons of gas is consumed for each mile traveled, and the car is traveling at 30 miles/hour, **how fast is gas being consumed?**
- Unit= gallon/hour
- Rate of gas consumed with respect to distance = $\frac{dG}{ds} = 0.05$ gallons/mile.
- Rate of distance increasing with respect to time = $\frac{ds}{dt} = 30$ miles/hour.
- $\frac{dG}{dt} = \frac{dG}{ds} \cdot \frac{ds}{dt} = 0.05 \cdot 30 = 1.5$ gallons/hour.

Examples

Find the derivative of the following functions

- (a) $f(t) = \ln(3t + 1)$

Examples

Find the derivative of the following functions

- (a) $f(t) = \ln(3t + 1)$
- (b) $y = 5(2x^2 + 4)^2$

Examples

Find the derivative of the following functions

- (a) $f(t) = \ln(3t + 1)$
- (b) $y = 5(2x^2 + 4)^2$
- (c) $P = (3x + 1) + 4e^{9x+3}$

Example

(a) $f(t) = \ln(3t + 1)$

- Write the composite form with new variable u : $f(u) = \ln u$ where $u = 3t + 1$.

Example

(a) $f(t) = \ln(3t + 1)$

- Write the composite form with new variable u : $f(u) = \ln u$ where $u = 3t + 1$.
- We need to find $df/dt = df/du \cdot du/dt$, so we need to find df/du and du/dt .

Example

(a) $f(t) = \ln(3t + 1)$

- Write the composite form with new variable u : $f(u) = \ln u$ where $u = 3t + 1$.
- We need to find $df/dt = df/du \cdot du/dt$, so we need to find df/du and du/dt .
- $df/du = 1/u$ and $du/dt = 3$.

Example

(a) $f(t) = \ln(3t + 1)$

- Write the composite form with new variable u : $f(u) = \ln u$ where $u = 3t + 1$.
- We need to find $df/dt = df/du \cdot du/dt$, so we need to find df/du and du/dt .
- $df/du = 1/u$ and $du/dt = 3$.
- $df/dt = df/du \cdot du/dt = (\ln u) \cdot 3 = 3 \ln(3t + 1)$, since $u = 3t + 1$.

Theorem

$$\frac{d}{dx}(x^n) = nx^{n-1}, \quad \frac{d}{dx}(a^x) = (\ln a)a^x, \quad \frac{d}{dx}(\ln x) = \frac{1}{x}.$$

Example

$$(b) \ y = 5(2x^2 + 4)^2$$

Example

$$(c) \ P = (3x + 1) + 4e^{9x+3}$$

Examples

Find the derivative of

- $\sqrt{1 + 2e^{5t}}$

Examples

Find the derivative of

- $\sqrt{1 + 2e^{5t}}$
- $\ln(q + 4^q)$

Examples

Find the derivative of

- $\sqrt{1 + 2e^{5t}}$
- $\ln(q + 4^q)$
- e^{-x^2+3x+1}

Relative Rate of change and logarithm

The relative rate of change of a function $z = f(t)$ to be

$$\frac{f'(t)}{f(t)} = \frac{1}{z} \cdot f'(t) = \frac{1}{z} \frac{dz}{dt}.$$

Since

$$\frac{d}{dt}(\ln z) = \frac{1}{z} \frac{dz}{dt},$$

we have

Theorem

For any positive function $f(t)$,

$$\text{Relative rate of change of } f(t) = \frac{d}{dt}(\ln f(t)).$$

Find the relative rate of change of the exponential function
 $z = P_0 e^{kt}$.

Find the relative rate of change of the exponential function $z = P_0 e^{kt}$.

- $\ln z = \ln(P_0 e^{kt}) = \ln P_0 + \ln(e^{kt}) = \ln P_0 + kt$

Examples

Find the relative rate of change of the exponential function $z = P_0 e^{kt}$.

- $\ln z = \ln(P_0 e^{kt}) = \ln P_0 + \ln(e^{kt}) = \ln P_0 + kt$
- Relative rate of change of $f(t)$ is $\frac{d}{dt}(\ln z) = k$.

Examples

- The surface area S of a mammal, in cm^2 , is a functions of the body mass, M , of the mammal, in kg, and is given by $S = 1095 \cdot M^{2/3}$. Find the relative rate of change of S with respect to M and evaluate for human with body mass 70 kg. Interpret the answer.

Examples

- The surface area S of a mammal, in cm^2 , is a functions of the body mass, M , of the mammal, in kg, and is given by $S = 1095 \cdot M^{2/3}$. Find the relative rate of change of S with respect to M and evaluate for human with body mass 70 kg. Interpret the answer.
- $\ln S = \ln(1095 \cdot M^{2/3}) = \ln 1095 + \frac{2}{3} \ln M$

Examples

- The surface area S of a mammal, in cm^2 , is a functions of the body mass, M , of the mammal, in kg, and is given by $S = 1095 \cdot M^{2/3}$. Find the relative rate of change of S with respect to M and evaluate for human with body mass 70 kg. Interpret the answer.
- $\ln S = \ln(1095 \cdot M^{2/3}) = \ln 1095 + \frac{2}{3} \ln M$
- Relative rate of change is $\frac{d}{dM}(\ln S)$.

Examples

The surface area S of a mammal, in cm^2 , is a function of the body mass, M , of the mammal, in kg, and is given by $S = 1095 \cdot M^{2/3}$. Find the relative rate of change of S with respect to M and evaluate for human with body mass 70 kg. Interpret the answer.

- $\ln S = \ln(1095 \cdot M^{2/3}) = \ln 1095 + \frac{2}{3} \ln M$
- Relative rate of change is $\frac{d}{dM}(\ln S) = \frac{2}{3} \cdot \frac{1}{M}$.

Examples

The surface area S of a mammal, in cm^2 , is a functions of the body mass, M , of the mammal, in kg, and is given by $S = 1095 \cdot M^{2/3}$. Find the relative rate of change of S with respect to M and evaluate for human with body mass 70 kg. Interpret the answer.

- $\ln S = \ln(1095 \cdot M^{2/3}) = \ln 1095 + \frac{2}{3} \ln M$
- Relative rate of change is $\frac{d}{dM}(\ln S) = \frac{2}{3} \cdot \frac{1}{M}$.

Examples

The surface area S of a mammal, in cm^2 , is a function of the body mass, M , of the mammal, in kg, and is given by $S = 1095 \cdot M^{2/3}$. Find the relative rate of change of S with respect to M and evaluate for human with body mass 70 kg. Interpret the answer.

- $\ln S = \ln(1095 \cdot M^{2/3}) = \ln 1095 + \frac{2}{3} \ln M$
- Relative rate of change is $\frac{d}{dM}(\ln S) = \frac{2}{3} \cdot \frac{1}{M}$.
- For human with body mass $M = 70$ kg, we have:

$$\text{Relative rate of change} = \frac{2}{3} \frac{1}{70} = 0.00095 = 0.95\%.$$

Examples

The surface area S of a mammal, in cm^2 , is a function of the body mass, M , of the mammal, in kg, and is given by $S = 1095 \cdot M^{2/3}$. Find the relative rate of change of S with respect to M and evaluate for human with body mass 70 kg. Interpret the answer.

- $\ln S = \ln(1095 \cdot M^{2/3}) = \ln 1095 + \frac{2}{3} \ln M$
- Relative rate of change is $\frac{d}{dM}(\ln S) = \frac{2}{3} \cdot \frac{1}{M}$.
- For human with body mass $M = 70$ kg, we have:

$$\text{Relative rate of change} = \frac{2}{3} \frac{1}{70} = 0.00095 = 0.95\%.$$

- The surface area of a human with body mass 70 kg increases by about 0.95 % if the body mass increases by 1 kg.