


# The Definite Integral as Area

November 6, 2013


# The case when $f(x)$ is nonnegative

- Let  $f(x)$  is nonnegative and continuous for  $a \leq x \leq b$ .




# The case when $f(x)$ is nonnegative

- Let  $f(x)$  is nonnegative and continuous for  $a \leq x \leq b$ .
- The left- and right-hand sums tends to the area under the graph of  $f$  between  $a$  and  $b$  as  $\Delta t$  smaller and smaller.




# The case when $f(x)$ is nonnegative

- Let  $f(x)$  is nonnegative and continuous for  $a \leq x \leq b$ .
- The left- and right-hand sums tends to the area under the graph of  $f$  between  $a$  and  $b$  as  $\Delta t$  smaller and smaller.
- The definite integral  $\int_a^b f(x)dx$  is the area.




# The case when $f(x)$ is nonnegative

- Let  $f(x)$  is nonnegative and continuous for  $a \leq x \leq b$ .
- The left- and right-hand sums tends to the area under the graph of  $f$  between  $a$  and  $b$  as  $\Delta t$  smaller and smaller.
- The definite integral  $\int_a^b f(x)dx$  is the area.



# The case when $f(x)$ is nonnegative

- Let  $f(x)$  is nonnegative and continuous for  $a \leq x \leq b$ .
- The left- and right-hand sums tends to the area under the graph of  $f$  between  $a$  and  $b$  as  $\Delta t$  smaller and smaller.
- The definite integral  $\int_a^b f(x)dx$  is the area.




# Example 1

Find the area under the graph of  $y = 10x(3^{-x})$  between  $x = 0$  and  $x = 3$ .

# Example 1

Find the area under the graph of  $y = 10x(3^{-x})$  between  $x = 0$  and  $x = 3$ .




$$\text{Area shaded} = \int_0^3 f(x) \, dx, \text{ where } f(x) = 10x3^{-x}.$$



# Example 1

Find the area under the graph of  $y = 10x(3^{-x})$  between  $x = 0$  and  $x = 3$ .

- Since the function is nonnegative for  $0 \leq x \leq 3$ , the area is equal to  $\int_0^3 10x(3^{-x})dx$ .




# Example 1

Find the area under the graph of  $y = 10x(3^{-x})$  between  $x = 0$  and  $x = 3$ .

- Since the function is nonnegative for  $0 \leq x \leq 3$ , the area is equal to  $\int_0^3 10x(3^{-x})dx$ .
- Using calculator or computer, we can obtain

$$\int_0^3 10x(3^{-x})dx = 6.967$$



# Example 1

Find the area under the graph of  $y = 10x(3^{-x})$  between  $x = 0$  and  $x = 3$

- Since the function is nonnegative for  $0 \leq x \leq 3$ , the area is equal to  $\int_0^3 10x(3^{-x})dx$ .
- Using calculator or computer, we can obtain

$$\int_0^3 10x(3^{-x})dx = 6.967$$

- See the calculator manual on Oncourse.

# Example 1

Find the area under the graph of  $y = 10x(3^{-x})$  between  $x = 0$  and  $x = 3$

- Since the function is nonnegative for  $0 \leq x \leq 3$ , the area is equal to  $\int_0^3 10x(3^{-x})dx$ .
- Using calculator or computer, we can obtain

$$\int_0^3 10x(3^{-x})dx = 6.967$$

- See the calculator manual on Oncourse.
- Use math softwares, such as Maple, Mathlab,

# Example 1

Find the area under the graph of  $y = 10x(3^{-x})$  between  $x = 0$  and  $x = 3$

- Since the function is nonnegative for  $0 \leq x \leq 3$ , the area is equal to  $\int_0^3 10x(3^{-x})dx$ .
- Using calculator or computer, we can obtain

$$\int_0^3 10x(3^{-x})dx = 6.967$$


- See the calculator manual on Oncourse.
- Use math softwares, such as Maple, Mathlab,
- Online free resource:  
<http://www.numberempire.com/definiteintegralcalculator.php>

# Relationship Between Definite Integral and Area : When $f(x)$ is NOT nonnegative.

Consider the relationship between the definite integral  $\int_{-1}^1 (x^2 - 1) dx$  and the area between the parabola  $y = x^2 - 1$  and the  $x$ -axis.


# Relationship Between Definite Integral and Area : When $f(x)$ is NOT nonnegative.

Consider the relationship between the definite integral  $\int_{-1}^1 (x^2 - 1) dx$  and the area between the parabola  $y = x^2 - 1$  and the  $x$ -axis.



# Relationship Between Definite Integral and Area : When $f(x)$ is NOT nonnegative.

- The parabola lies below the  $x$ -axis between  $x = -1$  and  $x = 1$ .




Integral  $\int_{-1}^1 f(x) \, dx$ , where  $f(x) = x^2 - 1$ , is negative of shaded area.



# Relationship Between Definite Integral and Area : When $f(x)$ is NOT nonnegative.

- The parabola lies below the  $x$ -axis between  $x = -1$  and  $x = 1$ .
- $\int_{-1}^1 (x^2 - 1) dx = -\text{Area} = -1.33$



Integral  $\int_{-1}^1 f(x) dx$ , where  $f(x) = x^2 - 1$ , is negative of shaded area.

# Conclusion

When  $f(x)$  is positive for some  $x$ -values and negative for others, and  $a < b$ :


$\int_a^b f(x)dx$  is the sum of the areas above the  $x$ -axis, **counted positively**, and the areas below the  $x$ -axis, **counted negatively**.

## Example 3

Interpret the definite integral  $\int_0^4 (x^3 - 7x^2 + 11x) dx$  in terms of areas.


## Example 3

Interpret the definite integral  $\int_0^4 (x^3 - 7x^2 + 11x) dx$  in terms of areas.




## Example 3

- the graph of  $f(x)$  is crossing below the  $x$ -axis about  $x = 2.4$ .




## Example 3

- the graph of  $f(x)$  is crossing below the  $x$ -axis about  $x = 2.4$ .
- Use calculator  $\int_0^4 (x^3 - 7x^2 + 11x) dx = 2.67$




## Example 3

- the graph of  $f(x)$  is crossing below the  $x$ -axis about  $x = 2.4$ .
- Use calculator  $\int_0^4 (x^3 - 7x^2 + 11x) dx = 2.67$
- $\int_0^4 (x^3 - 7x^2 + 11x) dx = A_1 - A_2$ .



## Example 3


- the graph of  $f(x)$  is crossing below the  $x$ -axis about  $x = 2.4$ .






## Example 3

- the graph of  $f(x)$  is crossing below the  $x$ -axis about  $x = 2.4$ .
- Use calculator  $\int_0^4 (x^3 - 7x^2 + 11x) dx = 2.67$




## Example 3

- the graph of  $f(x)$  is crossing below the  $x$ -axis about  $x = 2.4$ .
- Use calculator  $\int_0^4 (x^3 - 7x^2 + 11x) dx = 2.67$
- $\int_0^4 (x^3 - 7x^2 + 11x) dx = A_1 - A_2$ .



## Example 3

- the graph of  $f(x)$  is crossing below the  $x$ -axis about  $x = 2.4$ .
- Use calculator  $\int_0^4 (x^3 - 7x^2 + 11x) dx = 2.67$
- $\int_0^4 (x^3 - 7x^2 + 11x) dx = A_1 - A_2$ .
- Break the integral into two parts  
 $A_1 \approx \int_0^{2.4} (x^3 - 7x^2 + 11x) dx \approx 7.72$  and  
 $-A_2 \approx \int_{2.4}^4 (x^3 - 7x^2 + 11x) dx \approx -5.05$ .




Assume  $f(x)$  is continuous from  $a$  to  $b$ , and  $a < c < b$ . Then

$$\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$$


## Example 4

For each functions graphed in the figure, decide whether  $\int_0^5 f(x)dx$  is positive, negative or approximately zero.




## Example 4

For each functions graphed in the figure, decide whether  $\int_0^5 f(x)dx$  is positive, negative or approximately zero.




## Example 4

For each functions graphed in the figure, decide whether  $\int_0^5 f(x)dx$  is positive, negative or approximately zero.



# Area between two curves





# Area between two curves

If  $g(x) \leq f(x)$  for  $a \leq x \leq b$ . Then

Area between graphs of  $f(x)$  and  $g(x)$  for  $a \leq x \leq b$  is equal to  $\int_a^b (f(x) - g(x)) dx$ .

Assume  $f(x)$  and  $g(x)$  are two continuous functions for  $a \leq x \leq b$ , and  $A$  and  $B$  are two constants. Then


$$\int_a^b (Af(x) + Bg(x))dx = A \int_a^b f(x)dx + B \int_a^b g(x)dx$$

## Example 5

Graph  $f(x) = 4x - x^2$  and  $g(x) = \frac{1}{2}x^{3/2}$  for  $x \geq 0$ . Use a definite integral to estimate the area enclosed by the graphs of these two functions.

## Example 5


Graph  $f(x) = 4x - x^2$  and  $g(x) = \frac{1}{2}x^{3/2}$  for  $x \geq 0$ . Use a definite integral to estimate the area enclosed by the graphs of these two functions.



## Example 5

Graph  $f(x) = 4x - x^2$  and  $g(x) = \frac{1}{2}x^{3/2}$  for  $x \geq 0$ . Use a definite integral to estimate the area enclosed by the graphs of these two functions.


- Two graphs cross at  $x = 0$  and  $x \approx 3.1$ .



## Example 5

Graph  $f(x) = 4x - x^2$  and  $g(x) = \frac{1}{2}x^{3/2}$  for  $x \geq 0$ . Use a definite integral to estimate the area enclosed by the graphs of these two functions.


- Two graphs cross at  $x = 0$  and  $x \approx 3.1$ .
- Between  $x = 0$  and  $x = 3.1$  the graph of  $f(x) = 4x - x^2$  is above the graph of  $g(x) = \frac{1}{2}x^{3/2}$ .



## Example 5

Graph  $f(x) = 4x - x^2$  and  $g(x) = \frac{1}{2}x^{3/2}$  for  $x \geq 0$ . Use a definite integral to estimate the area enclosed by the graphs of these two functions.

- Two graphs cross at  $x = 0$  and  $x \approx 3.1$ .
- Between  $x = 0$  and  $x = 3.1$  the graph of  $f(x) = 4x - x^2$  is above the graph of  $g(x) = \frac{1}{2}x^{3/2}$ .
- The Area is (approximately)  $\int_0^{3.1} (4x - x^2 - \frac{1}{2}x^{3/2}) dx$ .



# Area between two curves

For  $a \leq x \leq b$

$\int_a^b (f(x) - g(x)) dx$  is the sum of the areas when  $f(x) > g(x)$ , counted positively, and the areas when  $f(x) < g(x)$ , counted negatively.



# Area between two curves

$$\int_0^{10} (\sin(x) - \sin(-x)) dx$$

