

# Second derivative

September 26, 2013

# What does the second derivative tell us?

- The derivative of derivative is called *second derivative*, and written by  $f''$ .

# What does the second derivative tell us?

- The derivative of derivative is called *second derivative*, and written by  $f''$ .
- If  $y = f(x)$ , then the second derivative can also be written by  $\frac{d^2y}{dx^2}$

# What does the second derivative tell us?

- The derivative of derivative is called *second derivative*, and written by  $f''$ .
- If  $y = f(x)$ , then the second derivative can also be written by  $\frac{d^2y}{dx^2}$
- which means  $\frac{d}{dx} \left( \frac{dy}{dx} \right)$ , i.e. the derivative of  $\frac{dy}{dx}$ .

# What does the second tell us?

## Theorem

- *If  $f' > 0$  on an interval, then  $f$  is increasing over that interval.*

## Theorem

- *If  $f'' > 0$  on an interval, then  $f'$  is increasing over that interval.*

# What does the second tell us?

## Theorem

- If  $f' > 0$  on an interval, then  $f$  is increasing over that interval.
- If  $f' < 0$  on an interval, then  $f$  is decreasing over that interval.

## Theorem

- If  $f'' > 0$  on an interval, then  $f'$  is increasing over that interval.
- If  $f'' < 0$  on an interval, then  $f'$  is decreasing over that interval.

# What does the second tell us?

## Theorem

- *If  $f'' > 0$  on an interval, then  $f'$  is increasing over that interval, so the graph of  $f$  is concave up there.*

# What does the second tell us?

## Theorem

- If  $f'' > 0$  on an interval, then  $f'$  is increasing over that interval, so the graph of  $f$  is concave up there.
- If  $f'' < 0$  on an interval, then  $f'$  is decreasing over that interval, so the graph of  $f$  is concave down there.



# Interpretation of the second derivative as a rate of change

- We think that the derivative as a rate of change, then the second derivative as the *rate of change of rate of change*.

# Interpretation of the second derivative as a rate of change

- We think that the derivative as a rate of change, then the second derivative as the *rate of change of rate of change*.
- If the second derivative is positive, the rate of change is increasing; if the second derivative is negative, the rate of change is decreasing.

# Example

- A population,  $P$ , growing in a confined environment often follow a *logistic* growth curve.

# Example

- A population,  $P$ , growing in a confined environment often follow a *logistic* growth curve.
- $f(x) = \frac{1}{1 + e^{-x}}$ .

# Example

- A population,  $P$ , growing in a confined environment often follow a *logistic* growth curve.
- $f(x) = \frac{1}{1 + e^{-x}}$ .
- Describe how the rate at which the population is increasing changes over time.

# Example

- A population,  $P$ , growing in a confined environment often follow a *logistic* growth curve.
- $f(x) = \frac{1}{1 + e^{-x}}$ .
- Describe how the rate at which the population is increasing changes over time.
- What is the sign of the second derivative  $\frac{d^2P}{dx^2}$ ?

# Example

- A population,  $P$ , growing in a confined environment often follow a *logistic* growth curve.
- $f(x) = \frac{1}{1 + e^{-x}}$ .
- Describe how the rate at which the population is increasing changes over time.
- What is the sign of the second derivative  $\frac{d^2P}{dx^2}$ ?
- What is the practical interpretation of  $t^*$  and  $L$ ?

# Example

- Initially, the population is increasing, and at an increasing rate.



# Example

- Initially, the population is increasing, and at an increasing rate.
- At  $t^*$  the population is increasing fastest.

# Example

- Initially, the population is increasing, and at an increasing rate.
- At  $t^*$  the population is increasing fastest.
- $L$  is the limiting value of the population.

# Example

Year, $t$	1972	1975	1980	1985	1990	1995	2000	2005
1000s of abortions, $A$	587	1034	1554	1589	1609	1359	1313	1206

- Table show the number of abortions per year.

# Example

Year, $t$	1972	1975	1980	1985	1990	1995	2000	2005
1000s of abortions, $A$	587	1034	1554	1589	1609	1359	1313	1206

- Table show the number of abortions per year.
- (a) Calculate the average rate of change for the time interval shown between 1972 an 2005.

# Example

Year, t	1972	1975	1980	1985	1990	1995	2000	2005
1000s of abortions, A	587	1034	1554	1589	1609	1359	1313	1206

- Table show the number of abortions per year.
- (a) Calculate the average rate of change for the time interval shown between 1972 an 2005.
- (b) What can you say about the sign of  $d^2A/dt^2$  during the period 1972-1995?