

A CHANGE OF RINGS RESULT FOR MATLIS REFLEXIVITY

DOUGLAS J. DAILEY AND THOMAS MARLEY

ABSTRACT. Let R be a commutative Noetherian ring and E the minimal injective cogenerator of the category of R -modules. An R -module M is (Matlis) reflexive if the natural evaluation map $M \rightarrow \text{Hom}_R(\text{Hom}_R(M, E), E)$ is an isomorphism. We prove that if S is a multiplicatively closed subset of R and M is an R_S -module which is reflexive as an R -module, then M is a reflexive R_S -module. The converse holds when S is the complement of the union of finitely many nonminimal primes of R , but fails in general.

1. INTRODUCTION

Let R be a commutative Noetherian ring and E the minimal injective cogenerator of the category of R -modules; i.e., $E = \bigoplus_{m \in \Lambda} E_R(R/m)$, where Λ denotes the set of maximal ideals of R and $E_R(-)$ denotes the injective hull. An R -module M is said to be (*Matlis reflexive*) if the natural evaluation map $M \rightarrow \text{Hom}_R(\text{Hom}_R(M, E), E)$ is an isomorphism. In [1], the authors assert the following “change of rings” principal for Matlis reflexivity ([1, Lemma 2]): *Let S be a multiplicatively closed subset of R and suppose M is an R_S -module. Then M is reflexive as an R -module if and only if M is reflexive as an R_S -module.* However, the proof given in [1] is incorrect (see Examples 3.1-3.3) and in fact the “if” part is false in general (cf. Proposition 3.4). In this note, we prove the following:

Theorem 1.1. *Let R be a Noetherian ring, S a multiplicatively closed subset of R , and M an R_S -module.*

- (a) *If M is reflexive as an R -module then M is reflexive as an R_S -module.*
- (b) *If $S = R \setminus (p_1 \cup \dots \cup p_r)$ where each p_i is a maximal ideal or a nonminimal prime ideal, then the converse to (a) holds.*

2. MAIN RESULTS

Throughout this section R will denote a Noetherian ring and S a multiplicatively closed subset of R . We let E_R (or just E if the ring is clear) denote the minimal injective cogenerator of the category of R -modules as defined in the introduction. A semilocal ring is said to be complete if it is complete with respect to the J -adic topology, where J is the Jacobson radical.

We will make use of the main result of [1]:

Date: May 11, 2016.

2010 Mathematics Subject Classification. Primary 13C05; Secondary 13C13.

Key words and phrases. Matlis reflexive, minimal injective cogenerator.

The first author was partially supported by U.S. Department of Education grant P00A120068 (GAANN).

Theorem 2.1. ([1, Theorem 12]) *Let R be a Noetherian ring, M an R -module, and $I = \text{Ann}_R M$. Then M is reflexive if and only if R/I is a complete semilocal ring and there exists a finitely generated submodule N of M such that M/N is Artinian.*

We remark that the validity of this theorem does not depend on [1, Lemma 2], as the proof of [1, Theorem 12] uses this lemma only in a special case where it is easily seen to hold. (See the proof of [1, Theorem 9], which is the only instance [1, Lemma 2] is used critically.)

Lemma 2.2. ([1, Lemma 1]) *Let M be an R -module and I an ideal of R such that $IM = 0$. Then M is reflexive as an R -module if and only if M is reflexive as an R/I -module.*

Proof. Since $E_{R/I} = \text{Hom}_R(R/I, E_R)$, the result follows readily by Hom-tensor adjunction. \square

Lemma 2.3. *Let $R = R_1 \times \cdots \times R_k$ be a product of Noetherian local rings. Let $M = M_1 \times \cdots \times M_k$ be an R -module. Then M is reflexive as an R -module if and only if M_i is reflexive as an R_i -module for all i .*

Proof. As finite sums and direct summands of reflexive modules are reflexive, it suffices to prove that M_i is reflexive as an R -module if and only if M_i is reflexive as an R_i -module for each i . But this follows immediately from Lemma 2.2. \square

Theorem 2.4. *Let S be a multiplicatively closed subset of R and M an R_S -module which is reflexive as an R -module. Then M is reflexive as an R_S -module.*

Proof. By Lemma 2.2, we may assume $\text{Ann}_R M = \text{Ann}_{R_S} M = 0$. Thus, R is semilocal and complete by Theorem 2.1. Hence, $R = R_1 \times \cdots \times R_k$ where each R_i is a complete local ring. Then $R_S = (R_1)_{S_1} \times \cdots \times (R_k)_{S_k}$ where S_i is the image of S under the canonical projection $R \rightarrow R_i$. Write $M = M_1 \times \cdots \times M_k$, where $M_i = R_i M$. As M is reflexive as an R -module, M_i is reflexive as an R_i -module for all i . Thus, it suffices to show that M_i is reflexive as an $(R_i)_{S_i}$ -module for all i . Hence, we may reduce the proof to the case (R, m) is a complete local ring with $\text{Ann}_R M = 0$ by passing to $R/\text{Ann}_R M$, if necessary. As M is reflexive as an R -module, we have by Theorem 2.1 that there exists an exact sequence

$$0 \rightarrow N \rightarrow M \rightarrow X \rightarrow 0$$

where N is a finitely generated R -module and X is an Artinian R -module. If $S \cap m = \emptyset$, then $R_S = R$ and there is nothing to prove. Otherwise, as $\text{Supp}_R X \subseteq \{m\}$, we have $X_S = 0$. Hence, $M \cong N_S$, a finitely generated R_S -module. To see that M is R_S -reflexive, it suffices to show that R_S is Artinian (hence semilocal and complete). Since $\text{Ann}_R N_S = \text{Ann}_R M = 0$, we have that $\text{Ann}_R N = 0$. Thus, $\dim R = \dim N$. Since M is an R_S -module and $S \cap m \neq \emptyset$, we have $H_m^i(M) \cong H_{mR_S}^i(M) = 0$ for all i . Further, as X is Artinian, $H_m^i(X) = 0$ for $i \geq 1$. Thus, from the long exact sequence on local cohomology, we conclude that $H_m^i(N) = 0$ for $i \geq 2$. Thus, $\dim R = \dim N \leq 1$, and hence, $\dim R_S = 0$. Consequently, R_S is Artinian, and M is a reflexive R_S -module. \square

To prove part (b) of Theorem 1.1, we will need the following result on Henselian local rings found in [2] (in which the authors credit it to F. Schmidt). As we need a slightly different version of this result than what is stated in [2] and the proof is short, we include it for the convenience of the reader:

Proposition 2.5. ([2, Satz 2.3.11]) *Let (R, m) be a local Henselian domain which is not a field and F the field of fractions of R . Let V be a discrete valuation ring with field of fractions F . Then $R \subseteq V$.*

Proof. Let k be the residue field of R and $a \in m$. As R is Henselian, for every positive integer n not divisible by the characteristic of k , the polynomial $x^n - (1 + a)$ has a root b in R . Let v be the valuation on F associated to V . Then $nv(b) = v(1 + a)$. If $v(a) < 0$ then $v(1 + a) < 0$ which implies $v(b) \leq -1$. Hence, $v(1 + a) \leq -n$. As n can be arbitrarily large, this leads to a contradiction. Hence, $v(a) \geq 0$ and $a \in V$. Thus, $m \subseteq V$. Now let $c \in R$ be arbitrary. Choose $d \in m, d \neq 0$. If $v(c) < 0$ then $v(c^\ell d) < 0$ for ℓ sufficiently large. But this contradicts that $c^\ell d \in m \subseteq V$ for every ℓ . Hence $v(c) \geq 0$ and $R \subseteq V$. \square

For a Noetherian ring R , let $\text{Min } R$ and $\text{Max } R$ denote the set of minimal and maximal primes of R , respectively. Let $\text{T}(R) = (\text{Spec } R \setminus \text{Min } R) \cup \text{Max } R$.

Lemma 2.6. *Let R be a Noetherian ring and $p \in \text{T}(R)$. If R_p is Henselian then the natural map $\varphi : R \rightarrow R_p$ is surjective; i.e., $R/\ker \varphi \cong R_p$.*

Proof. By replacing R with $R/\ker \varphi$, we may assume φ is injective. Then p contains every minimal prime of R . Let $u \in R, u \notin p$. It suffices to prove that the image of u in R/q is a unit for every minimal prime q of R . Hence, we may assume that R is a domain. (Note that $(R/q)_p = R_p/qR_p$ is still Henselian.) If R_p is a field, then, as $p \in \text{T}(R)$, we must have R is a field (as p must be both minimal and maximal in a domain). So certainly $u \notin p = (0)$ is a unit in R . Thus, we may assume R_p is not a field. Suppose u is not a unit in R . Then $u \in n$ for some maximal ideal n of R . Now, there exists a discrete valuation ring V with same field of fractions as R such that $m_V \cap R = n$ ([5, Theorem 6.3.3]). As R_p is Henselian, $R_p \subseteq V$ by Proposition 2.5. But as $u \notin p$, u is a unit in R_p , hence in V , contradicting $u \in n \subseteq m_V$. Thus, u is a unit in R and $R = R_p$. \square

Proposition 2.7. *Let R be a Noetherian ring and $S = R \setminus (p_1 \cup \dots \cup p_r)$ where $p_1, \dots, p_r \in \text{T}(R)$. Suppose R_S is complete with respect to its Jacobson radical. Then the natural map $\varphi : R \rightarrow R_S$ is surjective.*

Proof. First, we may assume that $p_j \not\subseteq \bigcup_{i \neq j} p_i$ for all j . Also, by passing to the ring $R/\ker \varphi$, we may assume φ is injective. (We note that if p_{i_1}, \dots, p_{i_t} are the ideals in the set $\{p_1, \dots, p_r\}$ containing $\ker \varphi$, it is easily seen that $(R/\ker \varphi)_S = (R/\ker \varphi)_T$ where $T = R \setminus (p_{i_1} \cup \dots \cup p_{i_t})$. Hence, we may assume each p_i contains $\ker \varphi$.) As R_S is semilocal and complete, the map $\psi : R_S \rightarrow R_{p_1} \times \dots \times R_{p_r}$ given by $\psi(u) = (\frac{u}{1}, \dots, \frac{u}{1})$ is an isomorphism. For each i , let $\rho_i : R \rightarrow R_{p_i}$ be the natural map. Since $R \rightarrow R_S$ is an injection, $\bigcap_i \ker \rho_i = (0)$. It suffices to prove that u is a unit in R for every $u \in S$. As R_{p_i} is complete, hence Henselian, we have that ρ_i is surjective for each i by Lemma 2.6. Thus, u is a unit in $R/\ker \rho_i$ for every i ; i.e., $(u) + \ker \rho_i = R$ for $i = 1, \dots, r$. Then $(u) = (u) + (\bigcap_i \ker \rho_i) = R$. Hence, u is a unit in R . \square

We now prove part (b) of the Theorem 1.1:

Theorem 2.8. *Let R be a Noetherian ring and M a reflexive R_S -module, where S is the complement in R of the union of finitely many elements of $\text{T}(R)$. Then M is reflexive as an R -module.*

Proof. We may assume $M \neq 0$. Let $S = R \setminus (p_1 \cup \cdots \cup p_r)$, where $p_1, \dots, p_r \in T(R)$. Let $I = \text{Ann}_R M$, whence $I_S = \text{Ann}_{R_S} M$. As in the proof of Proposition 2.7, we may assume each p_i contains I . Then by Lemma 2.2, we may reduce to the case $\text{Ann}_R M = \text{Ann}_{R_S} M = 0$. Note that this implies the natural map $R \rightarrow R_S$ is injective. As M is R_S -reflexive, R_S is complete with respect to its Jacobson radical by Theorem 2.1. By Proposition 2.7, we have that $R \cong R_S$ and hence M is R -reflexive. \square

3. EXAMPLES

The following examples show that $\text{Hom}_R(R_S, E_R)$ need not be the minimal injective cogenerator for the category of R_S -modules, contrary to what is stated in the proof of [1, Lemma 2]:

Example 3.1. Let (R, m) be a local ring of dimension at least two and p any prime which is not maximal or minimal. By [3, Lemma 4.1], every element of $\text{Spec } R_p$ is an associated prime of the R_p -module $\text{Hom}_R(R_p, E_R)$. In particular, $\text{Hom}_R(R_p, E_R) \not\cong E_{R_p}$.

Example 3.2. ([3, p. 127]) Let R be a local domain such that the completion of R has a nonminimal prime contracting to (0) in R . Let Q be the field of fractions of R . Then $\text{Hom}_R(Q, E_R)$ is not Artinian.

Example 3.3. Let R be a Noetherian domain which is not local. Let $m \neq n$ be maximal ideals of R . By a slight modification of the proof of [3, Lemma 4.1], one obtains that (0) is an associated prime of $\text{Hom}_R(R_m, E_R(R/n))$, which is a direct summand of $\text{Hom}_R(R_m, E_R)$. Hence, $\text{Hom}_R(R_m, E_R) \not\cong E_{R_m}$.

We now show that the converse to part (a) of Theorem 1.1 does not hold in general. Let R be a domain and Q its field of fractions. Of course, Q is reflexive as a $Q = R_{(0)}$ -module. But as the following theorem shows, Q is rarely a reflexive R -module.

Proposition 3.4. *Let R be a Noetherian domain and Q the field of fractions of R . Then Q is a reflexive R -module if and only if R is a complete local domain of dimension at most one.*

Proof. We first suppose R is a one-dimensional complete local domain with maximal ideal m . Let $E = E_R(R/m)$. By [4, Theorem 2.5], $\text{Hom}_R(Q, E) \cong Q$. Since the evaluation map of the Matlis double dual is always injective, we obtain that $Q \rightarrow \text{Hom}_R(\text{Hom}_R(Q, E), E)$ is an isomorphism.

Conversely, suppose Q is a reflexive R -module. By Theorem 2.1, R is a complete semilocal domain, hence local. It suffices to prove that $\dim R \leq 1$. Again by Theorem 2.1, there exists a finitely generated R -submodule N of Q such that Q/N is Artinian. Since $\text{Ann}_R N = 0$, $\dim R = \dim N$. Thus, it suffices to prove that $H_m^i(N) = 0$ for $i \geq 2$. But this follows readily from the facts that $H_m^i(Q) = 0$ for all i and $H_m^i(Q/N) = 0$ for $i \geq 1$ (as Q/N is Artinian). \square

Acknowledgments: The authors would like to thank Peder Thompson for many helpful discussions on this topic. They are also very grateful to Bill Heinzer for pointing out the existence of Proposition 2.5.

REFERENCES

1. R. Belshoff, E. Enochs, and J. García-Rozas, Generalized Matlis duality. Proc. Amer. Math. Soc. **128** (1999), no. 5, 1307-1312.
2. R. Berger, R. Kiehl, E. Kunz, and H.-J. Nastold, Differentialrechnung in der analytischen Geometrie. Lecture Notes in Mathematics **38**, Springer-Verlag, Berlin-New York, 1967.
3. L. Melkersson and P. Schenzel, The co-localization of an Artinian module. Proc. Edinburgh Math. Soc. **38** (1995), 121–131.
4. P. Schenzel, A note on the Matlis dual of a certain injective hull. J. Pure Appl. Algebra **219** (2015), no. 3, 666-671.
5. I. Swanson and C. Huneke, Integral closures of ideals, rings and modules. London Mathematical Society Lecture Note Series **336**, Cambridge University Press, Cambridge, 2006.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEBRASKA-LINCOLN, LINCOLN, NE 68588-0130
E-mail address: `ddailey2@math.unl.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEBRASKA-LINCOLN, LINCOLN, NE 68588-0130
E-mail address: `tmarley1@unl.edu`