

RIGIDITY OF EXT AND TOR WITH COEFFICIENTS IN RESIDUE FIELDS OF A COMMUTATIVE NOETHERIAN RING

LARS WINTHER CHRISTENSEN, SRIKANTH B. IYENGAR, AND THOMAS MARLEY

ABSTRACT. Let \mathfrak{p} be a prime ideal in a commutative noetherian ring R . It is proved that if an R -module M satisfies $\mathrm{Tor}_n^R(k(\mathfrak{p}), M) = 0$ for some $n \geq \dim R_{\mathfrak{p}}$, where $k(\mathfrak{p})$ is the residue field at \mathfrak{p} , then $\mathrm{Tor}_i^R(k(\mathfrak{p}), M) = 0$ holds for all $i \geq n$. Similar rigidity results concerning $\mathrm{Ext}_R^*(k(\mathfrak{p}), M)$ are proved, and applications to the theory of homological dimensions are explored.

1. INTRODUCTION

Flatness and injectivity of modules over a commutative ring R are characterized by vanishing of (co)homological functors and such vanishing can be verified by testing on cyclic R -modules. We discuss the flat case first, and in mildly greater generality: For any R -module M and integer $n \geq 0$, one has $\mathrm{flat\,dim}_R M < n$ if and only if $\mathrm{Tor}_n^R(R/\mathfrak{a}, M) = 0$ holds for all ideals $\mathfrak{a} \subseteq R$. When R is noetherian, and in this paper we assume that it is, it suffices to test on modules R/\mathfrak{p} where \mathfrak{p} varies over the prime ideals in R .

If R is local with unique maximal ideal \mathfrak{m} , and M is finitely generated, then it is sufficient to consider one cyclic module, namely the residue field $k := R/\mathfrak{m}$. Even if R is not local and M is not finitely generated, finiteness of $\mathrm{flat\,dim}_R M$ is characterized by vanishing of Tor with coefficients in fields, the residue fields $k(\mathfrak{p}) := R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}}$ to be specific. While vanishing of $\mathrm{Tor}_*^R(k(\mathfrak{p}), M)$ for any one particular residue field does not imply that $\mathrm{flat\,dim}_R M$ is finite, one may still ask if vanishing of a single group $\mathrm{Tor}_n^R(k(\mathfrak{p}), M)$ implies vanishing of all higher groups, a phenomenon known as *rigidity*. While this does not hold in general (cf. Example 4.2), we prove that it does hold if n is sufficiently large; see Theorem 4.1 for the proof.

Theorem 1.1. *Let \mathfrak{p} be a prime ideal in a commutative noetherian ring R and let M be an R -module. If one has $\mathrm{Tor}_n^R(k(\mathfrak{p}), M) = 0$ for some integer $n \geq \dim R_{\mathfrak{p}}$, then $\mathrm{Tor}_i^R(k(\mathfrak{p}), M) = 0$ holds for all $i \geq n$.*

As $\mathrm{flat\,dim}_R M < n$ holds if and only if one has $\mathrm{Tor}_i^R(k(\mathfrak{p}), M) = 0$ for all primes \mathfrak{p} and all $i \geq n$, Theorem 1.1 provides for an improvement of existing characterizations of modules of finite flat dimension.

In parallel to the flat case, the injective dimension of an R -module M is less than n if $\mathrm{Ext}_R^n(R/\mathfrak{p}, M) = 0$ for every prime ideal \mathfrak{p} . Moreover the injective dimension can be detected by vanishing *locally* of cohomology with coefficients in residue

Date: 27 July 2017.

2010 Mathematics Subject Classification. 13D07; 13D05.

Key words and phrases. Ext, Tor, rigidity, flat dimension, injective dimension.

L.W.C. was partly supported by NSA grant H98230-14-0140 and Simons Foundation collaboration grant 428308; S.B.I. was partly supported by NSF grant DMS-1503044.

fields. That is, $\text{inj dim}_R M < n$ holds if and only if one has $\text{Ext}_{R_{\mathfrak{p}}}^n(k(\mathfrak{p}), M_{\mathfrak{p}}) = 0$ for all primes \mathfrak{p} . By the standard isomorphisms $\text{Tor}_*^R(k(\mathfrak{p}), M) \cong \text{Tor}_*^{R_{\mathfrak{p}}}(k(\mathfrak{p}), M_{\mathfrak{p}})$ there is no local/global distinction for Tor vanishing. The following consequence of Proposition 3.2 is, therefore, a perfect parallel to Theorem 1.1.

Theorem 1.2. *Let \mathfrak{p} be a prime ideal in a commutative noetherian ring R and let M be an R -module. If one has $\text{Ext}_{R_{\mathfrak{p}}}^n(k(\mathfrak{p}), M_{\mathfrak{p}}) = 0$ for some integer $n \geq \dim R_{\mathfrak{p}}$, then $\text{Ext}_{R_{\mathfrak{p}}}^i(k(\mathfrak{p}), M_{\mathfrak{p}}) = 0$ holds for all $i \geq n$.*

In contrast to the situation for Tor, the cohomology groups $\text{Ext}_{R_{\mathfrak{p}}}^*(k(\mathfrak{p}), M_{\mathfrak{p}})$ and $\text{Ext}_R^*(k(\mathfrak{p}), M)$ can be quite different, and it was only proved recently, in [6, Theorem 1.1], that the injective dimension of an R -module can be detected by vanishing *globally* of cohomology with coefficients in residue fields. That is, $\text{inj dim}_R M < n$ holds if and only if one has $\text{Ext}_R^i(k(\mathfrak{p}), M) = 0$ for all $i \geq n$ and all primes \mathfrak{p} . One advantage of this global vanishing criterion is that it also applies to complexes of modules; per Example 6.3 the local vanishing criterion does not. For the proof of the following rigidity result for $\text{Ext}_R^*(k(\mathfrak{p}), M)$, see Remark 5.8.

Theorem 1.3. *Let \mathfrak{p} be a prime ideal in a commutative noetherian ring R and let M be an R -module. If one has $\text{Ext}_R^n(k(\mathfrak{p}), M) = 0$ for some integer $n \geq 2 \dim R$, then $\text{Ext}_R^i(k(\mathfrak{p}), M) = 0$ holds for all $i \geq n$.*

The case when \mathfrak{p} is the maximal ideal of a local ring merits comment, for the bound on n in Theorems 1.2 and 1.3 differs by a factor of 2. The proof shows that it is sufficient to require $n \geq \dim R_{\mathfrak{p}} + \text{proj dim}_R R_{\mathfrak{p}}$ in Theorem 1.3, and that aligns the two bounds in this special case. For a general prime \mathfrak{p} however the number $\text{proj dim}_R R_{\mathfrak{p}}$ may depend on the Continuum Hypothesis; see Osofsky [14].

In this introduction, we have focused on results that deal with rigidity of the Tor and Ext functors. In the text, we also establish results that track where vanishing of these functors starts, when indeed they vanish eventually.

* * *

Throughout R will be a commutative noetherian ring. Background material on homological invariants and local (co)homology is recalled in Section 2. Rigidity results for Ext and Tor over local rings are proved in Section 3, and applications to homological dimensions follow in Sections 4–5. The final section explores, by way of examples, the complicated nature of injective dimension of unbounded complexes.

2. LOCAL HOMOLOGY AND LOCAL COHOMOLOGY

Our standard reference for definitions and constructions involving complexes is [2]. We will be dealing with graded modules whose natural grading is the upper one and also those whose natural grading is the lower one. Therefore, we set

$$\inf H_*(M) := \inf\{i \mid H_i(M) \neq 0\} \quad \text{and} \quad \inf H^*(M) := \inf\{i \mid H^i(M) \neq 0\}$$

for an R -complex M , and analogously we define $\sup H_*(M)$ and $\sup H^*(M)$. We often work in the derived category of R -modules, and write \simeq for isomorphisms there. A morphism between complexes is a *quasi-isomorphism* if it is an isomorphism in homology; that is to say, if it becomes an isomorphism in the derived category.

The next paragraphs summarize the definitions and basic results on local cohomology and local homology, following [1, 13].

Local (co)homology. Let R be a commutative noetherian ring and \mathfrak{a} an ideal in R . The right derived functor of the \mathfrak{a} -torsion functor $\Gamma_{\mathfrak{a}}$ is denoted $\mathbf{R}\Gamma_{\mathfrak{a}}$, and the *local cohomology* supported on \mathfrak{a} of an R -complex M is the graded module

$$\mathbf{H}_{\mathfrak{a}}^*(M) := \mathbf{H}^*(\mathbf{R}\Gamma_{\mathfrak{a}}(M)).$$

There is a natural morphism $\mathbf{R}\Gamma_{\mathfrak{a}}(M) \rightarrow M$ in the derived category; M is said to be *derived \mathfrak{a} -torsion* when this map is an isomorphism. This is equivalent to the condition that $\mathbf{H}^*(M)$ is degreewise \mathfrak{a} -torsion; see [7, Proposition 6.12].

The left derived functor of the \mathfrak{a} -adic completion functor $\Lambda^{\mathfrak{a}}$ is denoted $\mathbf{L}\Lambda^{\mathfrak{a}}$ and the *local homology* of M supported on \mathfrak{a} is the graded module

$$\mathbf{H}_*^{\mathfrak{a}}(M) := \mathbf{H}_*(\mathbf{L}\Lambda^{\mathfrak{a}}(M)).$$

There is a natural morphism $M \rightarrow \mathbf{L}\Lambda^{\mathfrak{a}}(M)$ in the derived category and we say M is *derived \mathfrak{a} -complete* when this map is an isomorphism. This is equivalent to the condition that for each i , the natural map $\mathbf{H}^i(M) \rightarrow \mathbf{H}_0^{\mathfrak{a}}(\mathbf{H}^i(M))$ is an isomorphism; see [7, Proposition 6.15].

The morphisms $\mathbf{R}\Gamma_{\mathfrak{a}}(M) \rightarrow M$ and $M \rightarrow \mathbf{L}\Lambda^{\mathfrak{a}}(M)$ induce isomorphisms

$$(2.1) \quad \begin{aligned} \mathrm{Ext}_R^*(R/\mathfrak{a}, M) &\cong \mathrm{Ext}_R^*(R/\mathfrak{a}, \mathbf{R}\Gamma_{\mathfrak{a}}M) \quad \text{and} \\ \mathrm{Tor}_*^R(R/\mathfrak{a}, M) &\cong \mathrm{Tor}_*^R(R/\mathfrak{a}, \mathbf{L}\Lambda^{\mathfrak{a}}M). \end{aligned}$$

Indeed, the first one holds because the functor $\mathbf{R}\Gamma_{\mathfrak{a}}$ is right adjoint to the inclusion of the \mathfrak{a} -torsion complexes (that is to say, complexes whose cohomology is \mathfrak{a} -torsion) into the derived category of R ; see [13, Proposition 3.2.2]. Thus one has

$$(2.2) \quad \mathrm{RHom}(R/\mathfrak{a}, \mathbf{R}\Gamma_{\mathfrak{a}}M) \xrightarrow{\cong} \mathrm{RHom}(R/\mathfrak{a}, M),$$

and this gives the first isomorphism. As to the second isomorphism, consider the commutative square in the derived category

$$\begin{array}{ccc} \mathbf{R}\Gamma_{\mathfrak{a}}M & \longrightarrow & M \\ \cong \downarrow & & \downarrow \\ \mathbf{R}\Gamma_{\mathfrak{a}}\mathbf{L}\Lambda^{\mathfrak{a}}M & \longrightarrow & \mathbf{L}\Lambda^{\mathfrak{a}}M \end{array}$$

induced by the vertical morphism on right. The isomorphism on the left is part of Greenlees–May duality; see, for example, [1, Corollary (5.1.1)]. Applying the functor $R/\mathfrak{a} \otimes_R^{\mathbf{L}}(-)$ yields the commutative square

$$\begin{array}{ccc} R/\mathfrak{a} \otimes_R^{\mathbf{L}} \mathbf{R}\Gamma_{\mathfrak{a}}M & \xrightarrow{\cong} & R/\mathfrak{a} \otimes_R^{\mathbf{L}} M \\ \cong \downarrow & & \downarrow \\ R/\mathfrak{a} \otimes_R^{\mathbf{L}} \mathbf{R}\Gamma_{\mathfrak{a}}\mathbf{L}\Lambda^{\mathfrak{a}}M & \xrightarrow{\cong} & R/\mathfrak{a} \otimes_R^{\mathbf{L}} \mathbf{L}\Lambda^{\mathfrak{a}}M \end{array}$$

The horizontal maps are isomorphisms by (2.2) and [7, Proposition 6.5]. Thus the vertical map on the right is also an isomorphism; in homology, this is the desired isomorphism.

Let $C(\mathfrak{a})$ denote the Čech complex on a set of elements that generate \mathfrak{a} . The values of the functors $\mathbf{L}\Lambda^{\mathfrak{a}}$ and $\mathbf{R}\Gamma_{\mathfrak{a}}$ on an R -complex M can then be computed as

$$\mathbf{L}\Lambda^{\mathfrak{a}}(M) = \mathrm{RHom}_R(C(\mathfrak{a}), M) \quad \text{and} \quad \mathbf{R}\Gamma_{\mathfrak{a}}(M) = C(\mathfrak{a}) \otimes_R^{\mathbf{L}} M.$$

See for example [1, Theorem (0.3) and Lemma (3.1.1)].

Depth and width. In the remainder of this section (R, \mathfrak{m}, k) will be a local ring. This means that R is a commutative noetherian ring with unique maximal ideal \mathfrak{m} and residue field $k := R/\mathfrak{m}$.

The *depth* and *width* of an R -complex M are defined as follows:

$$\text{depth}_R M = \inf \text{Ext}_R^*(k, M) \quad \text{and} \quad \text{width}_R M = \inf \text{Tor}_*^R(k, M).$$

One has $\text{depth}_R M \geq \inf H^*(M)$ and if $i = \inf H^*(M)$ is finite, then equality holds if and only if $\text{Hom}_R(k, H^i(M)) \neq 0$. Similarly, one has $\text{width}_R M \geq \inf H_*(M)$ and if $j = \inf H_*(M)$ is finite, then equality holds if and only if $k \otimes_R H_j(M) \neq 0$.

If $\text{flat dim}_R M$ is finite, then one has an equality

$$(2.3) \quad \text{depth}_R M = \text{depth } R - \sup \text{Tor}_*^R(k, M).$$

This is an immediate consequence of [2, Lemma 4.4(F)]. For finitely generated modules it is the Auslander–Buchsbaum Formula.

Similarly, if $\text{inj dim}_R M$ is finite, then one has

$$(2.4) \quad \text{width}_R M = \text{depth } R - \sup \text{Ext}_R^*(k, M).$$

This is a consequence of [2, Lemma 4.4(I)]. For finitely generated modules the equality above yields Bass' formula $\text{inj dim}_R M = \text{depth } R$.

From [9, Definitions 2.3 and 4.3] one gets that the depth and width of an R -complex can be detected by vanishing of local (co)homology:

$$\text{depth}_R M = \inf H_{\mathfrak{m}}^*(M) \quad \text{and} \quad \text{width}_R M = \inf H_{\mathfrak{m}}^{\mathfrak{m}}(M).$$

Combining this with (2.1) and the isomorphisms

$$R\Gamma_{\mathfrak{m}}\mathbf{L}A^{\mathfrak{m}}(M) \simeq R\Gamma_{\mathfrak{m}}(M) \quad \text{and} \quad \mathbf{L}A^{\mathfrak{m}}R\Gamma_{\mathfrak{m}}(M) \simeq \mathbf{L}A^{\mathfrak{m}}(M)$$

from [1, Corollary (5.1.1)] one gets equalities

$$(2.5) \quad \text{depth}_R R\Gamma_{\mathfrak{m}}M = \text{depth}_R M = \text{depth}_R \mathbf{L}A^{\mathfrak{m}}M$$

$$(2.6) \quad \text{width}_R R\Gamma_{\mathfrak{m}}M = \text{width}_R M = \text{width}_R \mathbf{L}A^{\mathfrak{m}}M.$$

For later use, we note that for each R -complex M there are inequalities

$$(2.7) \quad \sup H_{\mathfrak{m}}^*(M) \leq \dim R + \sup H^*(M) \quad \text{and} \quad \sup H_{\mathfrak{m}}^*(M) \leq \sup \text{Ext}_R^*(k, M).$$

The first is immediate as one has $H_{\mathfrak{m}}^*(M) = H^*(C(\mathfrak{m}) \otimes_R^{\mathbf{L}} M)$ where $C(\mathfrak{m})$ is the Čech complex on a system of parameters for R ; the second is immediate once one recalls the isomorphism $H_{\mathfrak{m}}^*(M) \cong \varinjlim_i \text{Ext}_R^*(R/\mathfrak{m}^i, M)$.

The next result is a direct extension of [16, Proposition 2.1] by A.-M. Simon. Concerning the last assertion: $\text{Tor}_i^R(R, M) = H_i(M) \neq 0$, so n cannot equal $\sup H_*(M)$, unless both are infinite. However, for later applications it is convenient to have the statement in this form.

Lemma 2.1. *Let M be a derived \mathfrak{a} -complete R -complex with $\inf H_*(M) > -\infty$ and n an integer. If $\text{Tor}_n^R(R/\mathfrak{p}, M) = 0$ all prime ideals $\mathfrak{p} \supseteq \mathfrak{a}$, then $\text{Tor}_n^R(-, M) = 0$.*

When in addition $n \geq \sup H_(M)$, one has $\text{flat dim}_R M \leq n - 1$.*

Proof. First we claim that for any finitely generated R -module L and integer i , if $\mathfrak{a} \text{Tor}_i^R(L, M) = \text{Tor}_i^R(L, M)$, then $\text{Tor}_i^R(L, M) = 0$. Indeed, let F be a free resolution of L with each F_i finitely generated and equal to zero for $i < 0$. Let G be a semi-flat resolution of M with $G_i = 0$ for $i \ll 0$; this is possible as $\inf H_*(M)$

is finite. Since M is derived \mathfrak{a} -complete, the complex $\Lambda^{\mathfrak{a}}G$, which computes $L\Lambda^{\mathfrak{a}}M$, is quasi-isomorphic to M . Thus

$$\mathrm{Tor}_i^R(L, M) = \mathrm{H}_i(F \otimes_R \Lambda^{\mathfrak{a}}G).$$

Note that $F \otimes_R \Lambda^{\mathfrak{a}}G$ is a complex of \mathfrak{a} -adically complete R -modules; this is where we need that each F_i is finitely generated and that F_i and G_i are zero for $i \ll 0$. It remains to apply [16, Proposition 1.4].

For the stated result, it suffices to prove that the set

$$\{\mathfrak{b} \subset R \text{ an ideal} \mid \mathrm{Tor}_n^R(R/\mathfrak{b}, M) \neq 0\}$$

is empty. Suppose it is not. Pick a maximal element; say, \mathfrak{q} . We claim that this is a prime ideal. The argument is standard (see, for example, [12, 2.4]) and goes as follows: If it is not, let \mathfrak{q}' be an associated prime ideal of R/\mathfrak{q} , and $x \in R$ an element such that $\mathfrak{q}' = \{r \in R \mid xr \in \mathfrak{q}\}$. Then yields an exact sequence

$$0 \longrightarrow R/\mathfrak{q}' \xrightarrow{1 \mapsto x} R/\mathfrak{q} \longrightarrow R/((x) + \mathfrak{q}) \longrightarrow 0$$

Since both \mathfrak{q}' and $(x) + \mathfrak{q}$ strictly contain \mathfrak{q} , one obtains that

$$\mathrm{Tor}_n^R(R/\mathfrak{q}', M) = 0 = \mathrm{Tor}_n^R(R/((x) + \mathfrak{q}), M),$$

and hence $\mathrm{Tor}_n^R(R/\mathfrak{q}, M) = 0$, contradicting the choice of \mathfrak{q} . Thus \mathfrak{q} is prime.

By hypothesis, \mathfrak{q} does not contain \mathfrak{a} , so choose an element a in \mathfrak{a} but not in \mathfrak{q} and consider the exact sequence

$$0 \longrightarrow R/\mathfrak{q} \xrightarrow{a} R/\mathfrak{q} \longrightarrow R/((a) + \mathfrak{q}) \longrightarrow 0$$

Noting that $\mathrm{Tor}_n^R(R/((a) + \mathfrak{q}), M) = 0$ by the choice of \mathfrak{q} , it follows from the exact sequence above that the map $\mathrm{Tor}_n^R(R/\mathfrak{q}, M) \xrightarrow{a} \mathrm{Tor}_n^R(R/\mathfrak{q}, M)$ is surjective. By the claim in the first paragraph, this implies that $\mathrm{Tor}_n^R(R/\mathfrak{q}, M) = 0$, which is a contradiction. \square

3. LOCAL RINGS

In this section (R, \mathfrak{m}, k) is a local ring. Note from (2.7) that in the next statement n cannot equal $\sup \mathrm{H}^*(M)$, but, as with Lemma 2.1, this formulation is convenient for later applications.

Lemma 3.1. *Let M be a derived \mathfrak{m} -torsion R -complex with $\inf \mathrm{H}^*(M) > -\infty$. If one has $\mathrm{Ext}_R^n(k, M) = 0$ for some integer $n \geq \sup \mathrm{H}^*(M)$, then $\mathrm{inj\,dim}_R M \leq n - 1$.*

Proof. Let I be the minimal semi-injective resolution of M . One has

$$\mathrm{Ext}_R^n(k, M) = \mathrm{H}^n(\mathrm{Hom}_R(k, I)) = \mathrm{Hom}_R(k, I^n).$$

As M is derived \mathfrak{m} -torsion, each module I^i is a direct sum of copies of the injective envelope of k , so $\mathrm{Ext}_R^n(k, M) = 0$ implies $I^n = 0$. It follows from the assumption on n and minimality of I that $I^i = 0$ holds for all $i \geq n$; in particular, one has $\mathrm{inj\,dim}_R M \leq n - 1$. \square

The result below extends (2.4); its proof would be significantly shorter under the additional hypothesis that $\inf \mathrm{H}_{\mathfrak{m}}^*(M)$ is finite.

Proposition 3.2. *Let M be an R -complex. If $\mathrm{Ext}_R^n(k, M) = 0$ holds for some integer $n \geq \sup \mathrm{H}_{\mathfrak{m}}^*(M)$, then one has $\mathrm{Ext}_R^i(k, M) = 0$ for all $i \geq n$ and*

$$\sup \mathrm{Ext}_R^*(k, M) = \mathrm{depth} R - \mathrm{width}_R M.$$

Proof. Let J be the minimal semi-injective resolution of M and set $I := \Gamma_{\mathfrak{m}}(J)$. For every integer i one has

$$\mathrm{H}_{\mathfrak{m}}^i(M) = \mathrm{H}^i(I) \quad \text{and} \quad \mathrm{Ext}_R^i(k, M) = \mathrm{Hom}_R(k, J^i) = \mathrm{Hom}_R(k, I^i).$$

Let d be any integer with $n \geq d \geq \sup \mathrm{H}_{\mathfrak{m}}^*(M)$ and set $Z := Z^d(I)$, the submodule of cycles in degree d . Vanishing of $\mathrm{H}_{\mathfrak{m}}^i(M)$ and the identifications above yield

$$\mathrm{Ext}_R^i(k, M) \cong \mathrm{Ext}_R^{i-d}(k, Z) \quad \text{for all } i \geq d.$$

In particular, one has $\mathrm{Ext}_R^{n-d}(k, Z) = 0$. Since Z is an \mathfrak{m} -torsion R -module, Lemma 3.1 and the isomorphisms above yield $\mathrm{Ext}_R^i(k, M) = 0$ for all $i \geq n$.

It remains to verify the claim about the supremum of $\mathrm{Ext}_R^*(k, M)$. To this end, let K be the Koszul complex on a minimal set of generators for \mathfrak{m} . It follows from [9, Definition 2.3] that one has the second equivalence below

$$\begin{aligned} \sup \mathrm{Ext}_R^*(k, M) = -\infty &\iff \mathrm{Ext}_R^*(k, M) = 0 \\ &\iff \mathrm{H}^*(K \otimes_R M) = 0 \\ &\iff \mathrm{width}_R M = \infty. \end{aligned}$$

The first one is by definition while the last one is by [9, Theorem 4.1]. We may thus assume that $s := \sup \mathrm{Ext}_R^*(k, M)$ and $w := \sup \mathrm{H}^*(M \otimes_R K)$ are integers. Because K is a bounded complex of finitely generated free R -modules, there is a quasi-isomorphism

$$\mathrm{RHom}_R(k, M) \otimes_R^{\mathbb{L}} K \simeq \mathrm{RHom}(k, M \otimes_R K).$$

From this and the fact that $\mathrm{Ext}_R^*(k, M)$ is a graded k -vector space, it follows that

$$s = \sup \mathrm{H}^*(\mathrm{RHom}_R(k, M) \otimes_R^{\mathbb{L}} K) = \sup \mathrm{Ext}_R^*(k, M \otimes_R K).$$

Let E be the minimal injective resolution of $M \otimes_R K$. Since $M \otimes_R K$ is derived \mathfrak{m} -torsion, one has $\Gamma_{\mathfrak{m}} E = E$. From $\mathrm{Ext}_R^*(k, M \otimes_R K) = \mathrm{Hom}_R^*(k, E)$ it thus follows that $E^s \neq 0$ and $E^i = 0$ for all $i > s$. On the other hand, as $w = \sup \mathrm{H}^*(E)$ the complex $E^{\geq w}$ is the minimal injective resolution of the module $W := Z^w(E)$ of cycles in degree w , so one has $\mathrm{inj dim}_R W = s - w$.

It remains to show that $\mathrm{inj dim}_R W = \mathrm{depth} R$. Evidently one has $\mathrm{inj dim}_R W = \sup \mathrm{Ext}_R^*(k, W)$, so by (2.4) it suffices to show that W has width 0, that is to say, that $k \otimes_R W \neq 0$. But this is clear because $\mathrm{H}^w(E)$ is nonzero and annihilated by \mathfrak{m} , whence $\mathfrak{m}W \subseteq \mathrm{B}^w(E) \subsetneq W$. \square

Proposition 3.3. *Let M be an R -complex with $\mathrm{Tor}_n^R(k, M) = 0$ for some integer n , and assume that one of the following conditions is satisfied:*

- (1) $n \geq \sup \mathrm{H}_{\mathfrak{m}}^*(M)$ and $\inf \mathrm{H}_*(M) > -\infty$;
- (2) $n \geq \sup \mathrm{H}_{\mathfrak{m}}^*(\mathrm{Hom}_R(M, E(k)))$ where $E(k)$ is the injective envelope of k .

One then has $\mathrm{Tor}_i^R(k, M) = 0$ for all $i \geq n$ and

$$\sup \mathrm{Tor}_*^R(k, M) = \mathrm{depth} R - \mathrm{depth}_R M.$$

Proof. Assume first that (1) is satisfied. From (2.1) one gets $\mathrm{Tor}_n^R(k, \mathrm{L}A^{\mathfrak{m}}M) = 0$. The complex $\mathrm{L}A^{\mathfrak{m}}M$ is \mathfrak{m} -adically complete and

$$\inf \mathrm{H}_*(\mathrm{L}A^{\mathfrak{m}}M) \geq \inf \mathrm{H}_*(M) > -\infty.$$

Thus Lemma 2.1 applies and yields that $\text{flat dim}_R(LA^m M)$ is at most $n - 1$. Now (2.1) yields $\text{Tor}_i^R(k, M) = 0$ for all $i \geq n$, and then (2.3) and (2.5) yield the second and third equalities below

$$\begin{aligned} \sup \text{Tor}_*^R(k, M) &= \sup \text{Tor}_*^R(k, LA^m M) \\ &= \text{depth } R - \text{depth}_R(LA^m M) \\ &= \text{depth } R - \text{depth}_R M. \end{aligned}$$

Assume now that the hypothesis in (2) holds, and set $(-)^{\vee} := \text{Hom}_R(-, E(k))$. For each integer i , there is an isomorphism

$$\text{Tor}_i^R(k, M)^{\vee} \cong \text{Ext}_R^i(k, M^{\vee}).$$

The hypothesis and Proposition 3.2 yield $\text{Tor}_i^R(k, M) = 0$ for all $i \geq n$ and

$$\sup \text{Tor}_*^R(k, M) = \sup \text{Ext}_R^*(k, M^{\vee}) = \text{depth } R - \text{width}_R M^{\vee}.$$

Finally one has $\text{width}_R M^{\vee} = \text{depth}_R M$; see [9, Proposition 4.4]. \square

The final result of this section fleshes out a remark made by Fossum, Foxby, Griffith, and Reiten at the end of Section 1 in [8]. They phrase it as statement about nonvanishing: If M is an R -module and $\text{Ext}_R^n(k, M)$ is nonzero for some $n \geq \text{depth } R + 1$ then one has $\text{Ext}_R^i(k, M) \neq 0$ for all $i \geq n$. The formulation below makes for an easier comparison with Proposition 3.2.

Proposition 3.4. *Let M be an R -complex. If $\text{Ext}_R^n(k, M) = 0$ holds for some integer $n \geq \sup H^*(M) + \text{depth } R + 1$, then one has*

$$\text{Ext}_R^i(k, M) = 0 \quad \text{for every } i \text{ in the range } \sup H^*(M) + \text{depth } R + 1 \leq i \leq n.$$

Proof. We may assume that $n > \sup H^*(M) + \text{depth } R + 1$ holds. It suffices to verify that when $\text{Ext}_R^n(k, M)$ is zero, so is $\text{Ext}_R^{n-1}(k, M)$.

Let \mathbf{x} be a maximal regular sequence in R , set $S := R/(\mathbf{x})$ and $\mathfrak{n} := \mathfrak{m}/(\mathbf{x})$. Thus, (S, \mathfrak{n}, k) is a local ring of depth 0; in particular, $(0 : \mathfrak{n})$, the socle of S , is nonzero. Thus, there exists a positive integer, say s , such that $(0 : \mathfrak{n})$ is contained in \mathfrak{n}^s but not in \mathfrak{n}^{s+1} . Said otherwise, the composite of canonical maps

$$(0 : \mathfrak{n}) \longrightarrow \mathfrak{n}^s \longrightarrow \mathfrak{n}^s / \mathfrak{n}^{s+1}$$

is nonzero. Since the source and the target are k -vector spaces, this implies that k is a direct summand of \mathfrak{n}^s . It thus suffices to verify that $\text{Ext}_R^{n-1}(\mathfrak{n}^s, M) = 0$.

The Koszul complex on \mathbf{x} is a minimal free resolution of S over R , so one has $\text{proj dim}_R S = \text{depth } R$ and hence

$$\text{Ext}_R^j(S, M) = 0 \quad \text{for } j \geq \text{depth } R + 1 + \sup H^*(M).$$

Given this, the exact sequence

$$0 \longrightarrow \mathfrak{n}^s \longrightarrow S \longrightarrow S/\mathfrak{n}^s \longrightarrow 0$$

yields an isomorphism

$$\text{Ext}_R^{n-1}(\mathfrak{n}^s, M) \cong \text{Ext}_R^n(S/\mathfrak{n}^s, M).$$

Since the length of S/\mathfrak{n}^s is finite, $\text{Ext}_R^n(k, M) = 0$ implies $\text{Ext}_R^n(S/\mathfrak{n}^s, M) = 0$, and hence the isomorphism above yields $\text{Ext}_R^{n-1}(\mathfrak{n}^s, M) = 0$, as desired. \square

4. FLAT DIMENSION

Let R be a commutative noetherian ring, M be an R -complex, and n be an integer. Avramov and Foxby [2, Proposition 5.3.F] prove that $\text{flat dim}_R M < n$ holds if and only if one has $\text{Tor}_i^{R_{\mathfrak{p}}}(k(\mathfrak{p}), M_{\mathfrak{p}}) = 0$ for every prime \mathfrak{p} in R and all $i \geq n$. That is,

$$(4.1) \quad \text{flat dim}_R M = \sup\{i \in \mathbb{Z} \mid \text{Tor}_i^{R_{\mathfrak{p}}}(k(\mathfrak{p}), M_{\mathfrak{p}}) \neq 0 \text{ for some } \mathfrak{p} \in \text{Spec } R\}.$$

By way of the isomorphisms

$$(4.2) \quad \text{Tor}_i^{R_{\mathfrak{p}}}(k(\mathfrak{p}), M_{\mathfrak{p}}) \cong \text{Tor}_i^R(k(\mathfrak{p}), M)$$

this result compares—or may be it is the other way around—to [6, Theorem 1.1]; see (5.1). Combining (4.1) and (4.2) with (2.3) one gets

$$(4.3) \quad \text{flat dim}_R M = \sup_{\mathfrak{p} \in \text{Spec } R} \{\text{depth } R_{\mathfrak{p}} - \text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}\}$$

for every R -complex M of finite flat dimension. For modules of finite flat dimension this equality is known from work of Chouinard [5].

For rings of finite Krull dimension, the next theorem, which contains Theorem 1.1, represents a significant strengthening of (4.1).

Theorem 4.1. *Let R be a commutative noetherian ring and M be an R -complex.*

If for a prime ideal \mathfrak{p} and $n \geq \dim R_{\mathfrak{p}} + \sup H_(M)$ one has $\text{Tor}_n^R(k(\mathfrak{p}), M) = 0$, then*

$$\sup \text{Tor}_*^R(k(\mathfrak{p}), M) = \text{depth } R_{\mathfrak{p}} - \text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} < n.$$

In particular, if there exists an integer $n \geq \dim R + \sup H_(M)$ such that*

$$\text{Tor}_n^R(k(\mathfrak{p}), M) = 0 \quad \text{holds for every prime ideal } \mathfrak{p} \text{ in } R,$$

then the flat dimension of M is less than n .

Proof. It suffices to prove the first claim; the assertion about the flat dimension of M is a consequence, given (4.1).

Fix \mathfrak{p} and n as in the hypotheses. Given (4.2), this yields $\text{Tor}_n^{R_{\mathfrak{p}}}(k(\mathfrak{p}), M_{\mathfrak{p}}) = 0$. One has the following (in)equalities

$$\sup H_*(M) \geq \sup H_*(M_{\mathfrak{p}}) = \sup H^* \text{Hom}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}, E(k(\mathfrak{p}))).$$

Since $\dim R \geq \dim R_{\mathfrak{p}}$, it follows from (2.7) and Proposition 3.3 that

$$\sup \text{Tor}_*^{R_{\mathfrak{p}}}(k(\mathfrak{p}), M_{\mathfrak{p}}) = \text{depth } R_{\mathfrak{p}} - \text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} < n.$$

This is the desired result. \square

The next example shows that the constraint on n in Theorem 4.1 is needed.

Example 4.2. Let (R, \mathfrak{m}, k) be a local ring, N a finitely generated Cohen–Macaulay R -module of dimension d , and set $M := H_{\mathfrak{m}}^d(N)$. There are isomorphisms

$$\text{Tor}_i^R(k, M) \cong \text{Tor}_{i-d}^R(k, N) \quad \text{for all } i.$$

To see this, let F be the Čech complex on a maximal N -regular sequence \mathbf{x} . The complex $(\Sigma^d F) \otimes_R N$ is quasi-isomorphic to M , for $\inf H_{\mathfrak{m}}^*(N) = \sup H_{\mathfrak{m}}^*(N) = d$, by the hypothesis on N . Thus, there are quasi-isomorphisms

$$k \otimes_R^{\mathbb{L}} M \simeq k \otimes_R^{\mathbb{L}} ((\Sigma^d F) \otimes_R^{\mathbb{L}} N) \simeq \Sigma^d k \otimes_R^{\mathbb{L}} N.$$

The isomorphisms above follow. Thus one has

$$\mathrm{Tor}_i^R(k, M) = \begin{cases} 0 & \text{for } i < d \\ N/\mathfrak{m}N & \text{for } i = d. \end{cases}$$

In particular, one has $\inf \mathrm{Tor}_*^R(k, M) = d$, while $\mathrm{Tor}_*^R(k(\mathfrak{p}), M) = 0$ for every prime ideal $\mathfrak{p} \neq \mathfrak{m}$, since M is \mathfrak{m} -torsion.

Now, if in addition the inequality $d > \mathrm{depth} R$ holds, then $\mathrm{flat} \dim_R M$ is infinite. To see this apply Matlis duality $\mathrm{Tor}_i^R(k, M)^\vee \cong \mathrm{Ext}_R^i(k, M^\vee)$ and conclude from Proposition 3.4 that $\mathrm{Tor}_i^R(k, M)$ is nonzero for all $i \geq d$.

It remains to remark that such R and N exist: Let k be a field, d be a positive integer, and set

$$R := k[[x_1, \dots, x_{d+1}]]/(x_1^2, x_1x_2, \dots, x_1x_{d+1}).$$

This R is a local ring of dimension d and depth 0. The R -module $N = R/(x_1)$ is Cohen–Macaulay of dimension d .

5. INJECTIVE DIMENSION

Let M be an R -complex, by [6, Theorem 1.1] one has

$$(5.1) \quad \mathrm{inj} \dim_R M = \sup\{i \in \mathbb{Z} \mid \mathrm{Ext}_R^i(k(\mathfrak{p}), M) \neq 0 \text{ for some } \mathfrak{p} \in \mathrm{Spec} R\}.$$

In view of (4.2) this is a perfect parallel to the formula for flat dimension (4.1).

The equality $\mathrm{flat} \dim_R M = \sup_{\mathfrak{p} \in \mathrm{Spec} R} \{\mathrm{flat} \dim_{R_{\mathfrak{p}}} M_{\mathfrak{p}}\}$ is immediate from (4.1) and (4.2). The corresponding equality for the injective dimension only holds true under extra conditions, and the whole picture is altogether more complicated. If M satisfies $\inf H^*(M) > -\infty$, then [2, Proposition 5.3.I] yields

$$(5.2) \quad \begin{aligned} \mathrm{inj} \dim_R M &= \sup\{i \in \mathbb{Z} \mid \mathrm{Ext}_{R_{\mathfrak{p}}}^i(k(\mathfrak{p}), M_{\mathfrak{p}}) \neq 0 \text{ for some } \mathfrak{p} \in \mathrm{Spec} R\} \\ &= \sup_{\mathfrak{p} \in \mathrm{Spec} R} \{\mathrm{inj} \dim_{R_{\mathfrak{p}}} M_{\mathfrak{p}}\}. \end{aligned}$$

Without the boundedness condition on $H(M)$ the injective dimension may increase under localization; an example is provided in 6.3.

The next statement, which still requires homological boundedness, is folklore but not readily available in the literature.

Theorem 5.1. *Let R be a commutative noetherian ring and M an R -complex with $\inf H^*(M) > -\infty$. If there exists an integer $n \geq \sup H^*(M)$ such that*

$$\mathrm{Ext}_{R_{\mathfrak{p}}}^{n+1}(k(\mathfrak{p}), M_{\mathfrak{p}}) = 0 \quad \text{holds for every prime ideal } \mathfrak{p} \text{ in } R,$$

then the injective dimension of M is at most n .

Proof. Let I be a minimal semi-injective resolution of M ; as $\inf H^*(M) > -\infty$ holds one has $I^n = 0$ for $n \ll 0$. For every integer i one has $I^i = \coprod_{\mathfrak{p} \in \mathrm{Spec} R} E(R/\mathfrak{p})^{(\mu_i(\mathfrak{p}))}$, and to prove that $\mathrm{inj} \dim_R M$ is at most n it is sufficient to show that the index set $\mu_{n+1}(\mathfrak{p})$ is empty for every prime \mathfrak{p} . Fix \mathfrak{p} . Since $I_{\mathfrak{p}}$ is a complex of injectives with

$(I_{\mathfrak{p}})^n = 0$ for $n \ll 0$, it is a minimal semi-injective resolution of $M_{\mathfrak{p}}$, so one has

$$\begin{aligned} 0 &= \text{Ext}_{R_{\mathfrak{p}}}^{n+1}(k(\mathfrak{p}), M_{\mathfrak{p}}) \\ &= \text{H}^{n+1} \text{Hom}_{R_{\mathfrak{p}}}(k(\mathfrak{p}), I_{\mathfrak{p}}) \\ &= \text{Hom}_{R_{\mathfrak{p}}}(k(\mathfrak{p}), (I_{\mathfrak{p}})^{n+1}) \\ &= \text{Hom}_{R_{\mathfrak{p}}}(k(\mathfrak{p}), E(k(\mathfrak{p}))^{(\mu_{n+1}(\mathfrak{p}))}). \end{aligned}$$

It follows that $\mu_{n+1}(\mathfrak{p})$ is empty. \square

The next result corresponds to (4.3). It removes a restriction on the boundedness of M in Yassemi's version [17, Theorem 2.10] of Chouinard's formula [5, Corollary 3].

Proposition 5.2. *For every R -complex M of finite injective dimension one has*

$$\text{inj dim}_R M = \sup_{\mathfrak{p} \in \text{Spec } R} \{ \text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \}.$$

Proof. Without loss of generality, we can assume that M is semi-injective with $M^i = 0$ for all $i > d := \text{inj dim}_R M$. For every $u \leq d$ there is an exact sequence of semi-injective complexes

$$(1) \quad 0 \longrightarrow M^{\geq u} \longrightarrow M \longrightarrow M^{\leq u-1} \longrightarrow 0$$

with $\text{inj dim}_R M^{\leq u-1} \leq u-1$ and $\text{inj dim}_R M^{\geq u} = d$. The complex $M^{\geq u}$ is bounded, so (2.4) and (5.2) conspire to yield

$$(2) \quad d = \sup_{\mathfrak{p} \in \text{Spec } R} \{ \text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}^{\geq u} \}.$$

First we establish the inequality

$$(3) \quad \text{inj dim}_R M \geq \sup_{\mathfrak{p} \in \text{Spec } R} \{ \text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \}.$$

Let \mathfrak{p} be a prime. There are inequalities

$$\text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \geq \inf \text{H}_*(M_{\mathfrak{p}}) \geq \inf \text{H}_*(M) = -\sup \text{H}^*(M) \geq -d,$$

and without loss of generality one can assume that $\text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}$ is finite. Consider (1) for $u = -\text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}$ and localize at \mathfrak{p} . The associated exact sequence of Tor groups yields $\text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = \text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}^{\geq u}$, so the desired inequality $d \geq \text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}$ follows from (2). It remains to prove that equality holds for some prime.

Consider (1) for $u = d-1$ and choose by (2) a prime \mathfrak{p} with

$$d = \text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}^{\geq d-1}.$$

The second inequality in the next display is (3) applied to the complex $M^{\leq d-2}$.

$$d-2 \geq \text{inj dim}_R M^{\leq d-2} \geq \text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}^{\leq d-2}.$$

Eliminating d and $\text{depth } R_{\mathfrak{p}}$ between the two displays one gets the inequality

$$\text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}^{\geq d-1} \leq \text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}^{\leq d-2} - 2.$$

Finally, one gets $\text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = \text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}^{\geq d-1}$ from the exact sequence of Tor groups associated to (1). \square

We now aim for a characterization of complexes of finite injective dimension that does not require homological boundedness. It is based on the following observation, of independent interest.

Lemma 5.3. *Let M be an R -complex and \mathfrak{m} a maximal ideal in R . The localization maps $\rho: M \rightarrow M_{\mathfrak{m}}$ and $\sigma: R \rightarrow R_{\mathfrak{m}}$ induce quasi-isomorphisms*

$$\begin{aligned} \mathrm{RHom}_R(k(\mathfrak{m}), \rho): \mathrm{RHom}_R(k(\mathfrak{m}), M) &\xrightarrow{\simeq} \mathrm{RHom}_R(k(\mathfrak{m}), M_{\mathfrak{m}}) \quad \text{and} \\ k(\mathfrak{m}) \otimes_R^{\mathbb{L}} \mathrm{RHom}_R(\sigma, M): k(\mathfrak{m}) \otimes_R^{\mathbb{L}} \mathrm{RHom}_R(R_{\mathfrak{m}}, M) &\xrightarrow{\simeq} k(\mathfrak{m}) \otimes_R^{\mathbb{L}} M. \end{aligned}$$

Proof. In the derived category of R , consider the distinguished triangle

$$M \xrightarrow{\rho} M_{\mathfrak{m}} \rightarrow C \rightarrow \Sigma M.$$

The induced morphism $k(\mathfrak{m}) \otimes_R^{\mathbb{L}} \rho$ is a quasi-isomorphism, so $k(\mathfrak{m}) \otimes_R^{\mathbb{L}} C$ is acyclic. Then $\mathrm{RHom}_R(k(\mathfrak{m}), C)$ is also acyclic, by [3, Theorem 4.13], whence the map $\mathrm{RHom}_R(k(\mathfrak{m}), \rho)$ is a quasi-isomorphism, as claimed.

In the same vein, the distinguished triangle $R \rightarrow R_{\mathfrak{m}} \rightarrow D \rightarrow \Sigma R$ induces a distinguished triangle

$$\mathrm{RHom}_R(D, M) \rightarrow \mathrm{RHom}_R(R_{\mathfrak{m}}, M) \xrightarrow{\mathrm{RHom}_R(\sigma, M)} M \rightarrow \Sigma \mathrm{RHom}_R(D, M).$$

By adjunction $\mathrm{RHom}_R(k(\mathfrak{m}), \mathrm{RHom}_R(\sigma, M))$ is a quasi-isomorphism, so that

$$\mathrm{RHom}_R(k(\mathfrak{m}), \mathrm{RHom}_R(D, M))$$

is acyclic. Thus the complex $k(\mathfrak{m}) \otimes_R^{\mathbb{L}} \mathrm{RHom}_R(D, M)$ is acyclic by [3, Theorem 4.13], which justifies the second of the desired quasi-isomorphisms. \square

Proposition 5.4. *Let M be an R -complex and \mathfrak{m} a maximal ideal in R . If one has $\mathrm{Ext}_R^n(k(\mathfrak{m}), M) = 0$ for some $n \geq \dim R_{\mathfrak{m}} + \sup H^*(M_{\mathfrak{m}})$, then $\mathrm{Ext}_R^i(k(\mathfrak{m}), M) = 0$ holds for all $i \geq n$ and there are equalities*

$$\begin{aligned} \sup \mathrm{Ext}_R^*(k(\mathfrak{m}), M) &= \mathrm{depth} R_{\mathfrak{m}} - \mathrm{width}_{R_{\mathfrak{m}}} M_{\mathfrak{m}} \\ &= \mathrm{depth} R_{\mathfrak{m}} - \mathrm{width}_{R_{\mathfrak{m}}} \mathrm{RHom}_R(R_{\mathfrak{m}}, M). \end{aligned}$$

Proof. The first isomorphism below is by Lemma 5.3; the second is by adjunction.

$$\mathrm{Ext}_R^*(k(\mathfrak{m}), M) \cong \mathrm{Ext}_R^*(k(\mathfrak{m}), M_{\mathfrak{m}}) \cong \mathrm{Ext}_{R_{\mathfrak{m}}}^*(k(\mathfrak{m}), M_{\mathfrak{m}}).$$

In view of these isomorphisms and the assumption on n , Proposition 3.2 now yields

$$\sup \mathrm{Ext}_R^*(k(\mathfrak{m}), M) = \mathrm{depth} R_{\mathfrak{m}} - \mathrm{width}_{R_{\mathfrak{m}}} M_{\mathfrak{m}} < n.$$

It remains to observe that the width of $M_{\mathfrak{m}}$ and $\mathrm{RHom}_R(R_{\mathfrak{m}}, M)$ coincide, by the second quasi-isomorphism in Lemma 5.3. \square

Corollary 5.5. *Let R be an artinian ring and M an R -complex. If there exists an integer $n \geq \sup H^*(M)$ such that*

$$\mathrm{Ext}_R^n(k(\mathfrak{p}), M) = 0 \quad \text{holds for every prime ideal } \mathfrak{p} \text{ in } R,$$

then the injective dimension of M is less than n .

Proof. Every prime ideal \mathfrak{p} in R is maximal and there are inequalities

$$n \geq \sup H^*(M) \geq \sup H^*(M_{\mathfrak{p}}) = \dim R_{\mathfrak{p}} + \sup H^*(M_{\mathfrak{p}}).$$

Thus the claim follows from (5.1) and Proposition 5.4. \square

Remark 5.6. In the sequel we require the invariant

$$\text{splf } R = \sup\{\text{proj dim}_R F \mid F \text{ is a flat } R\text{-module}\}.$$

Every flat R -module is projective if and only if R is artinian, so $\text{splf } R > 0$ holds if $\dim R > 0$, and from work of Jensen [11, Proposition 6] and Raynaud and Gruson [15, thm. II.3.2.6] one gets the upper bound $\text{splf } R \leq \dim R$. A result of Gruson and Jensen [10, Theorem 7.10] yields another bound on the invariant $\text{splf } R$: If R has cardinality at most \aleph_m for some natural number m , then one has $\text{splf } R \leq m + 1$. Thus for countable rings, and for 1-dimensional rings, the bound in the theorem below is $n \geq \dim R + \sup H^*(M)$.

Theorem 5.7. *Let R be a commutative noetherian ring with $\dim R \geq 1$ and M an R -complex. If there exists an integer $n \geq \dim R - 1 + \text{splf } R + \sup H^*(M)$ such that*

$$\text{Ext}_R^n(k(\mathfrak{p}), M) = 0 \quad \text{holds for every prime ideal } \mathfrak{p} \text{ in } R,$$

then the injective dimension of M is less than n .

Proof. Fix a prime ideal \mathfrak{p} and consider the $R_{\mathfrak{p}}$ -complex $N := \text{RHom}_R(R_{\mathfrak{p}}, M)$. One has

$$H^i(N) = \text{Ext}_R^i(R_{\mathfrak{p}}, M) = 0 \quad \text{for } i > \text{proj dim}_R R_{\mathfrak{p}} + \sup H^*(M),$$

and standard adjunction yields

$$\text{Ext}_R^*(k(\mathfrak{p}), M) \cong \text{Ext}_{R_{\mathfrak{p}}}^*(k(\mathfrak{p}), N).$$

Since $R_{\mathfrak{p}}$ is a flat R -module, $\text{proj dim}_R R_{\mathfrak{p}}$ is at most $\text{splf } R$. Given this, (2.7) yields

$$H_{\mathfrak{p}R_{\mathfrak{p}}}^i(N) = 0 \quad \text{for } i > \dim R_{\mathfrak{p}} + \text{splf } R + \sup H^*(M).$$

Thus, if $n \geq \dim R_{\mathfrak{p}} + \text{splf } R + \sup H^*(M)$ holds, then Proposition 3.2 yields

$$\sup \text{Ext}_R^*(k(\mathfrak{p}), M) = \text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} N < n.$$

The same equality also holds when $n < \dim R_{\mathfrak{p}} + \text{splf } R + \sup H^*(M)$, for then the assumption on n forces $\dim R_{\mathfrak{p}} = \dim R$, so \mathfrak{p} is a maximal ideal and so Proposition 5.4 applies. Now the desired conclusions follows from (5.1). \square

Remark 5.8. As noted in Remark 5.6, one has $\text{splf } R \leq \dim R$. Thus Theorem 1.3 is a consequence of Corollary 5.5 and Theorem 5.7.

Confer the following result and Proposition 5.2.

Corollary 5.9. *For every R -complex of finite injective dimension one has*

$$\text{inj dim}_R M = \sup_{\mathfrak{p} \in \text{Spec } R} \{\text{depth } R_{\mathfrak{p}} - \text{width}_{R_{\mathfrak{p}}} \text{RHom}_R(R_{\mathfrak{p}}, M)\}.$$

Proof. Given Lemma 5.3, the desired equality is restatement of Proposition 5.2 in case R is artinian. If R is not artinian, then one has $\dim R \geq 1$ and the equality is immediate from (5.1) and the last display in the proof of Theorem 5.7. \square

Remark 5.10. Let R be a complete local domain of positive dimension. One has $\text{width}_{R_{(0)}} R_{(0)} = 0$, but the complex $\text{RHom}_R(R_{(0)}, R)$ is acyclic, see [3, Example 4.20], so $\text{width}_{R_{(0)}} \text{RHom}_R(R_{(0)}, R) = \infty$. We do not know how the numbers $\text{width}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}$ and $\text{width}_{R_{\mathfrak{p}}} \text{RHom}_R(R_{\mathfrak{p}}, M)$ from Proposition 5.2 and Corollary 5.9 compare in general.

6. EXAMPLES

In this section we describe examples to illustrate that, for complexes whose cohomology is not bounded below, finiteness of injective dimension does not behave well under localization or passage to torsion subcomplexes. This builds on [4].

Remark 6.1. Let R be a ring. A complex I of injective R -modules is semi-injective if and only if for each (equivalently, for some) integer n the quotient complex $I^{\leq n}$ is semi-injective. This is immediate from the exact sequence of complexes

$$0 \longrightarrow I^{>n} \longrightarrow I \longrightarrow I^{\leq n} \longrightarrow 0$$

since $I^{>n}$ is always semi-injective.

Remark 6.2. Let (R, \mathfrak{m}, k) be a local ring and E be the injective envelope of k . One has $(E^{\mathbb{N}})_{\mathfrak{p}} \neq 0$ for every prime ideal in R . Indeed, the claim is trivial if R is artinian. If R is not artinian, then one can choose an element $e = (e_n)_{n \in \mathbb{N}}$ in $E^{\mathbb{N}}$ with $\mathfrak{m}^n \subseteq (0 : e_n) \not\subseteq \mathfrak{m}^{n-1}$. The map $R \rightarrow E^{\mathbb{N}}$ given by $1 \mapsto e$ is injective by Krull's intersection theorem, so R is a submodule of $E^{\mathbb{N}}$.

Example 6.3. Let (R, \mathfrak{m}, k) be a local ring such that $(0 : x) = (x)$ holds for some $x \in \mathfrak{m}$; set $S = R/(x)$. The complex

$$\cdots \xrightarrow{x} R \xrightarrow{x} R \xrightarrow{x} R \longrightarrow 0$$

concentrated in nonnegative degrees has homology S in degree 0 and zero elsewhere. Dualizing with respect to E , the injective envelope of k over R , yields a complex

$$I := 0 \longrightarrow E \xrightarrow{x} E \xrightarrow{x} E \xrightarrow{x} \cdots$$

of injective R -modules. It is the minimal injective resolution of $E_S := \text{Hom}_R(S, E)$ over R . By periodicity, every injective syzygy of E_S is E_S . Consider the complex $J = \prod_{n>0} \Sigma^n I$, which is a semi-injective resolution of $\prod_{n>0} \Sigma^n E_S$.

Claim. The complex $M := J^{\leq 0}$ has injective dimension 0, whereas for each prime ideal $\mathfrak{p} \neq \mathfrak{m}$, one has that $\text{inj dim}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}$ is infinite.

Indeed, since J is semi-injective, so is M , by Remark 6.1. Since the cohomology module $H^0(M) \cong (E_S)^{\mathbb{N}}$ is nonzero, it follows that $\text{inj dim}_R M = 0$ holds.

Fix a prime ideal $\mathfrak{p} \neq \mathfrak{m}$. For $i < 0$ one has $H^i(M) = H^i(J) = E_S$ and, therefore, $H^i(M_{\mathfrak{p}}) = H^i(M)_{\mathfrak{p}} = 0$. This justifies the first quasi-isomorphism in the computation below; the rest are standard.

$$\begin{aligned} \text{RHom}_{R_{\mathfrak{p}}}(k(\mathfrak{p}), M_{\mathfrak{p}}) &\simeq \text{RHom}_{R_{\mathfrak{p}}}(k(\mathfrak{p}), ((E_S)^{\mathbb{N}})_{\mathfrak{p}}) \\ &\simeq \text{RHom}_R(R/\mathfrak{p}, (E_S)^{\mathbb{N}})_{\mathfrak{p}} \\ &\simeq (\text{RHom}_R(R/\mathfrak{p}, E_S)^{\mathbb{N}})_{\mathfrak{p}} \\ &\simeq (\text{Hom}_R(R/\mathfrak{p}, I)^{\mathbb{N}})_{\mathfrak{p}} \\ &\cong \left(\left(\prod_{i \geq 0} \Sigma^{-i} \text{Hom}_R(R/\mathfrak{p}, E) \right)^{\mathbb{N}} \right)_{\mathfrak{p}} \\ &\cong \prod_{i \geq 0} \Sigma^{-i} (\text{Hom}_R(R/\mathfrak{p}, E)^{\mathbb{N}})_{\mathfrak{p}}. \end{aligned}$$

The first isomorphism holds because $x = 0$ in R/\mathfrak{p} , for $x^2 = 0$, and hence the induced differential on the complex $\text{Hom}_R(R/\mathfrak{p}, I)$ is zero.

The module $E_{R/\mathfrak{p}} := \text{Hom}_R(R/\mathfrak{p}, E)$ is the injective envelope of k over the domain R/\mathfrak{p} . The computation above shows that for every $i \geq 0$ there is an isomorphism as R/\mathfrak{p} -modules

$$\text{Ext}_{R/\mathfrak{p}}^i(k(\mathfrak{p}), M_{\mathfrak{p}}) \cong ((E_{R/\mathfrak{p}})^{\mathbb{N}})_{(0)}.$$

Thus Remark 6.2 yields $\text{Ext}_{R/\mathfrak{p}}^i(k(\mathfrak{p}), M_{\mathfrak{p}}) \neq 0$ for all $i \geq 0$; hence $\text{inj dim}_{R/\mathfrak{p}} M_{\mathfrak{p}}$ is infinite, as claimed.

Example 6.4. Let k be a field and $R := k[[x, y]]/(x^2)$. Since $(0 : x) = (x)$, we are in the situation considered in the previous example. Let M be the complex of injectives with injective dimension zero constructed there. We claim that the injective dimension of the complexes M_y and $\Gamma_{(y)}M$ are infinite.

Indeed, observe that $M_y \cong M_{\mathfrak{p}}$ where \mathfrak{p} is the prime ideal (x) of R , so $\text{inj dim}_R M_y$ is infinite, by the claim in the previous example. Since $C(y) \otimes_R M$ is quasi-isomorphic to $\Gamma_{(y)}M$ and there is an exact sequence

$$0 \longrightarrow \Sigma^{-1}M_y \longrightarrow C(y) \otimes_R M \longrightarrow M \longrightarrow 0,$$

it follows that the injective dimension of $\Gamma_{(y)}M$ is infinite as well.

ACKNOWLEDGMENTS

We thank Peder Thompson for conversations that helped clarify the material in Section 5.

REFERENCES

- [1] Leovigildo Alonso Tarrío, Ana Jeremías López, and Joseph Lipman, *Local homology and cohomology on schemes*, Ann. Sci. École Norm. Sup. (4) **30** (1997), no. 1, 1–39. MR1422312
- [2] Luchezar L. Avramov and Hans-Bjørn Foxby, *Homological dimensions of unbounded complexes*, J. Pure Appl. Algebra **71** (1991), no. 2-3, 129–155. MR1117631
- [3] David J. Benson, Srikanth B. Iyengar, and Henning Krause, *Colocalizing subcategories and cosupport*, J. Reine Angew. Math. **673** (2012), 161–207. MR2999131
- [4] Xiao-Wu Chen and Srikanth B. Iyengar, *Support and injective resolutions of complexes over commutative rings*, Homology, Homotopy Appl. **12** (2010), no. 1, 39–44. MR2594681
- [5] Leo G. Chouinard, II, *On finite weak and injective dimension*, Proc. Amer. Math. Soc. **60** (1976), 57–60 (1977). MR0417158
- [6] Lars Winther Christensen and Srikanth B. Iyengar, *Tests for injectivity of modules over commutative rings*, Collect. Math. **68** (2017), 243–250.
- [7] William G. Dwyer and John P. C. Greenlees, *Complete modules and torsion modules*, Amer. J. Math., **124** (2002) 199–220.
- [8] Robert Fossum, Hans-Bjørn Foxby, Phillip Griffith, and Idun Reiten, *Minimal injective resolutions with applications to dualizing modules and Gorenstein modules*, Inst. Hautes Études Sci. Publ. Math. (1975), no. 45, 193–215.
- [9] Hans-Bjørn Foxby and Srikanth Iyengar, *Depth and amplitude for unbounded complexes*, Commutative algebra (Grenoble/Lyon, 2001) (Providence, RI), Contemp. Math., vol. 331, Amer. Math. Soc., 2003, pp. 119–137. MR2013162
- [10] Laurent Gruson and Christian U. Jensen, *Dimensions cohomologiques reliées aux foncteurs $\varprojlim^{(i)}$* , Paul Dubreil and Marie-Paule Malliavin Algebra Seminar, 33rd Year (Paris, 1980), Lecture Notes in Math., vol. 867, Springer, Berlin, 1981, pp. 234–294. MR0633523
- [11] Christian U. Jensen, *On the vanishing of $\varprojlim^{(i)}$* , J. Algebra **15** (1970), 151–166. MR0260839
- [12] T. Y. Lam and Manuel L. Reyes, *A prime ideal principle in commutative algebra*, J. Algebra **319** (2008), 3006–3027. MR2397420
- [13] Joseph Lipman, *Lectures on local cohomology and duality*, Local cohomology and its applications (Guanajuato, 1999), Lecture Notes in Pure and Appl. Math., vol. 226, Dekker, New York, 2002, pp. 39–89. MR1888195

- [14] Barbara L. Osofsky, *Homological dimension and the continuum hypothesis*, Trans. Amer. Math. Soc. **132** (1968), 217–230. MR0224606
- [15] Michel Raynaud and Laurent Gruson, *Critères de platitude et de projectivité. Techniques de “platification” d’un module*, Invent. Math. **13** (1971), 1–89. MR0308104
- [16] Anne-Marie Simon, *Some homological properties of complete modules*, Math. Proc. Cambridge Philos. Soc. **108** (1990), no. 2, 231–246. MR1074711
- [17] Siamak Yassemi, *Width of complexes of modules*, Acta Math. Vietnam. **23** (1998), no. 1, 161–169. MR1628029

L.W.C. TEXAS TECH UNIVERSITY, LUBBOCK, TX 79409, U.S.A.

E-mail address: `lars.w.christensen@ttu.edu`

URL: `http://www.math.ttu.edu/~lchrste`

S.B.I. UNIVERSITY OF UTAH, SALT LAKE CITY, UT 84112, U.S.A.

E-mail address: `iyengar@math.utah.edu`

URL: `http://www.math.utah.edu/~iyengar`

T.M. UNIVERSITY OF NEBRASKA-LINCOLN, LINCOLN, NE 68588, U.S.A.

E-mail address: `tmarley1@unl.edu`

URL: `http://www.math.unl.edu/~tmarley`